


Bijan Nekoueishahraki

Synthesis, Structure and Reactivity of Bismuth(III) and Aluminum(III) Complexes Supported by Nitrogen Donor Ligands



Göttingen 2009

Synthesis, Structure and Reactivity of Bismuth(III) and Aluminum(III) Complexes Supported by Nitrogen Donor Ligands

Dissertation
zur Erlangung des Doktorgrades
der Mathematisch-Naturwissenschaftlichen Fakultäten
der Georg-August-Universität zu Göttingen

vorgelegt von

Bijan Nekoueishahraki
aus Shahrekord
(IRAN)

Göttingen 2009

D 7

Referent: Prof. Dr. Dr. h.c. mult. Herbert W. Roesky

Koreferent: Prof. Dr. Dietmar Stalke

Tag der mündlichen Prüfung:

Dedicated to my mother and in the memory of my father

for their love and affection

Acknowledgment

The work described in this doctoral dissertation has been carried out under the guidance and supervision of Professor Dr. Dr. h. c. mult. H. W. Roesky at the Institut für Anorganische Chemie der Georg-August-Universität in Göttingen between January 2006 and January 2010.

I would like to express my deepest gratitude to my supervisor

Professor Dr. Dr. h. c. mult. H. W. Roesky

for his constant guidance, motivation, suggestions, and discussion throughout this work.

I thank Mr. D. Stern, Prof. D. Stalke, Dr. G. Schwab, and Prof. C. Schulzke for their help in the X-ray crystal structural investigations and refinement of disordered molecules and twinned crystals. I thank Mr. W. Zolke, Mr. R. Schöne, Dr. M. John (NMR spectra), Dr. D. Böhler, Mr. T. Schuchardt, Mr. J. Schöne (mass spectra), Mr. M. Hesse (IR spectra), Mr. J. Schimkowiak, Mr. M. Schlote, and the staff of the Analytical Laboratories, and Werkstatt for their timely support during this research work. I am also thankful to all the members of glass blowing section, work shop, chemical store as well the security of our institute for their co-operation and help in all kind of situations.

I am grateful to Dr. A. Jana, Mr. S. S. Sen, Mr. Z. Zhang, Mr. A. Döring, Dr. S. Nembenna, Dr. R. S. Ghadwal, Dr. S. K. Mandal, Dr. Y. Ying, and Dr. V. M. Jimenez-Perez.

My special thanks to S. P. Sarish, P. P. Samuel, and P. M. Guru, for their friendliness and help during my stay at Göttingen.

The full support and encouragement from my parents, family and friends made this work possible.

Table of Contents

1. Introduction	1
1.1. Organobismuth(III) Compounds	1
1.2. Organoaluminum(III) Compounds	7
1.3. Aims and Objectives of the Present Work	11
2. Results and Discussion	12
2.1. Bismuthamidation of Unsaturated Organic Compounds	12
2.1.1. Preparation of 1,8-$C_{10}H_6(NSiMe_3)_2BiNMe_2$ (1)	13
2.1.2. Reaction of L^1BiNMe_2 (1) with Ketone and Aldehyde: Synthesis of $L^1BiOCPh(2-Py)(NMe_2)$ (2) and $L^1BiOCH(3-Py)(NMe_2)$ (3)	14
2.1.3. Reaction of L^1BiNMe_2 (1) with Alkene and Alkyne: Synthesis of $L^1BiC(Me)(CN)CH_2NMe_2$ (4) and $L^1BiC(CO_2Et)CNMe_2(CONMe_2)$ (5)	18
2.1.4. Reaction of L^1BiNMe_2 (1) with <i>N,N'</i>-Diisopropylcarbodiimide: Synthesis of 1,8-$C_{10}H_6(NSiMe_3)_2Bi(NiPr)_2CNMe_2$ (6)	21
2.1.5. Reaction of L^1BiNMe_2 (1) with Phenylisocyanate: Synthesis of 1,8-$C_{10}H_6(NSiMe_3)_2BiNPhCONMe_2$ (7)	23
2.2. Heterobimetallic Bismuth Complexes with Main Group and Transition Metals	25
2.2.1. Reaction of L^1BiNMe_2 (1) with $LM(Me)(OH)$: Synthesis of 1,8-$C_{10}H_6(NSiMe_3)_2Bi(\mu-O)MMeL$ ($M = Al$ 8; Ga, 9)	26
2.2.2. Reaction of L^1BiNMe_2 (1) with $LGe(OH)$: Synthesis of 1,8-$C_{10}H_6(NSiMe_3)_2Bi(\mu-O)GeL$ (10)	29
2.2.3. Reaction of L^1BiNMe_2 (1) with $Cp^*_2MR(OH)$: Synthesis of 1,8-$C_{10}H_6(NSiMe_3)_2Bi(\mu-O)MRCp^*_2$ ($M = Zr$ 11, $R = Me$; Hf, 12, $R = Cl$)	31
2.2.4. Structural Comparison of compounds 8-12	35
2.3. Synthesis, Structure and Reactivity of Organobismuth(III), Bismuth Chalcogenide and Dibismuthine Complexes	35
2.3.1. Synthesis, Spectroscopic Characterization and Structure of Complexes [1,8-$C_{10}H_6(NSiMe_3)_2]BiR$ ($R = Cl$(13), $CCPh$(14), Cp^*(15), Me(16), OPh(17),	

NH_2 (18).....	36
2.3.2. <i>Molecular Structure Description of Compounds 13-16</i>	37
2.3.3. <i>Synthesis, Spectroscopic Characterization and Structure of Organobismuth Chalcogenide Complex: Synthesis of [1,8-$C_{10}H_6(NSiMe_3)_2Bi]_2S$ (19)</i>	40
2.3.4. <i>Synthesis, Spectroscopic Characterization and Structure of Low-valent Dibismuthine Complex: Synthesis of [1,8-$C_{10}H_6(NSiMe_3)_2Bi]_2$ (20)</i>	42
2.4. <i>Synthesis, Characterization, and X-ray Single Crystal Structures of N,N'-Bis(trimethylsilyl)diphenyliminophosphonamide Aluminum Halide, Hydride, Amide and Alkylate</i>	45
2.4.1. <i>Synthesis of $Ph_2P(NSiMe_3)_2Li$ (22)</i>	45
2.4.2. <i>Reaction of L^2Li with $RAiCl_2$: Synthesis of $Ph_2P(NSiMe_3)_2AlRCl$ ($R = Cl$, 23; Me, 24)</i>	46
2.4.3. <i>Reaction of L^2H with $AlH_3 \cdot NMe_3$: Synthesis of $[Ph_2P(NSiMe_3)_2]_2AlH$ (25)</i>	50
2.4.4. <i>Reaction of L^2H with $AlEt_3$ and $[Al(NMe_2)_3]_2$: Synthesis of $Ph_2P(NSiMe_3)_2AlEt_2$ (26) and $Ph_2P(NSiMe_3)_2Al(NMe_2)_2$ (27)</i>	52
2.4.5. <i>Structural Comparison of L^2AlCl_2 (23), $L^2AlMeCl$ (24), L^2_2AlH (25), and $L^2Al(NMe_2)_2$ (27)</i>	54
3. Summary and Outlook	55
3.1. <i>Summary</i>	55
3.2. <i>Outlook</i>	61
4. Experimental Section	62
4.1. <i>General Procedures</i>	62
4.2. <i>Physical Measurements</i>	62
4.3. <i>Starting Materials</i>	63
4.4. <i>Synthesis</i>	63
5. Handling and Disposal of Solvents and Residual Waste	75

6. References.....	76
List of Publications.....	87

Abbreviations

av	Average
Ar	Aryl
br	Broad
<i>t</i> Bu	<i>tert</i> -butyl
C	Celsius
calcd	calculated
Cp	cyclopentadienyl
Cp*	pentamethylcyclopentadienyl
d	doublet
dec	decomposition
δ	chemical shift
EI	electron impact ionization
g	grams
<i>i</i> Pr	isopropyl
IR	infrared
h	hour(s)
η	hapto
Hz	Hertz
λ	wavelength
M ⁺	molecular ion
<i>m/z</i>	mass/charge
M.p.	melting point
μ	bridging
NMR	nuclear magnetic resonance
ppm	parts per million

q	quartet
RT	room temperature
s	singlet
sept	septet
t	triplet
THF	tetrahydrofuran
TMS	tetramethylsilane
V	Volume
$\tilde{\nu}$	Wave number
Z	number of molecules in the unit cell
L	$\text{CH}((\text{CMe})\text{NAr})_2$, Ar = 2,6- <i>i</i> Pr ₂ C ₆ H ₃
L ¹	1,8-C ₁₀ H ₆ (NSiMe ₃) ₂
L ²	Ph ₂ P(NSiMe ₃) ₂

1. Introduction

This section of the thesis gives the background and an overview of the area in two sections before the present work is put forward.


1.1. Organobismuth(III) Compounds

Bismuth was first mentioned in the 1450s as Wismutton or Bisemutun, probably derived from the old German word Weissmuth or Wismut, meaning white substances. Bismuth, Bi, the 83rd element of the periodic table is the most metallic and the least abundant of the elements in the nitrogen family (group 15). The bismuth atom with a ground state electronic configuration of $[\text{Xe}] 4f^{14}5d^{10}6s^26p^3$ usually utilizes the three $6p$ electrons in bond formation and retains the two $6s$ electrons as an inert pair, hence the oxidation state +3 exhibited bismuth in the vast majority of its compounds. However, a variety of organobismuth compounds can contain the element in the +5 oxidation state. Coordination numbers are 2, 3, 4, 5 and 6. It is the heaviest member of group 15 and the heaviest stable element in the periodic table. Although it belongs to the group 15 family, the chemistry of bismuth differs considerably from that of other lighter members, such as phosphorus, arsenic and antimony. Elemental bismuth is inert in dry air at room temperature, but oxidizes slowly to become covered with a thin film of the oxide. Bismuth compounds are attracting increasing attention due to their application in heterogeneous catalysis,¹ catalysts for organic synthesis,² superconducting materials,³ and also biological activity.

Bismuth amides, $\text{Bi}(\text{NRR}')_3$, are a series of compounds that have covalent Bi-N bonds (R and R' can be alkyl, aryl or silyl groups). These compounds can be promising precursors for bismuth oxide based materials^{4,5} and have been used as precursors for metal-organic chemical vapor deposition (MOCVD) of bismuth-containing thin films due to their low M-N bond strength.⁶ Relatively few bismuth amides with sterically crowded substituents are known. The

reported bismuth amides were synthesized by reaction of the lithium amide salt with a bismuth halide (Scheme 1).


Scheme 1



R and R' = alkyl, aryl or silyl R'' = alkyl n = 0, 1, or 2


Depending on the different R and R' groups, bismuth amides $\text{Bi(NRR}')_3$ have varying properties. When R and R' are alkyl groups (R = R' = Me, Et, or Pr), bismuth amides are air sensitive and thermosensitive.⁷ In the preparation of these simple alkyl bismuth amides, the use of THF as solvent is indispensable; all attempts in hexane, cyclohexane, petroleum ether and ether failed. These amides gradually turn jet black at room temperature; however, they can be stored indefinitely at dry ice temperature. All of these bismuth amides are volatile. The best conditions for sublimation of $\text{Bi(NMe}_2)_3$ are using a pressure of about 10^{-2} mbar and warming the flask to 30 °C. $\text{Bi(NMe}_2)_3$ is light sensitive and with exposure to bright sunlight it will turn black. For R = aryl, both the compounds with simple phenyl group and substituted phenyl groups have been reported. When R = R' = Ph, the bismuth amide is soluble in toluene, THF and CH_2Cl_2 and is very sensitive to traces of moisture. When Li(NRH) was used (R = 2,6- $\text{Me}_2\text{C}_6\text{H}_3$), a trinuclear complex $[\text{Bi}_3(\mu\text{-NR})_4(\text{NHR})]$ was characterized by single crystal X-ray diffraction.⁸ A pure sample of this compound could not be obtained due to contamination by other unidentified complexes. When the steric bulk was increased by introducing ortho isopropyl substituents on the phenyl groups, a bismuth amide with a cyclic dimeric structure was obtained (Scheme 2).⁹

Scheme 2



When $\text{Li}(\text{NHR})$ ($\text{R} = 2,4,6\text{-tri-}i\text{-tert-butylphenyl}$) was used, the steric bulk was further increased and monomeric $\text{Bi}[\text{N}(\text{H})(\text{C}_6\text{H}_2t\text{Bu}_3)]_3$ was obtained.¹⁰ Burford et al. pointed out that the monomeric compounds $\text{Bi}[\text{N}(\text{H})(\text{C}_6\text{H}_2t\text{Bu}_3)]_3$ and $\text{Bi}[\text{N}(\text{H})(\text{C}_6\text{H}_3i\text{Pr}_2)]_3$ should form first, and then the corresponding dimers will form.¹⁰ They suggested a different mechanism (Scheme 3) to account for the formation of the dimer $[\text{RNBi}(\text{NHR})]_2$, $\text{R} = 2,6\text{-diisopropylphenyl}$, through the bismazene intermediate **A**. James et al. also believe that their trinuclear complex $[\text{Bi}_3(\mu\text{-NR})_4(\text{NHR})]$ is formed from the corresponding tricoordinate compound by elimination of the amine.⁸


Scheme 3



Bismuth silylamides, $\text{Bi}(\text{NRR}')_3$ ($\text{R} = \text{SiMe}_3$), have also been reported. A series of monosilylated bismuth amides ($\text{Me}_2\text{Bi}[\text{NMe}(\text{SiMe}_3)]$, $\text{MeBi}[\text{NMe}(\text{SiMe}_3)]_2$, and $\text{Bi}[\text{NMe}(\text{SiMe}_3)]_3$) were first reported by Scherer et al.¹¹ The corresponding disilylated

derivative, $\text{Bi}[\text{N}(\text{SiMe}_3)_2]_3$, was later reported by Lappert et al.¹² All of these bismuth amides have been characterized spectroscopically but without X-ray structure determinations. $\text{Me}_2\text{Bi}[\text{NMe}(\text{SiMe}_3)]$ and $\text{MeBi}[\text{NMe}(\text{SiMe}_3)]_2$ are liquids; $\text{Bi}[\text{NMe}(\text{SiMe}_3)]_3$ has a melting point of 27-29 °C; and $\text{Bi}[\text{NMe}(\text{SiMe}_3)]_3$ has a melting point around 90 °C (decomposition). Veith et al. obtained $\text{Bi}_2[(\text{N}t\text{Bu})_2\text{SiMe}_2]_3$ (Scheme 4) starting from the lithium salt of the diamine ($t\text{BuN}(\text{H})\text{-SiMe}_2\text{-N}(\text{H})t\text{Bu}$).¹³ If the ratio of BiCl_3 and the lithium salt is 1:1, the corresponding bismuth chlorine compound $[\text{Me}_2\text{Si}(\text{N}t\text{Bu})_2\text{Bi}]\text{Cl}$ (Scheme 4) can also be isolated.


Scheme 4



Metal-amido complexes have been proposed as key intermediates in important catalytic processes and especially in the fixation and activation of nitrogen.¹⁴ Insertion reactions of carbonyl and alkene compounds into metal nitrogen bonds have been known for lanthanides and early and late transition metals complexes. However, the insertion of carbon-carbon, carbon-oxygen and carbon-nitrogen bonds into the metal-nitrogen bond has not well been studied with main-group compounds. This may be due the poor reactivity of compounds with nitrogen bonds. Lappert et al. described the addition of aminostannanes to a variety of

alkines and alkenes.¹⁵ They also reported aminosilylation and aminophosphination reactions with highly electrophilic substrates.¹⁶ Recently Hartwig and coworkers reported transamination of alkenes and vinylarenes by rhodium (I) amides.¹⁷ The insertion of alkyne into a molybdenum amid bond has also been described.¹⁸

The design and synthesis of heterobimetallic bismuth complexes with a stoichiometry appropriate for use as direct precursors for advanced materials is an ambitious challenge. The coordination chemistry of all of the heavy main group elements, and bismuth in particular, is complicated by the high latent Lewis acidity of the metal center combined with its ability to easily expand its coordination sphere. Most work has concentrated on bimetallic systems which possess bridging ligands coordinated to two contiguous metals.¹⁹⁻²³ Other constructions involve tethering two metal complexes by non-coordinating links.²⁴⁻²⁶ Metals may be linked by one (1),²⁷⁻²⁹ two (2)³⁰⁻³³ or three (3)³⁴⁻³⁶ bridging groups of which the most prevalent is the two-bridge system 2.



In reviewing literature, several general routes to the formation of heterobimetallic coordination complexes can be surmised.³⁷ Specifically, these are salt metathesis reactions, volatile elimination reactions, bifunctional ligands, Lewis acid-base adduct formation and alkali metal reduction. Not all of these synthetic approaches are available for the development of precursors for advanced oxide materials. Significant efforts have been made to develop solution-based routes to the formation of oxide based materials, and numerous solution-based approaches for the formation of bi- or multimetallic oxide systems have been reported including sol-gel, metal organic decomposition (MOD) routes. These methods have been used successfully to produce numerous heterobimetallic systems. It has been reported that oxides

synthesized by these approaches have properties that are superior to those produced in solid-state reactions.³⁸ Further, it has also been observed that the properties of many oxides produced by wet chemical routes, including the reactivity of material and conditions required for crystallization of the phase, are sensitive to the manner in which the different metal precursors are brought together prior to decomposition.³⁹ Recently these complexes have attracted increasing interest due to their potential applications as high T_c superconductors,⁴⁰ in nonlinear optics,⁴¹ in oxidation reactions, as catalysts,⁴² and as thermoelectric and ferroelectric materials. It has been observed that the bismuth germanium oxide system is the best candidate for electro-optic voltage sensors⁴³ and the mixed oxides of aluminum and bismuth show high oxide-ion conductivity.⁴⁴


Bismuth chalcogenides are an interesting class of compounds with amazing structural and compositional complexity. The stereochemical activity of the $6s^2$ lone pair of electrons of Bi influences the structures, and thus the properties of the compounds. Diorganobismuth chalcogenides of the type $(R_2Bi)_2X$ (X = chalcogen, R = alkyl, aryl) have been under investigation for a long time but little is known about their structural chemistry. These compounds are of interest as potential precursors for semiconducting bismuth chalcogenides, Bi_2E_3 (E =chalcogen), and therefore they can display interesting molecular and supramolecular architectures. Several well-defined organobismuth chalcogenides have been synthesized and their crystal structures were determined such as $(R_2Bi)_2E$ [R = Mes, E = O, S, Se].⁴⁵

Dibismuthines R_2BiBiR_2 are a type of compound which contain a bismuth-bismuth bond. R can be an alkyl group ($HC(Me_3Si)_2$,⁴⁶ Me, Et, Pr, Bu, or iPr ⁴⁷), or an alkene or aromatic group ($CH_2=C(CH_3)$ -,⁴⁷ $(CH_3)_2C=CH$ -,⁴⁷ Ph,⁴⁸ 2,4,6-trimethylphenyl,⁴⁹ 2-(Me_2NCH_2) C_6H_4 ,⁵⁰ 2,6-(Me_2NCH_2) $_2C_6H_3$,⁵¹ or p -methylphenyl⁵²). They have characteristic physical and chemical properties due to the Bi-Bi bond.⁵³ The most interesting is thermochromism: when melted or dissolved in organic solvents, some dibismuthines show drastic color changes. Recent studies have revealed that the intermolecular $Bi \cdots Bi$ interaction of dibismuthines in solid state has a


The importance of organoaluminum compounds is due to their use in a variety of applications, including organic synthesis and industrial catalytic processes. They have also been utilized as precursors in chemical vapor deposition (CVD) processes.⁶² Compounds such as hexakis(dimethylamido)dialuminum, $\text{Al}_2(\text{NMe}_2)_6$,⁶³⁻⁶⁵ azides of formula $(\text{R}_2\text{AlN}_3)_3$ ($\text{R} = \text{Me}$ or Et)⁶⁶⁻⁶⁸ and amides of formula $(\text{R}^1_2\text{AlNR}^2)_n$ ($\text{R}^1 = \text{Me}$; $\text{R}^2 = \text{H}, \text{Me}$ or $i\text{Pr}$; $n = 2$ or 3)^{69,70} have been used to deposit polycrystalline AlN films, although an atmosphere of ammonia is required to obtain good-quality material. In polymerization catalysis, main group compounds play important roles as activators in the generation of cationic transition-metal alkyl species that polymerize α -olefins.⁷¹ It has been observed that stable aluminum compounds involving nitrogen substituents or neutral nitrogen donors are useful in the preparation of aluminum nitrides, AlN based semiconductors, and AlN ceramics.⁷²⁻⁷⁴ The chemistry of compounds containing AlN bonds has flourished over the past several years due mainly to current interest in developing optimum AlN precursors.^{75,76} Earlier work on the structures and properties of the alkyl aluminum amides has focused largely on the alkyl and arylamine derivatives $\text{R}_2\text{AlNR}'\text{R}''$ where both R' and R'' are organic groups or where one of these substituents is hydrogen. These aluminum derivatives show a strong tendency to oligomerize due to the Lewis acidity of the aluminum center and to the presence of the electron lone pair of the amido group. The steric bulk of the substituents is the major factor that determines the degree of association.⁷⁷ Therefore, these compounds are generally found in the form of oligomers of the type $[\text{R}_2\text{AlNR}'\text{R}'']_n$ with either four- ($n=2$) or six-membered ($N=3$) $(\text{AlN})_n$ rings. Alkylaluminum amides have a very rich and diverse structural chemistry. When a trialkylalane reacts with primary or secondary amines, the amides **A**, **B**, and **C** are formed in addition to simple adducts (Schemes 6, 7). Compounds of type **A** or **B** are cyclic and those of type **C** are polycyclic, as shown by spectroscopic data or X-ray structure determinations. $(\text{Me}_2\text{-AlNMe}_2)_2$ forms a planar four-membered cycle⁷⁸ but $(\text{Me}_2\text{AlNHMe})_3$

has a six-membered Al_3N_3 ring.⁷⁹ The polycyclic oligomers $(\text{RAINR})_n$ (type **C** compounds) may adopt an $\text{Al}_n\text{-N}_n$ skeleton that resemble either a cube ($n = 4$) or a hexagonal prism ($n = 6$)

Scheme 6




Scheme 7



As in the compounds of type **A** or **B**, both the aluminum and nitrogen atoms always have the coordination number four. When the substituents on the secondary amine in Scheme 7 are bulky or when trimethylsilyl groups are employed, compounds are isolated that have only one amide function in the product (Scheme 8, $\text{R} = \text{Ph}, \text{SiMe}_3$).⁸⁰

Scheme 8



There are, however, only a few examples known of monomeric aluminum amides in addition to compounds of type **A** and **B**, some of which have been claimed to be monomeric [e.g. $\text{Me}_2\text{AlN}(\text{SiEt}_3)_2$, $\text{Me}_2\text{Al-NPhSiMe}_3$].^{80,81}

Particularly, there has been immense research interest in synthesizing aluminum chlorides and methyl derivatives which can act as precursors for heterobimetallic complexes and aluminum hydrides which can be useful for preparing aluminum chalcogenides of the formula $(\text{RAIE})_n$ ($\text{R} = \text{organic ligand}; \text{E} = \text{S, Se, Te}; n = 2, 4$)⁸², and also can be used for preparing

aluminum containing heteroatom rings⁸³ by H₂ elimination reactions. Aluminum alkyls can be converted to aluminum halides or aluminum oxides. Recently a series of aluminum oxides and aluminum hydroxides were synthesized by the hydrolysis reaction of aluminum alkyls.⁸⁴

1.3. Aims and Scope of the Present Work

As discussed above, organobismuth and organoaluminum compounds have been widely investigated as cocatalysts, reagents as well as precursors for material science and industrial application. The aim of this thesis is the following by using the appropriate ligand system:

- To synthesize bismuthamide complex and investigate its property and reaction with compounds containing unsaturated C-O, C-C, and C-N bonds;
- To use this species as a synthon to assemble a variety of heterobimetallic, chalcogenide and low-valent systems for potential application;
- To use spectral methods such as NMR spectroscopy, IR spectroscopy and X-ray structural analysis to unambiguously characterize the products obtained;
- To synthesize aluminum compounds stabilised by the iminophosphonamide ligand.

2. Results and Discussion

2.1. Bismuthamidation of Unsaturated Organic Compounds


A bulky organic ligand is usually employed to stabilize the bismuth center electronically and/ or sterically. 1,8-Diaminonaphthalene ligands have been shown to be a very useful platform for the preparation of organometallic compounds with both transition and main group metals because the diamide ligand system generates a unique metallic center in terms of the geometry and the electronic property. In 1998, a series of new divalent bulky 1,8-diaminonaphthalene ligands were reported, which are currently exhibiting growing interest due to their specific steric and electronic properties to control effectively the geometry at the metal center. The known complexes containing these ligands include main group (Al, Ge, In, Li, Mg, Si, Sn, Tl) and transition metal (Ni, Ti, Zr) derivatives.⁸⁵⁻⁸⁸

The insertion of carbon-carbon, carbon-nitrogen and carbon-oxygen bonds into the bismuth-nitrogen bond has not well been studied. This may be attributed to the poor reactivity of compounds with nitrogen bonds. The only known compounds to this category are the reactions of bismuth amide with heterocumulenes by Ando et al.⁸⁹ The work reported herein was fueled by the design and synthesis of main group complexes with the focus on their reactivity toward insertion reactions. We have recently reported the synthesis, structural characterization and reactions of tin(II) hydride species with compounds containing unsaturated C-O, C-C, and C-N bonds, which resulted in simultaneous hydrogen and {LSn} transfer to the organic substrates.⁹⁰ Herein we report the synthesis and structural characterization of a bismuth amide complex and its reaction with compounds containing unsaturated C-O, C-C, and C-N bonds.

2.1.1 Preparation of 1,8- $C_{10}H_6(NSiMe_3)_2BiNMe_2$ (**1**)

Compound 1,8-bis[(trimethylsilyl)amino]naphthalene (L^1H_2) was prepared by a previously reported method.⁸⁸ The reaction of 1,8-bis[(trimethylsilyl)amino]naphthalene with $Bi(NMe_2)_3$ ⁶ in *n*-hexane at room temperature results in the formation of L^1BiNMe_2 (**1**) [$L^1=1,8-C_{10}H_6(NSiMe_3)_2$] in good yield (Scheme 9). **1** is an orange solid, soluble in *n*-hexane, THF and toluene. Compound **1** is moderately air and moisture sensitive, and is slowly oxidized and hydrolyzed on exposure to air. It was characterized by multinuclear NMR, elemental analysis, and X-ray structural analysis. The 1H NMR spectrum of **1** exhibits two singlets for the $SiMe_3$ and NMe_2 groups, respectively. (δ 0.25 and 2.48 ppm). The ^{13}C NMR spectrum of **1** reveals two resonances (δ 41.98, 3.05 ppm), which can be assigned to the carbon resonances arising from NMe_2 and $SiMe_3$ groups, respectively. The ^{29}Si NMR spectrum of **1** exhibits one singlet (δ 3.306 ppm) attributed to the silicon resonances arising from $SiMe_3$ groups.

Scheme 9



The molecular structure of L^1BiNMe_2 (**1**) has been determined by single crystal X-ray diffraction. **1** crystallizes in the monoclinic space group $P2_1/n$ and the molecular structure (Figure 1) features a trigonal pyramidal coordinated bismuth atom. The N–Bi–N bond angles sum to 281.42° , hence prove the stereochemically active lone pair in the apical position. The

Bi–N distances for **1** (average 2.168 Å) are comparable to those observed in structurally characterized $\text{Bi}(\text{NMe}_2)_3$.^{6,91}





Figure 1. Crystal structure of **1**, anisotropic displacement parameters depicted at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1–N1 2.150(1), Bi1–N2 2.176(1), Bi1–N3 2.179(1), N1–Bi1–N2 85.02(7), N1–Bi1–N3 100.06(7), N2–Bi1–N3 96.34(6).

2.1.2. Reaction of $L^1\text{BiNMe}_2$ (**1**) with Ketone and Aldehyde: Synthesis of $L^1\text{BiOCPh}(2\text{-Py})(\text{NMe}_2)$ (**2**) and $L^1\text{BiOCH}(3\text{-Py})(\text{NMe}_2)$ (**3**)

The reaction of **1** with 2-benzoylpyridine proceeds smoothly at 25 °C in *n*-hexane to give product **2**, which was subsequently isolated in 83% yield (Scheme 10). The ^1H and ^{13}C NMR spectra indicate the formation of an addition product in which 2-benzoylpyridine inserts into the Bi–N bond of **1** under formation of **2**. The ^1H NMR spectrum of **2** exhibits two different resonances (δ 0.47 and 0.32 ppm) for the SiMe_3 groups and a singlet for the NMe_2 substituent at δ 1.59 ppm. The ^{13}C NMR spectrum of **2** shows an upfield-shifted carbonyl resonance of the inserted ketone which appears at δ 94.8 ppm in comparison to the starting material (δ

193.7 ppm) and three resonances (δ 38.68, 3.15, 3.06 ppm), which can be assigned to the carbon resonances arising from NMe_2 and two SiMe_3 groups, respectively. The ^{29}Si NMR spectrum of **2** exhibits the SiMe_3 to resonate at δ 0.034 and -0.81 ppm with upfield-shifted signals compared to those of **1**.

Scheme 10



Crystals of **2** suitable for an X-ray diffraction study were grown from *n*-hexane solution at -30 °C. **2** is monomeric in the solid state and crystallizes in the triclinic space group $P\bar{1}$ (Figure 2). The coordination environment around the bismuth atom is distorted trigonal pyramidal with the bond angles in the range of $83.46(12)$ to $97.03(11)^\circ$ around the bismuth atom. The Bi–O distance ($2.089(2)$ Å) is in the range of a Bi–O single bond when compared with other structural characterized bismuth compounds.⁹² The Bi–N4 (donor side-arm) distance of $2.615(4)$ Å is considerably longer than the Bi–N bond lengths of the ligand ($2.194(3)$ and $2.123(3)$ Å).




Figure 2. Crystal structure of **2**, anisotropic displacement parameters depicted at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1-O1 2.089(2), Bi1-N1 2.194(3), Bi1-N2 2.123(3), Bi1-N4 2.615(4), O1-C17 1.405(4), O1-Bi1-N1 85.44(11), O1-Bi1-N2 97.03(11), N1-Bi1-N2 83.46 (12), Bi1-O1-C17 125.4(2).

In a similar manner a *n*-hexane solution of **1** smoothly reacts with 3-pyridinecarboxaldehyde at room temperature to afford crystalline **3** in 75% yield (Scheme 10). The characterization of **3** indicates that the insertion of 3-pyridinecarboxaldehyde occurs selectively into the Bi–N bond of **1**. The ^1H NMR spectrum of **3** exhibits singlet resonances for the dimethylamine and quaternary CH protons, respectively (δ 1.89 and 5.05 ppm). The three methyl groups of the SiMe_3 ligands appear at δ 0.3 and 0.26 ppm. The ^{13}C NMR spectrum of **3** shows a singlet resonance (δ 89.23 ppm), which can be assigned to the carbonyl group of the inserted aldehyde, whereas the carbon atoms of SiMe_3 and NMe_2 groups resonate

at 3.06, 3.05 and 39.35 ppm respectively. **3** shows a resonance at 1.92 and 1.77 ppm in its ^{29}Si NMR spectrum.

3 crystallizes in the triclinic space group $P\bar{1}$ with two molecules in the asymmetric unit and the bismuth atom exhibits a trigonal pyramidal environment with bond angles of 83.96(8) to 91.85(9) $^\circ$ (Figure 3). The Bi–O distance in **3** (2.10(2) Å) is in the range of Bi–O single bonds when compared to literature examples (2.058 and 2.109 Å in $[\text{Bi}(\text{ONp})_3(\text{Py})_2]$).⁹² In this structure also the Bi–N4 (donor side-arm) distance of 2.891(4) Å is considerably longer than the Bi–N bond lengths of the ligand (2.170(2) and 2.164(2) Å).





Figure 3. Crystal structure of **3**, anisotropic displacement parameters depicted at the 50% probability level. Hydrogen atoms (except for those of C17) and disorder in one SiMe_3 group are omitted for clarity. Selected bond lengths [Å] and angles [$^\circ$]: Bi1–O1 2.10(2), Bi1–N1 2.170(2), Bi1–N2 2.164(2), Bi–N4 2.891(4), O1–C17 1.388(4), O1–Bi1–N1 83.96(11), O1–Bi1–N2 91.85(9), N1–Bi1–N2 84.02 (9), Bi1–O1–C17 113.70(18)

2.1.3. Reaction of $L^1\text{BiNMe}_2$ (1) with Alkene and Alkyne: Synthesis of $L^1\text{BiC(Me)(CN)CH}_2\text{NMe}_2$ (4) and $L^1\text{BiC(CO}_2\text{Et)CNMe}_2(\text{CONMe}_2)$ (5)

The driving force for the formation of compounds **2** and **3** is the Bi–O bond energy. For monomeric BiO it was determined by mass spectrometric measurement ($338.9 \pm 5.9 \text{ kJmol}^{-1}$).⁹³ This value may well differ from a single Bi–O bond energy in the solid. Additionally the intramolecular coordination between bismuth and nitrogen (see Figures 2 and 3) may play an important role for the stability of these compounds. To extend those findings to bismuth-carbon bonds, we reacted **1** with an alkene and alkyne, respectively, although aware that the Bi–C bond is considerably weaker (143 kJmol^{-1})⁹⁴ than the Bi–O bond. The reaction of **1** and 2-methyl-2-propenenitrile in *n*-hexane at 25 °C gives the yellow complex **4** in 74 % yield (Scheme 11). Formation of **4** proceeds with good yield and no decomposition products are observed. The ¹H NMR spectrum of **4** contains two nonequivalent CH protons of the methylene residue (δ 2.43 and 4.53 ppm) and two singlets for NMe₂ and Me (δ 2.08 and 1.17 ppm). The ¹³C NMR spectrum indicates the presence of a new resonance (δ 62.6 ppm), which can be assigned to the carbon atom that binds to the bismuth atom. **4** shows a resonance at δ 5.41 and 3.46 ppm in its ²⁹Si NMR spectrum.

Scheme 11

Like the other structures, **4** is monomeric in the solid state and crystallizes in the monoclinic space group $P2_1/c$ and in this structure again the bismuth atom shows the anticipated trigonal pyramidal coordination geometry with the bond angles in the range of 84.95(12) to 100.15(12)° around the bismuth atom. Moreover, the crystal structure of **4** reveals that the bismuth atom and the NMe₂ group are arranged at the same side of the molecule. This might be caused by the packing of the molecules in the crystal. The C17–C20 bond length of 1.540(5) Å is in the range of normal C–C single bonds (Figure 4). The Bi1–C17 bond length (2.341(3) Å) in **4** is longer than those in [*t*BuN(CH₂C₆H₄)₂BiMe] (2.264(6) Å),⁹⁵ whereas the Bi–N bond lengths (Bi1–N1 2.171(3), Bi1–N2 2.155(3) Å) are comparable to those observed in **1**. In this structure also the Bi–N4 (donor side-arm) distance of 2.936(6) Å is considerably longer than the Bi–N bond lengths of **1**.





Figure 4. Crystal structure of **4**, anisotropic displacement parameters depicted at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1–N1 2.171(3), Bi1–N2 2.155(3), Bi1–N4 2.936(6) Bi1–C17 2.341(3), N4–C20

2. Results and Discussion

1.462(5), C17-C20 1.540(5), N1-Bi1-C17 100.15(12), N2-Bi1-C17 94.37(12), N1-Bi1-N2 84.95(12), C20-C17-Bi1 100.5(2), N4-C20-C17 112.8(3).

The reaction of **1** with diethyl acetylene dicarboxylate at 25 °C in a molar ratio of 2:1 resulted in the yellow product **5** in 55% yield (Scheme 12). During the formation of **5** the exchange of one of the OEt groups by a NMe₂ substituent occurred. The ¹H NMR spectrum of **5** displays three resonances as a singlet (δ 2.43, 0.52 and 0.53 ppm) which can be attributed to the Me protons of NMe₂ and SiMe₃ groups, whereas non equivalent Me protons of the substituent NMe₂ resonate as a singlet at δ 2.42 and 1.20 ppm. The ¹³C NMR spectrum of **5** reveals three resonances (δ 40.95, 3.31 and 3.206 ppm), which can be assigned to the carbon resonances arising from inserted NMe₂ and SiMe₃ groups, respectively, whereas the carbon atoms of the resulting C=C double bond in **5** resonate at δ 169.3 and 174.4 ppm. **5** shows a resonance at δ 2.154 and 1.951 ppm in its ²⁹Si NMR spectrum.

Scheme 12



The crystal structure of **5** is shown in Figure 5. **5** is monomeric in the solid state and crystallizes in the triclinic space group *P* $\bar{1}$ with a trigonal pyramidal geometry at the bismuth atom. The structure reveals that the addition resulted in a *Z*-orientation of bismuth and the amide relative to the double bond. The Bi–O1 (O_{carbonyl} donor) distance (2.70(40) Å) is

comparable with the Bi-O_{carbonyl} interaction that ranges from 2.56 to 2.86 Å in other known structures.⁹⁶ The Bi1-C18 bond length (2.29(8) Å) in **5** is comparable to those in [*t*BuN(CH₂C₆H₄)₂BiCCPh] (2.289(4) Å)⁹⁵ but shorter than in **4** (2.341(3) Å).




Figure 5. Crystal structure of **5**, anisotropic displacement parameters depicted at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1-N1 2.178(3), Bi1-N2 2.20(30), Bi1-O1 2.70(40), Bi1-C18 2.29(8), C18-C19 1.381(5), N1-Bi1-C18 92.2(5), N2-Bi1-C18 103(6), N1-Bi1-N2 81.64(16).


2.1.4. Reaction of *L*¹BiNMe₂ (**1**) with *N,N'*-diisopropylcarbodiimide: Synthesis of 1,8-*C*₁₀*H*₆(*NSiMe*₃)₂Bi(*NiPr*)₂CNMe₂ (**6**)

Compounds **2-5** show reactivity of bismuth-nitrogen bond toward carbon-carbon and carbon-oxygen bonds. To extend those findings to carbon-nitrogen bonds, we reacted **1** with diisopropylcarbodiimide and phenylisocyanate, respectively.

Reaction of diisopropylcarbodiimide with **1** in *n*-hexane results in the formation of **6**, indicating that one carbodiimide molecule is inserted into the Bi-N bond (Scheme 13).

Compound **6** is an orange crystalline solid, which is soluble in organic solvents such as *n*-hexane, THF, and toluene. The ^1H and ^{13}C NMR spectra also confirm the insertion of diisopropylcarbodiimide into the Bi–N bond of **1** under formation of **6**. The ^1H NMR spectrum of **6** displays two resonances (δ 1.99, 0.34 ppm) which can be attributed to the NMe_2 and SiMe_3 groups, respectively, whereas the two and twelve protons of N-isopropyl groups resonate at δ 3.93 and 0.94 ppm. The ^{13}C NMR spectrum of **6** shows upfield-shifted resonances of NMe_2 (δ 39.7 ppm), in comparison to **1** (δ 41.9 ppm), whereas the carbon atoms of the N-isopropyl and SiMe_3 groups resonate at δ 39.7, 24.9, and 2.7 ppm, respectively. The ^{29}Si NMR spectrum of **6** exhibits a singlet resonance (δ 0.21 ppm) which can be assigned to SiMe_3 groups.

Scheme 13



Single crystals of **6**, suitable for X-ray diffraction studies, were obtained from a saturated solution of *n*-hexane at -30 °C. In **6** the bismuth atom is tetracoordinate in an irregular tetragonal-pyramidal geometry. Compound **6** crystallizes in the monoclinic space group $P2_1/n$ with one molecule in the asymmetric unit (Figure 6). The bismuth center in **6** is coordinate to two nitrogen atoms of the anionic ligand and to two nitrogen atoms of carbodiimide unit. The C–N bond distances associated with the central sp^2 carbon of the carbodiimide species are almost equal and are considerably shorter than a typical C–N single bond, suggesting that the

three C-N bonds possess a partial double-bond character with typical C(17)-N(3), C(17)-N(4), and C(17)-N(5) distances of 1.380(2), 1.309(2), and 1.369(2) Å respectively.





Figure 6. Molecular structure of **6**. Thermal ellipsoids are set at 50% probability level. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi(1)-N(1) 2.2356(13), Bi(1)-N(2) 2.1418(14), Bi(1)-N(3) 2.2035(14), Bi(1)-N(4) 2.5259(14), N(1)-Si(1) 1.7316(14), N(2)-Si(2) 1.7319(14), C(17)-N(3) 1.380(2), C(17)-N(4) 1.309(2), C(17)-N(5) 1.369(2), N(1)-Bi(1)-N(2) 81.51, N(2)-Bi(1)-N(3) 96.87(5), N(3)-Bi(1)-N(4) 56.83(5), N(1)-Bi(1)-N(3) 91.34(5), N(1)-Bi(1)-N(4) 144.97(5), N(2)-Bi(1)-N(4) 87.78(5).

2.1.5. Reaction of $L^1\text{BiNMe}_2$ (**1**) with Phenylisocyanate: Synthesis of 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{BiNPhCONMe}_2$ (**7**)

In a similar manner to **6**, *n*-hexane solution of **1** smoothly reacts with phenylisocyanate at room temperature to afford crystalline **7** in 69% yield under isocyanate insertion into the Bi-N bond of **1** (Scheme 14). The ^1H NMR spectrum of **7** in C_6D_6 exhibits two singlets (δ 2.08, 0.31 ppm) for NMe_2 and SiMe_3 groups. The ^{13}C NMR spectrum of **7** shows an upfield-shifted resonance of NMe_2 (δ 37.5 ppm) in comparison to **1** (δ 41.9 ppm), confirming the insertion of

phenylisocyanate into the Bi–N bond of **1**, whereas the carbon atoms of SiMe₃ resonate as a singlet at δ 2.71 ppm. The ²⁹Si NMR spectrum of **6** displays one resonance at δ 2.93 ppm which can be attributed to the SiMe₃ groups.

Scheme 14



Yellow crystals of **7**, suitable for single-crystal X-ray diffraction study, were obtained from a saturated solution of *n*-hexane at -30 °C. The coordination geometry around the bismuth atom in **7** is a distorted pyramidal arrangement with a stereochemically active electron lone pair. Compound **7** crystallizes in the triclinic space group $P\bar{1}$ with two molecules in the asymmetric unit (Figure 6). In **7** the N–C (av 1.384 Å) and C–O (av 1.255 Å) bond distances of the ureato fragments are both in the range of single and double bonds, indicating that the π electrons of the ureato fragments are partially delocalized in the NC(N)O core. Additionally, the intramolecular coordination between bismuth and oxygen may play an important role for the stability of this compound.




Figure 7. Molecular structure of **7**. Thermal ellipsoids are set at 50% probability level. Only one of the two crystallographically independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^\circ$]: Bi1-N 2.1579(14), Bi1-N2 2.1183(14), Bi1-N3 2.2334(15), N1-Si1 1.7371(15), N2-Si2 1.7297(15), Bi1-O1 2.6364(13), C17-N3 1.373(2), C17-N4 1.351(2), C17-O1 1.266(2), N1-Bi1-N2 85.40(6), N1-Bi1-N3 92.88(5), N2-Bi1-N3 97.08(6), C17-N3-Bi1 101.30(11), N1-Bi1-O1 146.87(5), N2-Bi1-O1 93.36(5)

2.2. Heterobimetallic Bismuth Complexes with Main Group and Transition Metals


Since the first isolation and characterization of a bismuth-transition metal heterobimetallic alkoxide in 1996,⁹⁷ a wide variety of bismuth heterobimetallic complexes have been synthesized and reviewed.⁹⁸ Limberg and coworkers reported the first structurally characterized molecular complexes with Bi-O-Mo linkages. These moieties are thought to be the active oxo-transfer sites during allylic oxidation of propene.⁹⁹ The solid state structure of the first coordination complex containing bismuth and aluminum has also been described.¹⁰⁰ We have recently reported the synthesis and structural characterization of μ -oxo bridged Al-O-M (M= Zr, Ti, Hf), M-O-M¹ (M= Ti, Zr; M¹= Al, Ga), Ge-O-M (M= Zr, Hf), and Al-O-M-

O-Al (M= Ti, Zr) complexes.¹⁰¹ Herein we report a general synthetic approach and structural characterization of five bismuth heterobimetallic complexes with main group and transition metals.

2.2.1. Reaction of $L^1\text{BiNMe}_2$ (**1**) with $\text{LM}(\text{Me})(\text{OH})$: Synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{MMeL}$ (M = Al **8**; Ga, **9**)

Treatment of **1** with stoichiometric amounts of $\text{LM}(\text{Me})(\text{OH})$ (M=Al, Ga) in *n*-hexane at room temperature results in the formation of **8** and **9** exhibiting the Bi-O-M (M=Al **8**, Ga **9**) moiety (Scheme 15). Compounds **8** and **9** are soluble in toluene, THF, and *n*-hexane and are characterized by analytical, spectroscopic, and single-crystal X-ray diffraction studies. Compounds **8** and **9** are yellow crystalline solids that melt at 174 °C and 191 °C. The ^1H NMR spectra of the reaction mixture of compounds **8-9** show almost quantitative conversion of the reactants to products, as revealed by the absence of any M-OH and NMe_2 resonances. The ^1H NMR spectrum of **8** displays two resonances (δ -0.69 and 0.24 ppm) which can be attributed to the Me protons of AlMe and SiMe₃ groups, respectively, whereas the particular GaMe and SiMe₃ groups in compound **9** resonate at δ -0.56 and 0.28 ppm. In addition, a set of resonances assignable to the isopropyl and methyl protons associated with the β -diketiminato ligand is found in the range between δ 1.31 and 0.99 ppm, and the absence of the OH proton resonance features both **8** and **9**. The mass spectral data of compounds **8-9** are in agreement with the assigned structures. Compound **8** is moderately air and moisture sensitive, but thermally quite stable as indicated by the high melting point and EI mass spectrum in which the molecular ion (M^+) was observed with 100% intensity. The next most intense peak at m/z 969.5 was assigned to $[\text{M-Me}]^+$.

Scheme 15



Yellow crystals of **8** suitable for single-crystal X-ray diffraction study were obtained from a saturated solution of *n*-hexane at -30 °C and for those of **9** from a mixture of *n*-hexane/toluene solution at -30 °C. Compounds **7** and **8** crystallize in the monoclinic space group $P2_1/c$, with two and one molecules in the asymmetric unit, respectively (Figures 8, 9). The aluminum and gallium atoms adopt each a distorted tetrahedral environment, whereas the coordination geometry around the bismuth atom is that of a distorted pyramidal arrangement with a stereochemically active electron lone pair. The Al–O distance (av 1.834 Å) is similar to that which has been reported for complex $[\{Bi(Hsal)_3\}_2\{Al(acac)_3\}]$ (av 1.871 Å).¹⁰⁰ The Al(μ -O)Bi angle (av 160.6°) in **2** is considerably bigger than the corresponding Al(μ -O)Ga bond angle in **3** (123.25(7)°).




Figure 8. Molecular structure of **8**. Only one of the two independent molecules in the crystal is shown. Anisotropic displacement parameters are depicted at the 50% probability level. H atoms and disordered SiMe₃-groups are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1-O1 1.995(4), Bi1-N2 2.115(4), Bi1-N1 2.157(4), Al1-O1 1.690(3), Al1-N3 1.902(4), Al1-N4 1.931(4), Al1-C46 1.959(5), O1-Bi1-N2 92.55(16), O1-Bi1-N1 95.16(16), N2-Bi1-N1 85.5417, O1-Al1-N3 111.13(18), O1-Al1-N4 111.14(18), N3-Al1-N4 95.97(16), O1-Al1-C46 116.2(2), N3-Al1-C46 111.8(2), N4-Al1-C46 108.8(2), Al1-O1-Bi1 160.7(2)

The Ga-N bond lengths (1.963(16) and 1.973(16) Å) and the N-Ga-N angle (95.87(6)°) are comparable to those observed in [Me₂Ga(2-NC₅H₄)₂CH].¹⁰² The angle for the bent Ga-O-Bi linkage is 123.25(7)°, which is considerably more acute than that in LGa(Me)(μ-OH)LnCp₃ (Ln=Sm, Nd; av 149.5°). The corresponding Ga-O bond length (1.815(13) Å) is shorter than the Ga-μ-OH bond in LGa(Me)(μ-OH)LnCp₃ (av 1.856 Å).¹⁰³ Obviously there is more charge accumulated at the O²⁻ in **3** than at the OH⁻ in the latter complex.





Figure 9. Molecular structure of **9**. Anisotropic displacement parameters are depicted at the 50% probability level. H atoms and one toluene solvent molecule are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1-O1 2.0282(13), Ga1-O1 1.8159(13), Ga1-C100 1.955(2), Bi1-N1 2.1425(17), Bi1-N2 2.1443(16), O1-Ga1-N4 103.90(6), C100-Ga1-N3 111.54(8), C100-Ga1-N4 113.33(8), O1-Bi1-N1 93.67(6), N1-Bi1-N2 85.13(6), N3-Ga1-N(4) 95.87(6), Ga1-O1-Bi1 123.25(7), O1-Bi1-N2 94.02(6), O1-Ga1-N3 109.26(6), O1-Ga1-C100 120.06(8).

2.2.2. Reaction of $L^1\text{BiNMe}_2$ (**1**) with LGe(OH) : Synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{GeL}$ (**10**)

Similarly the reaction of **1** with LGe(OH) in *n*-hexane at room temperature affords $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{GeL}$ (**10**) in moderate yield (63%) (Scheme 16). The ^1H NMR and ^{29}Si NMR spectra of **10** exhibit singlets at δ 0.24 and 0.99 ppm, respectively, corresponding to the SiMe_3 groups. Complex **10** is a yellow crystalline solid which is soluble in common organic solvents such as toluene, THF, and *n*-hexane and is stable at room temperature for several months under an inert atmosphere. In the mass spectrum of **10** the peak with the

highest mass corresponds to the molecular ion M^+ and the next most intense peak is observed at m/z 599 which can be assigned to the fragment $[M-CH(DippNCMe)_2]^+$.

Scheme 16



Compound **10** crystallizes in the monoclinic space group $P2_1/n$ with two molecules and one disordered *n*-hexane in the asymmetric unit. As expected, **10** contains the bismuth atom bonded through an oxygen atom to germanium (Figure 10). The bismuth and germanium atoms both exhibit a highly distorted pyramidal geometry, involving the chelating 1,8- $C_{10}H_6(NSiMe_3)_2$ ligand and β -diketiminato ligand, the (μ -O) atom and a stereochemically active lone pair at each atom. The Ge-N bond lengths (av 2.010 and 2.049 Å) and N-Ge-N angle (av 87.47°) are comparable to those of oxygen-bridged heterobimetallic complexes reported in reference 104. **10** features a bent Ge-O-Bi (av 118.89°) arrangement with a Ge-O distance of av 1.825 Å and a Bi-O bond length of av 2.052 Å. The Ge-O bond length is longer than those in Ge-O-Yb (1.769 Å) and Ge-O-Y (1.762 Å) species¹⁰⁵ and is comparable with those in Ge-O-Hf (1.799 Å) and Ge-O-Zr (1.797 Å) arrangements.¹⁰⁴




Figure 10. Molecular structure of **10**. Only one of the two independent molecules in the crystal is shown. Anisotropic displacement parameters are depicted at the 50% probability level. H atoms and half a molecule of *n*-hexane are omitted for clarity. Selected bond lengths [Å] and angles [°]: Ge1-O1 1.8231(11), Ge1-N3 2.0261(13), Ge1-N4 2.0556(13), Bi1-O1 2.0493(11), Bi1-N1 2.1389(14), Bi1-N2 2.1614(14), Ge1-O1-Bi(1) 125.20(6), O1-Bi1-N1 96.72(5), O1-Bi1-N2 93.05(5), N1-Bi1-N2 84.77(5), O1-Ge1-N3 97.57(5), O1-Ge1-N4 92.28(5), N3-Ge1-N4 87.68(5).


2.2.3. Reaction of $L^1\text{BiNMe}_2$ (**1**) with $\text{Cp}^*_2\text{MR}(\text{OH})$: Synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{MRCp}^*_2$ ($M = \text{Zr}$ **11**, $R = \text{Me}$; Hf , **12**, $R = \text{Cl}$)

The synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{ZrMeCp}^*_2$ (**11**) and $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{HfClCp}^*_2$ (**12**) are accomplished by reacting **1** with $\text{Cp}^*_2\text{ZrMe}(\text{OH})$ and the in situ reaction with $\text{Cp}^*_2\text{HfCl}(\text{OH})$, respectively (Scheme 17). $\text{Cp}^*_2\text{HfCl}(\text{OH})$ was prepared from $\text{Cp}^*_2\text{HfCl}_2$ and water in the presence of a N-heterocyclic carbene $[\text{CN}(\text{iPr})\text{C}_2\text{Me}_2\text{N}(\text{iPr})]$. The carbene is used for trapping quantitatively the generated HCl from hydrolysis of $\text{Cp}^*_2\text{HfCl}_2$.

2. Results and Discussion

Compounds **11** and **12** are yellow solids, which are thermally quite stable as evidenced by their high melting points (206 °C and 210 °C respectively) and the presence of molecular ions in their EI mass spectra. **11** is soluble in toluene, THF, and *n*-hexane, while compound **12** is soluble in toluene and THF only. The ^1H and ^{13}C NMR data of **11** and **12** are consistent with the determined single crystal structures. The ^1H NMR spectrum of **11** in C_6D_6 exhibits three singlets (δ 0.28, 1.63, -0.43 ppm) attributed to the proton resonances arising from SiMe_3 , $\eta^5\text{-C}_5\text{Me}_5$, and Zr-Me groups, whereas the respective SiMe_3 and $\eta^5\text{-C}_5\text{Me}_5$ groups in compound **12** resonate at δ 0.3 and 1.73 ppm. The ^{13}C NMR spectrum of **11** reveals two resonances (δ 25.7, 11.3 ppm), which can be assigned to the carbon resonances arising from Zr-Me and Zr-Cp*, respectively. In compound **12** the carbon atoms of the methyl groups (Hf-Cp*) resonate at δ 11.5 ppm. The mass spectrometry data for **11** revealing a signal of low intensity for the molecular ion and the base peak is observed at m/z 494.1 corresponding to $[\text{M-Cp}^*_2\text{Zr}(\text{Me})\text{-OMe}]^+$. In the mass spectrum of compound **12** the base peak is observed at m/z 875.3 representing $[\text{M-Cp}^*]^+$ and the peak with the highest mass corresponds to the molecular ion M^+ (m/z 1010.5).

Scheme 17



Yellow crystals of **11** suitable for single-crystal X-ray diffraction study were obtained from a saturated solution of *n*-hexane at $-30\text{ }^{\circ}\text{C}$ and for those of **12** from a mixture of *n*-hexane/toluene solution at $-30\text{ }^{\circ}\text{C}$. Compounds **11** and **12** crystallize in the monoclinic space group *Pn* with one molecule in the asymmetric unit. Both absolute structures could be determined unequivocally. In **11** the bismuth atom is bonded through an oxygen atom to zirconium and in **12** to hafnium, respectively, and both contain a bent Bi-O-M (M=Zr, Hf) core as revealed by the corresponding bond angles ($138.23(19)^{\circ}$ and $134.55(18)^{\circ}$). Like the other structures, the bismuth atom shows the anticipated trigonal pyramidal coordination geometry with two nitrogen atoms of the 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2$ ligand and one (μ -O) atom. The zirconium and hafnium metal atoms show a distorted tetragonal geometry each (Figures 11 and 12).




Figure 11. Molecular structure of **11**. Anisotropic displacement parameters are depicted at the 50% probability level. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^{\circ}$]: Bi1-O1 2.032(3), Zr1-O1 1.982(3), Zr1-C37 2.292(5), Bi1-N1 2.169(4), Bi1-N2 2.139(4), Zr1-O1-Bi1 138.23(19), N1-Bi1-N2 84.09(16), O1-Zr1-C37 92.07(18), O1-Bi1-N1 98.42(15), O1-Bi1-N2 93.82(15).

The coordination sphere around the transition metals is quite similar: the zirconium metal in **11** accommodates two of the η^5 -coordinated Cp* ligands, the μ -bridging O atom, and one Me group, whereas at the hafnium atom in **12** the latter is replaced by a chlorine atom. The Zr-O bond length (1.982(3) Å) falls between those found in LMeAl(μ -O)ZrRCp₂ (av 1.92 Å)¹⁰⁶ and Ti₄Zr₄O₆(OBu)₄(OMc)₁₆ (OMc= methacrylate, av Zr-O 2.17 Å).¹⁰⁷ In **6** the Hf-O bond length (1.957(3) Å) is longer than that in LAlMe(μ -O)HfMeCp₂ (1.919 Å).^{101c} The Bi-O bond length in compounds **8-12** (av 2.038 Å) is comparable to those observed in (BiTi₂O(O*i*Pr)₉) (2.090 Å)¹⁰⁸ and [(Cp*MoO₃)₂BiPh₃] (2.201 Å)¹⁰⁹ but considerably shorter than that found in [{Bi(Hsal)₃}]₂{Al(acac)₃} (2.765 Å).¹⁰⁰




Figure 12. Molecular structure of **12**. Anisotropic displacement parameters are depicted at the 50% probability level. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1-O1 2.045(3), Hf1-O1 1.957(3), Hf1-Cl1 2.4459(12), Bi1-N1 2.158(4), Bi1-N2 2.139(4), Hf1-O1-Bi1 134.55(18), N1-Bi1-N2 86.63(16), O1-Hf1-Cl1 92.5410, O1-Bi1-N1 98.68(15), O1-Bi1-N2 93.09(14).

2.2.4 Structural Comparison of compounds 8-12

The Bi–N bond distances span a very narrow range of 2.115(4) in **8** to 2.169(4) Å in **11** while the Bi–O distances vary by about 0.08 Å from 1.978(3) in **8** to 2.0552(11) Å in **10**. The shortest Bi–O bond in the aluminum derivative **8** correlates with the widest Bi–O–M angle (160.7(2)° in **8**) while the longer distance in **10** tolerates the most acute angle (112.58(6)°). The angle of the Bi–O bond to the BiN₂ plane is remarkably invariant and widens only by less than 4° from 95.2° in **9** to 98.9° in **11** and **12**, demonstrating that the lone-pair character at the bismuth atom is almost independent from the nature of the second organometallic fragment. The angle of the M–O bond relative to the ML₂ plane varies much more (from 99.1° in **10** to 140.3° in **11**), reflecting the dependence of the angle on the steric requirements of the organometallic fragment.

2.3 Synthesis, Structure and Reactivity of Organobismuth(III), Bismuth Chalcogenide and Dibismuthine Complexes


Organobismuth compounds have attracted interest because of the Lewis acidity and the nontoxicity of the bismuth atom. Trivalent organobismuth compounds with electronegative substituents, such as halogen and oxygen atoms, are well-known to behave as good Lewis acids and to form tetracoordinate structures easily by the coordination with Lewis bases.¹¹⁰ Bismuth halide fragments have been employed as a building block for the formation of two-dimensional coordination networks with tunable conducting/semiconducting properties.¹¹¹ The application of organobismuth reagents in organic synthesis is still very limited.¹¹² Organobismuth compounds are normally nontoxic and potentially meet the recent requirement for the reagent, process, and product of little or no risk to humans and environment, which is a major contemporary concern in the chemistry community. The unique performance of bismuth salts as Lewis acid catalysts or reagents is becoming

evident¹¹³, and organobismuth(III) has been used in the oxidation of alcohols to aldehydes and ketones under mild conditions.¹¹⁴ Herein we report the preparation and stabilization of organobismuth(III) compounds, low-valent bismuthane and organobismuth chalcogenides.

2.3.1 Synthesis, Spectroscopic Characterization and Structure of Complexes [1,8-C₁₀H₆(NSiMe₃)₂]BiR (R= Cl(**13**), CCPh(**14**), Cp*(**15**), Me(**16**), OPh(**17**), NH₂(**18**))

Metathesis of amide to chloride can be achieved by treatment of *n*-hexane solution of **1** with ClSiMe₃. Organic groups can be easily introduced in **1** by elimination of HNMe₂ using acidic reagents. Treatment of **1** with stoichiometric amounts of PhCCH, Cp*H and PhOH in *n*-hexane results in the formation of LBiCCPh (**14**), LBiCp*(**15**), and LBiOPh (**17**) [L= 1,8-C₁₀H₆(NSiMe₃)₂], respectively under elimination of HNMe₂ as shown in Scheme 18. LBiNH₂

Scheme 18



(**18**) was prepared as an oily compound by bubbling dry ammonia gas through the *n*-hexane solution of **1**, whereas LBiMe (**16**) can be synthesized by slow addition of AlMe₃ in *n*-hexane to the *n*-hexane solution of **1**. The ¹H NMR spectrum of each reaction mixture of compounds **13-18** shows almost quantitative conversion of the precursors to products, as revealed by the absence of NMe₂ resonance. The ¹H NMR spectra of compounds **13-18** in C₆D₆ at room temperature show singlet resonances for SiMe₃ groups. The ¹H NMR spectrum of **15** displays one resonance at δ 1.86 ppm which can be attributed to the Me protons of Cp*, whereas the

particular BiMe group in **16** resonates as a singlet at δ 0.43 ppm. In compound **18** a broad singlet resonance at δ 5.38 ppm can be assigned to the protons of the NH₂ group. The ¹³C NMR spectrum of **15** reveals two resonances (δ 10.4, 3.3 ppm), which can be assigned to the carbon resonances arising from SiMe₃ and Bi-Cp*, respectively. Compound **13** is a red crystalline solid which is soluble in common organic solvents such as toluene and THF, and it is stable in the solid state at room temperature for several months under an inert atmosphere. Organobismuth compounds **14-17** are orange crystalline solids which are soluble in toluene, THF, and *n*-hexane. Compounds **14-18** are air and moisture sensitive and hydrolyze upon exposure to air.




Figure 13. Molecular structure of **13**. Thermal ellipsoids are set at 50% probability level. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^\circ$]: Bi1–N1 2.1569(18), Bi1–N2 2.127(2), Bi1–Cl 2.4784(6), Si1–N1 1.735(2), Si2–N2 1.745(2), N1–Bi1–N2 83.88, Cl–Bi1–N1 92.10(6), Cl–Bi1–N2 95.72(6), Si1–N1–Bi1 119.61(10), Si2–N2–Bi1 118.80(10).

2.3.2 Molecular Structure Description of Compounds 13-16

Single crystals suitable for X-ray structures of **14-16** were obtained from saturated solution of *n*-hexane at -30 $^\circ\text{C}$ and for those of **13** from a mixture of *n*-hexane/toluene solution at -30 $^\circ\text{C}$. The coordination geometry around the bismuth atom in compounds **13-16**

is that of a distorted trigonal-pyramidal arrangement with a stereochemically active electron lone pair. Compound **13** crystallizes in the tetragonal space group $I\bar{4}$ with one molecule in the asymmetric unit (Figure 13). The Bi–Cl distance (2.4784(6) Å) is shorter than those that have been reported for complexes [2,6-(Me₂NCH₂)₂C₆H₃]₂-BiCl (2.6086(13) Å)¹¹⁵ and [{HC(Et₂NCH₂CH₂NCMe)₂}]BiCl₂] (av 2.7055 Å)¹¹⁶ but is comparable to that of characteristic Bi–chlorine covalent bond ($\Sigma_{\text{cov}}(\text{Bi-Cl})$ 2.51 Å).¹¹⁷ Compounds **14** and **16** crystallize in the monoclinic space group $P2_1/n$ with two molecules in the asymmetric unit (Figures 14, 16).




Figure 14. Molecular structure of **14**. Thermal ellipsoids are set at 50% probability level. Only one of the two crystallographically independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi2–N1 2.142(2), Bi2–N2 2.145(2), Bi2–C17 2.212(3), C17–C18 1.202(4), Si1–N1 1.733(2), Si2–N2 1.738(2), N1–Bi2–N2 84.88(8), C17–Bi2–N1 89.81(9), C17–Bi2–N2 94.91(9), Si1–N1–Bi2 119.55(12), Si2–N2–Bi2 120.86(11).

The Bi-C bond length (av 2.2115 Å) in **14** is shorter than that in [tBuN(CH₂C₆H₄)₂BiCCPh] (2.289(4) Å).⁹⁵ The Bi-Me distance in **16** (av 2.2485 Å) falls between those found in [2,6-(Me₂NCH₂)₂C₆H₃](Me)BiI (2.224(10) Å)¹¹⁵ and [tBuN(CH₂C₆H₄)₂BiMe] (2.264(6) Å).⁹⁵ Compound **15** crystallizes in the triclinic space group $P\bar{1}$ with two molecules in the asymmetric unit (Figure 15). The Bi-C bond length pattern with one short, two intermediate and two longer distances (av 2.3638, 2.7616, 2.7745, 3.1900, 3.1998 Å) between Cp* and bismuth resembles a scenario between an η^1 or η^3 chelating mode. The Bi-XB_{1A_B} (XB_{1A_B} = centroid of the Cp* ring) distance (2.6059 Å) in **15** is longer than those in [(C₅HR₄)BiCl₂]₂ (R=CHMe₂) (Bi-X_{1A}, av 2.3495 Å).¹¹⁸ In compounds **13-16** the Bi-N bond lengths are in the range of 2.127(2) Å (in **13**) to 2.1995(16) Å (in **15**) and the N-Bi-N angles are ranging from 82.47(7)° (in **16**) to 84.88(8)° (in **14**), which are comparable to those observed in **1**.




Figure 15. Molecular structure of **15**. Thermal ellipsoids are set at 50% probability level. Only one of the two crystallographically independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi2-N1 2.1854(16), Bi2-N2 2.1568(15), Bi1-X_{1A} 2.62137(10), Si1-N1 1.7373(16), Si2-N2 1.7285(17), N1-Bi2-N2

83.46(6), X1A–Bi2–N1 123.53(4), X1A–Bi2–N2 118.55(4), Si1–N1–Bi2 117.64(8), Si2–N2–Bi2 117.65(8). (X1A = centroid of the Cp* ring)




Figure 16. Molecular structure of **16**. Thermal ellipsoids are set at 50% probability level. Only one of the two crystallographically independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Bi1–N1 2.1561(19), Bi1–N2 2.1733(19), Bi1–C17 2.249(3), Si1–N1 1.737(2), Si2–N2 1.745(2), N1–Bi2–N2 83.69(8), C17–Bi1–N1 93.32(9), C17–Bi1–N2 94.75(9), Si1–N1–Bi1 121.10(10), Si2–N2–Bi1 118.45(8).

2.3.3. Synthesis, Spectroscopic Characterization and Structure of Organobismuth Chalcogenide Complex: Synthesis of [1,8-C₁₀H₆(NSiMe₃)₂Bi]₂S (**19**)


The first bis(diorganobismuth) chalcogenides, (R₂Bi)₂E (E = O, S, Se, Te) were prepared several years ago by chalcogen insertion into bismuth–bismuth bonds.^{119,120} Later also reactions between diorganobismuth halides and sodium chalcogenides were accomplished.¹²¹ The examples with known crystal structures are the mesityl derivatives (R₂Bi)₂E [E = O, S,

2. Results and Discussion

Se; R=2,4,6-Me₃C₆H₂],¹²² [2,6-(Me₂NCH₂)₂C₆H₃BiS]₂,⁴⁵ (R₂Bi)₂E [E = S, Te; R=CH(SiMe₃)₂]¹²³ and [2,6-(*t*-BuOCH₂)₂C₆H₃BiS]₂.¹²⁴

The reaction of **1** in the presence of equivalent amounts of elemental sulfur at room temperature in *n*-hexane leads after three days to compound **19** in good yield (Scheme 19). Compound **19** is an orange crystalline solid, which is soluble in organic solvents such as *n*-hexane, THF, and toluene. The ¹H NMR spectrum of **19** exhibits two different resonances (δ 0.169 and 0.131 ppm) for the SiMe₃ groups.

Scheme 19



In **19** the bismuth atoms are bound by a bent Bi(μ -S)Bi core as revealed by the bond angle of av 113.18°. The bismuth-sulfur distances in **19** (av 2.5383 Å) are similar to those in [{2-(Me₂NCH₂)C₆H₄}₂Bi]₂S (2.5558(17) Å)⁴⁵ and (Mes₂Bi)₂S (av 2.5325 Å)¹²¹, while the Bi(μ -S)Bi angle in **19** is considerably larger when compared with those of the latter compounds (98.17(8)°, 98.7(2)°).




Figure 17. Molecular structure of **19**. Thermal ellipsoids are set at 50% probability level. Only one of the two crystallographically independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^\circ$]: Bi1-N1 2.167(2), Bi1-N2 2.138(2), Bi2-N3 2.144(2), Bi2-N4 2.170(2), N1-Si1 1.743(2), N2-Si2 1.733(2), N3-Si3 1.725(2), N4-Si4 1.751(2), Bi1-S1 2.5386(7), Bi2-S1 2.5381(7), Bi1-S1-Bi2 113.47(3), N1-Bi1-N2 84.68(8), N3-Bi2-N4 84.00(9), N1-Bi1-S1 92.10(6), N2-Bi1-S1 95.10(6), N3-Bi2-S1 94.96(6), N4-Bi2-S1 98.22(6), Bi1-S1-Bi2 112.89(3).


2.3.4. Synthesis, Spectroscopic Characterization and Structure of Low-valent Dibismuthine Complex: Synthesis of $[1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}]_2$ (**20**)

There are very few compounds with known crystal structures of compounds containing bismuth-bismuth bonds. Interesting aspects of the structures of dibismuthines are the conformation and the possibility of the formation of bismuth chains through alternating Bi-Bi bonds and intermolecular Bi \cdots Bi contacts. The first crystallographically characterized tetraorganyl derivative of bismuth(II) is tetraphenyldibismuthine, $\text{Ph}_2\text{Bi-BiPh}_2$.¹²⁵ The molecule has a staggered trans conformation. The coordination at the bismuth atoms is

essentially trigonal pyramidal. The bond angles about bismuth are significantly smaller than the tetrahedral value. The dibismuthine $(\text{Me}_3\text{Si})_2\text{Bi-Bi}(\text{SiMe}_3)_2$, which does not contain a direct Bi-C bond, has the same molecular structure as tetraphenyldibismuthine with a staggered trans conformation of the substituents.¹²⁶ However, the crystal structure is different. There are linear zigzag bismuth chains with alternating Bi-Bi bonds and shortened intermolecular $\text{Bi}\cdots\text{Bi}$ contacts in the crystalline state. The shortest Bi-Bi single bond (2.947 Å) was found in the unusual bismuth-arene adduct, $(\mu\text{-}\eta^6\text{-C}_6\text{Me}_6)[\text{Bi}_2(\mu\text{-O}_2\text{CCF}_3)_4]_2$,¹²⁷ which is the first example of a carboxylato derivative containing the bismuth atoms in a formal +2 oxidation state.

Reaction of PhSiH_3 with **1** in *n*-hexane results in a Bi-Bi bond formation (Scheme 20). Compound **20** is obtained as a red crystalline solid which is air and moisture sensitive but stable in the solid state under an inert atmosphere at room temperature. In the ^1H NMR spectrum of **20** the 18 protons of SiMe_3 resonate as a singlet (δ 0.15 ppm) and the aromatic protons appear in the range of δ 7.13 - 7.22 ppm.

Scheme 20



Compound LBi-BiL (**20**) crystallizes in the monoclinic space group $P2_1/c$ with one molecule in the asymmetric unit (Figure 18). The N,N'-disubstituted 1,8-diaminonaphthalene ligands of **20** are in a *trans* conformation to each other and the N-Bi-Bi angles lie in the range

between av 89.88° and av 89.03° . The Bi-Bi bond distance of $3.0197(2) \text{ \AA}$ corresponds to a normal bismuth-bismuth single bond as found in $(\text{Me}_3\text{Si})_4\text{Bi}_2$ ($3.035(3) \text{ \AA}$)¹²⁶ or R_4Bi_4 ($\text{R}=(\text{Me}_3\text{Si})_2\text{CH}$; av 3.005 \AA)⁵⁶.




Figure 18. Molecular structure of **20**. Thermal ellipsoids are set at 50% probability level. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^\circ$]: Bi(1)-N(1) 2.1648(11), Bi(1)-N(2) 2.1660(12), Bi1A-N1A 2.1621(13), Bi1A-N2A 2.1646(11), N1-Si1 1.7432(12), N2-Si2 1.7422(13), N1A-Si1A 1.7470(13), N2A-Si2A 1.7475(12), Bi1-Bi1A 3.0197, N1-Bi1-N2 84.51(5), N1A-Bi1A-N2A 85.02(4), N1-Bi1-Bi1A 89.82(3), N2-Bi1-Bi1A 88.87(3), N1A-Bi1A-Bi1 89.95(3), N2A-Bi1A-Bi1 89.19(3).

2.4 Synthesis, Characterization, and X-ray Single Crystal Structures of *N,N'*-Bis(trimethylsilyl)diphenyliminophosphonamide Aluminum Halide, Hydride, Amide and Alkylate.

Monoanionic *N,N'*-chelating organic ligands such as β -diketiminato, guanidinate and amidinate have attracted growing interest in organometallic chemistry in the last decade due to their capability to coordinate very flexible to the metal center as monodentate (η^1), chelating (η^2) or bridging monodentate (μ - η^1 - η^1) four-electron donor.¹²⁸ One of the most fruitful systems has been the $[\text{R-N-E}(\text{R}')_n\text{N-R}]^-$ ($\text{E}=\text{C}, \text{S}, n=1; \text{Si}, \text{P}, n=2$) ligand coordinating with two nitrogen atoms. Moreover, they were found very useful in catalysis, material science (i.e. precursors for CVD) and organic-inorganic hybrids since their steric and electronic properties can easily be tuned by modification of the organic substituents R and R'.


A number of PN_2 complexes of the metals have been prepared and characterized;¹²⁹⁻¹³² many of these compounds were synthesized by the reaction of the $[\text{Ph}_2\text{P}(\text{NSiMe}_3)_2]^-$ ligand with the corresponding metal chloride. Monosubstituted complexes are generally accessible via this route and are typically dinuclear with halide bridges. Amine elimination is also a complementary route for the preparation of both mono- and disubstituted PN_2 complexes.¹³³ Herein, we report the synthesis and solid-state structures of aluminum complexes with the $\text{Ph}_2\text{P}(\text{NSiMe}_3)_2^-$ ligand.

2.4.1 Synthesis of $\text{Ph}_2\text{P}(\text{NSiMe}_3)_2\text{Li}$ (22**)**

Compound $\text{Ph}_2\text{P}(\text{NSiMe}_3)_2\text{H}$ (**21**) was prepared by a previously reported method.¹³⁴ Treatment of iminophosphonamide with MeLi or *n*BuLi yielded the lithium salt of iminophosphonamide L^2Li (**22**) ($\text{L}^2=\text{Ph}_2\text{P}(\text{NSiMe}_3)_2$) in high yield (Scheme 21).¹³⁵ The ^1H NMR spectrum of **22** exhibits one singlet at δ 0.12 ppm which can be attributed to SiMe_3

groups while the aromatic protons appear between δ 6.98 and 7.80 ppm. **22** shows a resonance at δ 7.4 ppm in its ^{31}P NMR spectrum.

Scheme 21




2.4.2. Reaction of $L^2\text{Li}$ with RAlCl_2 : Synthesis of $\text{Ph}_2\text{P}(\text{NSiMe}_3)_2\text{AlRCl}$ ($\text{R} = \text{Cl}$, **23**; Me , **24**)

Synthesis of $L^2\text{AlCl}_2$ (**23**) (Scheme 22) was accomplished by reacting the lithiated precursor **22** with AlCl_3 under elimination of LiCl in moderate yield. The solution of AlCl_3 in toluene was added drop by drop to the solution of **22** in a 1:1 stoichiometric ratio in toluene, and stirred at 25 °C for 14 h which yields the colorless complex **23**. In a similar route complex **24** was prepared by slow addition of AlMeCl_2 (1.0 M in *n*-hexane) to the toluene solution of **22**, and stirring at room temperature for 14 h. Compounds **23** and **24** are soluble in common organic solvents such as toluene, THF, and *n*-hexane, and they are stable in the solid state at room temperature for several months under an inert atmosphere. **23** is easily hydrolyzed on exposure to air as indicated by a color change from colorless to a light yellow oily solid, and we were not able to measure elemental analysis of **23** due to its sensitivity. The ^{31}P NMR spectra of **23** and **24** exhibit downfield shifts when compared with those of complex **22** (δ 37.8 ppm for **23**, 33.2 ppm for **24**, 7.4 ppm for **22**). The ^1H NMR of **24** exhibits a single

resonance (δ 0.04 ppm) which can be assigned to Al-Me. The 18 protons of SiMe₃ resonate as a singlet at δ -0.03 ppm and the aromatic protons of Ph are shown at δ 7.46 - 7.79 ppm.

Scheme 22



Complexes **23** and **24** were characterized further by X-ray crystallography, and the molecular structures are presented in Figures 21 and 22. Compound [Ph₂P(NSiMe₃)₂AlCl₂] (**23**) crystallizes in the triclinic space group $P\bar{1}$ with two molecules in the asymmetric unit whereas [Ph₂P(NSiMe₃)₂AlMeCl] (**24**) is monomeric in the solid state and crystallizes in the monoclinic space group $P2_1/c$. Colorless crystals of **23** and **24** suitable for X-ray structure analysis were grown from toluene and *n*-hexane at -78 °C respectively. In both structures the aluminum atoms show a distorted tetrahedral environment, and the metrical parameters associated with the PN₂ ligand are quite similar.¹³⁶ All N atoms bearing SiMe₃ groups show a planar arrangement; the Si-N bonds are slightly shorter than the sum of the covalent radii of 1.87 Å¹³⁷, and the P-N distances in the range of 1.614(12)-1.629(16) Å have been previously rationalized to range between single and double P-N bonds.¹³⁸ However, recent experimental charge density studies proved the shortening mainly to be caused by electrostatic contributions.¹³⁹ In addition to the P-N σ -bonds, there is Coulomb attraction between the positively charged phosphorus atoms and the negatively charged nitrogen atoms, and in return, repulsion between the competing N-atoms across the central P-atom. This ‘seesaw’

effect is most prominent in the structures from the tetra(amino) phosphonium cation, $[\text{P}(\text{NHPh})_4]^+$, to the tetra(imino) phosphate trianion, $[\text{P}(\text{NPh})_4]^{3-}$.




Figure 21. Molecular structure of **23**. Anisotropic displacement parameters are drawn at the 50 % probability level. Only one of the two crystal independent molecules is shown. H atoms are omitted for clarity. Selected bond lengths [\AA] and angles [$^\circ$]: Al1-N1 1.880(17), Al1-N2 1.885(17), Si1-N1 1.739(16), Si2-N2 1.747(17), P1-N1 1.629(16), P1-N2 1.623(16), Al1-Cl1 2.132(8), Al1-Cl2 2.130(8), Cl1-Al1-Cl2 109.51(3), Al1-N1-P1 90.59(8), Al1-N2-P1 90.57(8), N1-P1-N2 97.65(8), N1-Al1-N2 81.11(7), C1-P1-C7 107.16(9), Si1-N1-Al1 132.34(9), Si1-N1-P1 132.44(10), Si2-N2-Al1 132.11(9), Si2-N2-P1 137.02(10), Cl1-Al1-N1 114.04(6), Cl2-Al1-N1 116.42(6), Cl1-Al1-N2 115.83(6), Cl2-Al1-N2 117.67(6).

Despite the diversity of the P–N bond lengths, the average P–N bond length within the PN_4 tetrahedra is very similar ($\Sigma(d_{\text{P-N}})/4 = 1.62(1)\text{\AA}$).¹⁴⁰ On this basis we state that there is no P=N double bonding, and hence no hypervalent phosphorus present in the iminophosphonamide ligands presented in this paper.¹⁴¹

The Al-Cl bond lengths in compound **23** are shorter than that of compound **24**, while the ligand bite (N1-Al1-N2) in **23** (av 81.09°) is slightly bigger than that in **24** (80.04(5)°). The Al-Cl distances in **23** (av 2.129 Å) are quite similar to those observed in the mono 1-aza-allyl complex [N(SiMe₃)C(Ph)C(SiMe₃)₂]AlCl₂ (av 2.071 Å) but the core angle is significantly different due to non equivalent chelating backbone.¹⁴²





Figure 22: Molecular structure of **24**. Anisotropic displacement parameters are drawn at the 50 % probability level. H atoms (except for those of C19) and half toluene solvent molecule are omitted for clarity. Selected bond lengths [Å] and angles [°]: Al1-N1 1.899(13), Al1-N2 1.909(13), Si1-N1 1.736(12), Si2-N2 1.733(12), P1-N1 1.614(12), P1-N2 1.617(12), Al1-Cl1 2.155(6), Al1-C19 1.947(15), Cl1-Al1-C19 111.89(6), Al1-N1-P1 90.89(6), Al1-N2-P1 90.41(6), N1-P1-N2 98.56(6), N1-Al1-N2 80.04(5), C1-P1-C7 106.846(7), Si1-N1-Al1 131.52(7), Si1-N1-P1 136.69(8), Si2-N2-Al1 128.83(7), Si2-N2-P1 134.17(8), Cl1-Al1-N1 110.81(4), C19-Al1-N1 120.92(7), Cl1-Al1-N2 109.69(4), C19-Al1-N2 119.69(6).

2.4.3. Reaction of L^2H with $AlH_3 \cdot NMe_3$: Synthesis of $[Ph_2P(NSiMe_3)_2]_2AlH$ (**25**)

Organoaluminum hydrides of low aggregation have proved to be effective reagents for preparing chalcogenide compounds with elemental chalcogens or organochalcogenides,¹⁴³ such as $LAl(SeH)_2$, $LAl(SeH)Se(SeH)AlL$ ($L = HC(CMeNAr)_2$, $Ar = 2,6\text{-}iPr_2C_6H_3$),¹⁴⁴ $LAl(SH)_2$ ($L = HC(CMeNAr)_2$, $Ar = 2,6\text{-}iPr_2C_6H_3$),¹⁴⁵ $Al_4Se_5(H)_2(NMe_3)_4$.¹⁴⁶ As an extension of this type of reaction, we explored iminophosphonamide **21** to react with $AlH_3 \cdot NMe_3$ to afford an aluminum monohydride $[Ph_2P(NSiMe_3)_2]_2AlH$ (**25**) under elimination of hydrogen.

Reaction of iminophosphonamide **21** with $AlH_3 \cdot NMe_3$ in 2:1 stoichiometric ratio in *n*-hexane/THF at -78 °C afforded the iminophosphonamide aluminum complex $[Ph_2P(NSiMe_3)_2]_2AlH$ (**25**) in moderate yield (Scheme 23). The 1H NMR spectrum of **25** exhibits a broad singlet at δ 5.1 ppm assigned to the AlH and two singlets at δ -0.22 and -0.13 ppm, which can be assigned to the 36 protons of the $SiMe_3$ units. The presence of two non equivalent $SiMe_3$ groups indicates a restricted fluctuonality of these groups at room temperature. In the IR spectrum one weak absorption for the AlH stretching frequency is observed at 1853 cm^{-1} . ^{31}P NMR spectrum of **25** exhibits two single resonances at δ 29.2 and 34.6 ppm. **25** is stable in the solid state under an inert atmosphere and is slowly oxidized and hydrolyzed on exposure to air.

Scheme 23



Complex **25** was characterized further by X-ray crystallography and the molecular structure is presented in Figure 23. The X-ray structure reveals that in **25** the aluminum is five coordinate and in a distorted trigonal bipyramidal geometry with N1 and N4 in axial positions and N2, N3, and H in the equatorial plane. The N1-Al1-N4 axis is almost linear [N1-Al1-N4 175.80(5)°] in which the Al-N bond lengths of 2.08(13) Å for Al1-N1 and 2.100(13) for Al1-N4 are almost the same. The equatorial Al1-N2 and Al1-N3 having almost the same bond lengths (Al1-N2 1.94(13) Å, Al1-N3 1.93(13) Å). The deviation from the ideal trigonal-bipyramidal geometry can be described by an angle of 73.94(26)° between the least-squares plane of N1, Al, N4, H and the equatorial plane. Clearly, the deviation from ideal trigonal-bipyramidal geometry arises from the steric requirements of the two chelating ligands. The axial Al-N bond lengths are in good agreement with those which are observed in similar five coordinate aluminum complexes¹⁴⁷ where Al-N bond lengths are found in the range from 2.051(18) Å to 2.179(7) Å.





Figure 23. Molecular structure of **25**. Anisotropic displacement parameters are drawn at the 50 % probability level. H atoms (except for those of Al1) and one THF solvent molecule are omitted for clarity. Selected bond lengths [Å] and angles [°]: Al1-N1 2.080(13), Al1-N2 1.947(13), Si1-N1 1.7252(13), Si2-N2 1.7381(13), P1-N1 1.5923(13), P1-N2 1.6162(12),

Al1-N3 1.939(13), Al1-N4 2.100(13), N3-Al1-N4 75.66 (5), Al1-N1-P1 89.23(6), Al1-N2-P1 93.90(6), N1-P1-N2 101.10(6), N1-Al1-N2 75.75(5), C1-P1-C7 106.58(7), Si1-N1-Al1 133.46(7), Si1-N1-P1 133.13(8), Si2-N2-Al1 135.69(7), Si2-N2-P1 128.08(8), N3-Al1-N1 102.38(5), N4-Al1-N1 175.80(5), N3-Al1-N2 110.34(6), N4-Al1-N2 101.33(5).

2.4.4. Reaction of L^2H with $AlEt_3$ and $[Al(NMe_2)_3]_2$: Synthesis of $Ph_2P(NSiMe_3)_2AlEt_2$ (26**) and $Ph_2P(NSiMe_3)_2Al(NMe_2)_2$ (**27**)**

Compound **26** was prepared as a white crystalline solid from the reaction of iminophosphonamide **21** with $AlEt_3$ in *n*-hexane at -78 °C. Reaction of **21** with $[Al(NMe_2)_3]_2$ in *n*-hexane at reflux temperature for 6 h afforded the aluminum amide complex **27** (Scheme 24). Both compounds are soluble in common organic solvents such as *n*-hexane, toluene, and tetrahydrofuran. **27** is stable in the solid state under an inert atmosphere and is slowly hydrolyzed on exposure to air whereas **26** is not stable and very sensitive to air. Therefore we were not able to determine the elemental analysis. ^{31}P NMR spectra of **26** and **27** exhibit single resonances at δ 30.3 and 28.6 ppm, respectively. In both compounds the 1H NMR and ^{29}Si NMR spectra exhibit only one singlet for the $SiMe_3$ groups. In compound **27** the 12 protons correspond to NMe_2 and resonate at δ 3.06 ppm while the aromatic protons appear between δ 7.03 and 7.83 ppm.

Scheme 24



Colorless compound **27** crystallizes in the monoclinic space group $P2_1/n$, with one monomer in the asymmetric unit from *n*-hexane at -78 °C. The coordination polyhedron around the aluminum atom features a distorted tetrahedral environment and the metrical parameters associated with the PN_2 ligand are quite similar to those of **22** and **23**. The Al-N3 and Al-N4 bond distances are comparable to those reported in structurally characterized Al-bound terminal NMe_2 groups.¹⁴⁸ The Al-N1 and Al-N2 bond lengths (1.937(12)–1.950(12) Å) are longer than the Al-N3 and Al-N4 distances (1.790(13)–1.795(12) Å), due to the electronically delocalized PN_2 ligand.




Figure 24. Molecular structure of **27**. Anisotropic displacement parameters are drawn at the 50 % probability level. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Al1-N1 1.937(12), Al1-N2 1.950(12), Si1-N1 1.727(12), Si2-N2 1.728(12), P1-N1 1.614(11), P1-N2 1.606(12), Al1-N3 1.795(12), Al1-N4 1.790(13), N3-Al1-N4 111.61(6), Al1-N1-P1 91.38(6), Al1-N2-P1 91.14(6), N1-P1-N2 99.15(6), N1-Al1-N2 78.21(5), C1-P1-C7 104.23(7), Si1-N1-Al1 131.04(6), Si1-N1-P1 133.52(7), Si2-N2-Al1 130.33(6), Si2-N2-P1 133.57(7), N3-Al1-N1 113.12(6), N4-Al1-N1 118.87(6), N3-Al1-N2 119.33(6), N4-Al1-N2 112.27(6).

2.4.5 Structural Comparison of LAiCl_2 (**23**), LAiMeCl (**24**), L_2AlH (**25**), and $\text{LAl}(\text{NMe}_2)_2$ (**27**)

The superposition of the PN_2Al four-membered rings in the four structures in Figure 25 reveals that not only this motif remains widely unchanged but even the phenyl groups adopt the same orientation. Also the remaining substituents at aluminum stay almost in place. Remarkably, only one of the two Me_3Si groups responds to the various steric requirements of those two exocyclic substituents. Switching one of the chlorine substituents in **23** to a methyl group in **24** leaves the angle sum at the nitrogen atom virtually unchanged (359.7 vs 359.1°) and the silyl atom almost shares the plane with the PN_2Al four-membered ring. Only the second iminophosphonamide ligand in **25** and the two dimethylamino groups in **27** remove that silyl group from the plane, indicated by an angle sum of only 356.16 and 355.94° , respectively.





Figure 25. Superposition of **23**, **24**, **25**, and **27**, depicting the response of the trimethylsilyl group at N1 to the bulk of the two other substituents at the aluminum atom.

3. Summary and Outlook


3.1. Summary

Organometallic molecules with useful functionalities have gained tremendous attention. Substitution of functionalities leads to the synthesis of a variety of other new derivatives, and condensation in a controlled manner with suitable metal synthons affords a route to homo- and heterobimetallic complexes which are mostly useful as catalysts.

The first part of the thesis deals with the synthesis of bismuth amide complex and its reaction with compounds containing unsaturated C-O, C-C, and C-N bonds. The main motivation for the research in the field of the bismuth chemistry is, without any doubt, the role of bismuth compounds in catalysis and due to their non toxicity in pharmacology. $L^1\text{BiNMe}_2$ (**1**) was prepared by the reaction of 1,8-bis[(trimethylsilyl)amino]naphthalene with $\text{Bi}(\text{NMe}_2)_3$. To investigate the reactivity of bismuth-nitrogen bond toward carbon-oxygen bond we treated **1** with aldehyde and keton. The reaction of **1** with 2-benzoylpyridine and 3-pyridinecarboxaldehyde proceeds smoothly at room temperature to give addition products in which 2-benzoylpyridine and 3-pyridinecarboxaldehyde insert into the Bi-N bond of **1** under formation of **2** and **3**, respectively.




Molecular Structure of 2




Molecular Structure of 3

3. Summary and Outlook

To extend those findings to carbon-carbon bonds we reacted **1** with alkene and alkyne. Compounds **4** and **5** were obtained by reaction of **1** with 2-methyl-2-propenenitrile and diethyl acetylene dicarboxylate, respectively. During the formation of **5** the exchange of one of the OEt groups by a NMe₂ substituent occurred.




Molecular Structure of 4




Molecular Structure of 5

Compounds **6** and **7** are the products of bismuth-nitrogen bond insertion to carbon-nitrogen bonds. Reaction of **1** with diisopropylcarbodiimide and phenylisocyanate results in the formation of **6** and **7**, indicating that carbodiimide and isocyanate molecules are inserted into the Bi-N bond, respectively.




Molecular Structure of 6




Molecular Structure of 7

3. Summary and Outlook


Due to the basic character of the NMe₂ group, compound **1** was utilized to prepare a series of heterobimetallic oxides. The second part of the thesis describes the efforts to synthesize oxygen-bridged heterobimetallic complexes of bismuth with main group and transition metals which are difficult to achieve by other methods. The bismuthamide (**1**) complex acts as a building block for the preparation of the first structurally characterized oxygen bridged heterobimetallic complexes of groups 4, 13 and 14 metals. Thus, the reaction of L¹BiNMe₂ (**1**) with LM(Me)(OH) and LGe(OH) (M=Al, Ga, L= CH((CMe)NAr)₂, Ar = 2,6-*i*Pr₂C₆H₃), provides the complexes L¹Bi(μ-O)MMeL (M = Al **8**; Ga, **9**) and L¹Bi(μ-O)GeL (**10**).



Molecular Structure of 8




Molecular Structure of 9




Molecular Structure of 10

The study of the half-metallocenes of zirconium and hafnium bearing terminal methyl and chloride groups is limited because of their thermal and kinetic instability. A series of heterobimetallic complexes of half-metallocenes bearing terminal methyl and chloride groups have been prepared. The transition metal hydroxides $\text{Cp}^*_2\text{ZrMe}(\text{OH})$ and $\text{Cp}^*_2\text{HfCl}(\text{OH})$ react with **1** to generate the corresponding complexes $\text{L}^1\text{Bi}(\mu\text{-O})\text{ZrMeCp}^*_2$ (**11**) and $\text{L}^1\text{Bi}(\mu\text{-O})\text{HfClCp}^*_2$ (**12**) under elimination of HNMe_2 . X-ray structural analysis reveals the presence of bent $\text{Bi}(\mu\text{-O})\text{Zr}$ and $\text{Bi}(\mu\text{-O})\text{Hf}$ cores.




Molecular Structure of 11



Molecular Structure of 12


From the point of view of the applications in catalysis and organic synthesis the organobismuth(III) complexes play a very important role. Under given reaction conditions the amide group can be selectively substituted or replaced by another similar or different functional group. Thus, the reaction of L^1BiNMe_2 (**1**) with ClSiMe_3 smoothly yields the corresponding chloride L^1BiCl (**13**).




Molecular Structure of 13

3. Summary and Outlook


Treatment of **1** with HCCPh, Cp*H, AlMe₃, PhOH and NH₃ resulted in the formation of the organobismuth(III) complexes L¹BiCCPh (**14**), L¹BiCp*(**15**), L¹BiMe (**16**), L¹BiOPh (**17**) and L¹BiNH₂ respectively. The molecular structures of **14**, **15** and **16** have been determined by X-ray structure analysis. Compound **15** is an example of a half-sandwich bismuth(III) complex with bonding to Cp* in an η¹ or η³ chelating mode. LBiMe (**16**) can be synthesized by slow addition of AlMe₃ in *n*-hexane to the *n*-hexane solution of **1**.



Molecular Structure of 14




Molecular Structure of 15




Molecular Structure of 16

Reaction of **1** with elemental sulfur and PhSiH₃ affords a bimetallic derivative of composition (L¹Bi)₂S (**19**) with a bent Bi(μ-S)Bi core and a low valent dibismuthine (L¹Bi)₂ (**20**) complex, respectively.




Molecular Structure of 19




Molecular Structure of 20

The last part of the thesis deals with the synthesis of four-membered aluminum rings bearing halide, methyl, and hydride substituents. Aluminum hydrides, alkylates, and halides are involved in most of aluminum containing reactions and they are extraordinary important for aluminum chemistry

Iminophosphonamide aluminum dichloride and methylchloride L^2AlCl_2 (**23**) and $L^2AlMeCl$ (**24**) ($L^2=Ph_2P(NSiMe_3)_2$) were easily obtained from the reactions of L^2Li and selected simple aluminum halide precursors.



Molecular Structure of 23




Molecular Structure of 24


3. Summary and Outlook

The reaction of the very bulky iminophosphonamide ligand L^2H with $AlH_3 \cdot NMe_3$, $AlEt_3$ and $[Al(NMe_2)_3]_2$ afforded a dimeric aluminum monohydride L^2_2AlH (**25**), an aluminum ethyl L^2AlEt_2 (**26**) and an aluminum amide L^2AlNMe_2 (**27**) complex, respectively.

In **25** the aluminum is five coordinate and in a distorted trigonal bipyramidal geometry. It could be a very useful precursor for the preparation of aluminum chalcogenide complexes.



Molecular Structure of 25



Molecular Structure of 27

3.2. Outlook

The thesis presented here has focussed on generating functionalities on bismuth and aluminum centers such as NMe_2 , halide, alkyl and NH_2 and studying their reactivity. This resulted in the development of new synthetic strategies for generating such species.

Extension of this work may be the application of the carbene method for conversion of metal halides to the corresponding hydroxides and amides and as possible starting materials for reduction with the aim of obtaining metals in low oxidation states, and also for studying their involvement in different metathesis reactions. In summary, generation of such hydroxides has a huge synthetic potential in the field of well-defined heterobimetallic oxides. Moreover, other p-, d- or f- block metals are potential candidates for the generation of heterobimetallic complexes containing Bi-O-M and Al-O-M cores.

4. Experimental Section

4.1. General Procedures

All experiments were carried out under an atmosphere of dry nitrogen or argon using Schlenk techniques or inside a glove-box filled with dry nitrogen, where the O₂ and H₂O level were strictly maintained below 1 ppm. Toluene, benzene, hexane, pentane, tetrahydrofuran, and diethylether were purified with the M-Braun solvent drying system. Dichloromethane (CaH₂) were dried and distilled prior to use. The samples for analytical measurements were prepared inside the glove box

4.2. Physical Measurements

The *melting points* of all compounds described in this thesis were measured on a Büchi B-540 apparatus in sealed capillaries and are uncorrected.

¹H, ¹³C, ²⁷Al, ²⁹Si, and ³¹P NMR spectra were recorded on Bruker-Avance-500, Avance-300 and Bruker-Avance-200 instruments. Deuterated NMR solvents C₆D₆, C₇H₈, and THF-D₈ were dried by stirring for 2 days over Na/K alloy followed by distillation *in vacuo* and degassed. The chemical shifts are reported in ppm with reference to external standards, positive shifts being downfield, SiMe₄ for ¹H, ¹³C and ²⁹Si nuclei, and 85% H₃PO₄ for ³¹P NMR. The operation temperature was in the range from 293 to 300 K. The single types are assigned as follows: s = singlet, d = dublet, t = triplet, q = quartet, m = mutiplet.

IR spectra were recorded on a Bio-Rad Digilab FTS-7 spectrometer as Nujol mulls on KBr plates. Intensities were abbreviated as follows: s (strong), m (medium), w (weak). Only the characteristic absorptions are listed.

Mass spectra were obtained on a Finnigan MAT 8230 or a varian MAT CH5 instrument by EI-, FI- and FAB-MS techniques. Only the highest peak of an isotope distribution is given.

4. Experimental Section

Elemental analyses were performed at the Analytical Laboratory of the Institute of Inorganic Chemistry, University of Göttingen.

4.3 Starting materials

Commercially available chemicals were purchased from Fluka, Acros Organics or Aldrich and used as received. The other compounds used in this thesis were prepared according to published procedures: BiCl₃ (Aldrich), Li(NMe₂) (Aldrich), AlEt₃ (Aldrich), MeAlCl₂ (Aldrich), AlCl₃ (Aldrich), AlMe₃ (2.0 M solution in *n*-hexane, Acros), MeLi (1.6 M in Et₂O, Acros Organics), sulfur (Aldrich), 1,8-diaminonaphthalene (Aldrich), hexamethyldisilazane (Fluka), Cp*₂ZrMe₂ (Fluka), Cp*₂HfCl₂ (Fluka), AlH₃·NMe₃,¹⁴⁹ N,N'-bis(trimethylsilyl)diphenyliminophosphonamide¹⁵⁰, [Al(NMe₂)₃]₂,¹⁵¹ Bi(NMe₂)₃,⁶ 1,8-C₁₀H₆(NHSiMe₃)₂,^{88,152} LMeM(OH) (M= Al, Ga),^{153,154} LGe(OH),¹⁵⁵ (L=CH(NAr(CMe))₂, Ar=2,6-*i*Pr₂C₆H₃), Cp*₂MeZr(OH).¹⁵⁶

4.4. Synthesis

4.4.1. Synthesis of [1,8-C₁₀H₆(NSiMe₃)₂BiNMe₂] (1)

Compound 1,8-C₁₀H₆(NHSiMe₃)₂ (1.68g, 5.55 mmol) in *n*-hexane (30 mL) was added to Bi(NMe₂)₃ (1.68g, 5.55 mmol) in *n*-hexane (30 mL) at -30 °C. The reaction mixture was warmed to room temperature and stirred for additionally 12 h. Then the mixture was concentrated and stored at -30 °C in a freezer to obtain orange crystals. Yield: 2.39g (78%); Mp 105 °C. ¹H NMR (500.13 MHz, C₆D₆, 25 °C, TMS): δ 7.42 (d, 2H), 7.26 (t, 2H), 7.21 (d, 2H), 2.84(s, 6H), 0.25(s, 18H). ¹³C NMR (125.8 MHz, C₆D₆, TMS): δ 147.8, 130.7, 138.3, 126.5, 120.8, 116.8, 41.9, 3.0. ²⁹Si NMR (99.3 MHz, C₆D₆, 298 K): δ 3.3. Anal. Calcd for C₁₈H₃₀BiN₃Si₂ (553.18): C 39.05, H 5.46, N 7.59. Found: C 38.51, H 5.51, N 7.16.

4.4.2. Synthesis of $L^1\text{BiOCPH}(2\text{-Py})(\text{NMe}_2)$ (2)

A stirred solution of 2-benzoylpyridine (0.19g, 1.04 mmol) in *n*-hexane (20 mL) was added to a solution of $L^1\text{BiNMe}_2$ (**1**) [$L^1=1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2$] (0.57g, 1.04 mmol) in *n*-hexane (20 mL). The reaction was allowed to stir at room temperature for 1 h. Then the mixture was concentrated and stored at $-30\text{ }^\circ\text{C}$ in a freezer to obtain yellow crystals. Yield: 0.63g (83%). ^1H NMR (500.13 MHz, C_6D_6 , $25\text{ }^\circ\text{C}$, TMS): δ 8.06 (m, 1H), 7.42-7.53 (m, 5H), 7.30-7.33 (m, 2H), 7.09-7.16 (m, 3H), 6.91-6.97 (m, 3H), 6.50-6.53 (m, 1H), 1.59 (s, 6H), 0.47(s, 9H), 0.32 (s, 9H). ^{13}C NMR (125.8 MHz, C_6D_6 , TMS): δ 166.9, 149.5, 148.21, 145.5, 138.7, 137.1, 130.5, 126.9, 126.6, 124.3, 122.1, 119.8, 118.9, 116.1, 94.8, 38.68, 3.15, 3.06. ^{29}Si NMR (99.3 MHz, C_6D_6 , 298 K): δ 0.034, -0.81. Anal. Calcd for $\text{C}_{30}\text{H}_{39}\text{BiN}_4\text{OSi}_2$ (736.81): C 48.9, H 5.34, N 7.60. Found: C 48.18, H 5.39, N 7.35.

4.4.3. Synthesis of $L^1\text{BiOCH}(3\text{-Py})(\text{NMe}_2)$ (3)

A stirred solution of 3-pyridinecarboxaldehyde (0.214g, 2 mmol) in *n*-hexane (20 mL) was added to a solution of **1** (1.1g, 2 mmol) in *n*-hexane (20 mL). The reaction mixture was allowed to stir at room temperature for 1 h. Then the mixture was concentrated and stored at $-30\text{ }^\circ\text{C}$ in a freezer to obtain yellow crystals. Yield: 0.97g (75%); Mp $152\text{ }^\circ\text{C}$. ^1H NMR (500.13 MHz, C_6D_6 , $25\text{ }^\circ\text{C}$, TMS): δ 8.2-8.29 (m, 2H), 7.16-7.43 (m, 6H), 6.48-6.69 (m, 2H), 5.05(s, 1H), 1.89 (s, 6H), 0.30(s, 9H), 0.26 (s, 9H). ^{13}C NMR (125.8 MHz, C_6D_6 , TMS): δ 149.1, 148.5, 148.4, 148.1, 138.2, 137.9, 134.5, 130.5, 126.7, 122.7, 120.6, 116.7, 116.2, 89.23, 39.35, 3.06, 3.05. ^{29}Si NMR (99.3 MHz, C_6D_6 , 298 K): δ 1.92, 1.77. Anal. Calcd for $\text{C}_{24}\text{H}_{35}\text{BiN}_4\text{OSi}_2$ (660.71): C 43.63, H 5.34, N 8.48. Found: C 43.79, H 5.53, N 8.27.

4.4.4. Synthesis of $L^1\text{BiC}(\text{Me})(\text{CN})\text{CH}_2\text{NMe}_2$ (**4**)

A stirred solution of 2-methyl-2-propenenitrile (0.09g, 1.34 mmol) in *n*-hexane (20 mL) was added to a solution of **1** (0.74g, 1.34 mmol) in *n*-hexane (20 mL). The reaction was allowed to stir at room temperature for 12 h. Then the mixture was concentrated and stored at $-30\text{ }^\circ\text{C}$ in a freezer to obtain yellow crystals of **4**. Yield: 0.61g (74%); Mp $108\text{ }^\circ\text{C}$. ^1H NMR (500.13 MHz, C_6D_6 , $25\text{ }^\circ\text{C}$, TMS): δ 7.19-7.4 (m, 5H), 7.01-7.06 (m, 1H), 4.53 (d, 1H), 2.43 (d, 1H), 2.08 (s, 6H), 1.17 (s, 3H), 0.43 (s, 9H), 0.16 (s, 9H). ^{13}C NMR (125.8 MHz, C_6D_6 , TMS): δ 149.4, 147.2, 138.5, 130.5, 126.7, 125.8, 124.2, 122.5, 119.2, 116.8, 62.6, 46.7, 18.4, 3.1, 2.8. ^{29}Si NMR (99.3 MHz, C_6D_6 , 298 K): δ 5.41, 3.46. Anal. Calcd for $\text{C}_{22}\text{H}_{35}\text{BiN}_4\text{Si}_2$ (620.69): C 42.57, H 5.68, N 9.03. Found: C 41.61, H 5.71, N 9.03.

4.4.5. Synthesis of $L^1\text{BiC}(\text{CO}_2\text{Et})\text{CNMe}_2(\text{CONMe}_2)$ (**5**)

A stirred solution of diethyl acetylenedicarboxylate (0.104g, 0.614 mmol) in *n*-hexane (20 mL) was added to a solution of **1** (0.68g, 1.23 mmol) in *n*-hexane (20 mL). The reaction was allowed to stir at room temperature for 12 h. Then the mixture was concentrated and stored at $-30\text{ }^\circ\text{C}$ in a freezer to obtain yellow crystals of **5**. Yield: 0.24g (55%); Mp $138\text{ }^\circ\text{C}$. ^1H NMR (300.13 MHz, C_6D_6): δ 7.07-7.28 (m, 5H), 6.84-76.89 (m, 1H), 3.81-4.11 (m, 2H), 2.43 (s, 6H), 2.42 (s, 3H), 1.29 (s, 3H), 0.91 (t, 3H), 0.52 (s, 9H), 0.22 (s, 9H). ^{13}C NMR (75.47 MHz, C_6D_6): δ 174.4, 169.3, 159.6, 151.3, 150.5, 138.6, 133.0, 126.6, 125.9, 119.7, 118.2, 115.7, 114.7, 60.2, 40.9, 36.2, 33.4, 14.9, 3.3, 3.2. ^{29}Si NMR (59.62 MHz, C_6D_6): δ 2.154, 1.951. IR (Nujol, cm^{-1}): $\tilde{\nu}$ = 1638, 1554, 1475, 1375, 1327, 1280, 1245, 1068, 1041, 892, 862, 832, 755. Anal. Calcd for $\text{C}_{26}\text{H}_{41}\text{BiN}_4\text{O}_3\text{Si}_2$ (722.79): C 43.21, H 5.72, N 7.75. Found: C 43.41, H 5.46, N 7.46.

4.4.6. Synthesis of 1,8- $C_{10}H_6(NSiMe_3)_2Bi(NiPr)_2CNMe_2$ (6)

To a stirred solution of N,N'-diisopropylcarbodiimide (0.1g, 0.79 mmol) in *n*-hexane (20 mL) was added a solution of **1** (0.42g, 0.76 mmol) in *n*-hexane (20 mL). The reaction was allowed to stir at room temperature for 12 h. Then the mixture was concentrated and stored at $-30\text{ }^{\circ}\text{C}$ in a freezer to obtain yellow crystals of **6**. Yield: 0.4g (78%); Mp $116\text{ }^{\circ}\text{C}$. ^1H NMR (500.13 MHz, C_6D_6): δ 7.11-7.43 (m, 6H), 3.93 (sept, 2H), 1.99 (s, 6H), 0.94 (d, 12H), 0.34 (s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 166.5, 150.2, 138.2, 132.4, 126.2, 119.8, 117.3, 47.3, 39.7, 24.9, 2.7. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 0.21.

4.4.7. Synthesis of 1,8- $C_{10}H_6(NSiMe_3)_2BiNPhCONMe_2$ (7).

To a stirred solution of phenylisocyanate (0.19g, 1.6 mmol) in *n*-hexane (20 mL) was added a solution of **1** (0.88g, 1.6 mmol) in *n*-hexane (20 mL). The reaction was allowed to stir at room temperature for 12 h. Then the mixture was concentrated and stored at $-30\text{ }^{\circ}\text{C}$ in a freezer to obtain yellow crystals of **7**. Yield: 0.73g (69%); Mp $151\text{ }^{\circ}\text{C}$. ^1H NMR (500.13 MHz, C_6D_6): δ 7.31-7.36 (m, 2H), 7.16-7.22 (m, 2H), 6.97-7.03 (m, 2H), 6.84-6.91 (m, 2H), 6.59-6.66 (m, 1H), 6.18-6.23 (m, 2H), 2.08 (s, 6H), 0.31 (s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 148.2, 144.3, 138.4, 130.6, 126.5, 126, 123.8, 119.5, 115.5, 37.5, 2.71. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 2.93. Anal. Calcd for $C_{25}H_{35}BiN_4OSi_2$: C, 44.63; H, 5.24; N, 8.33. Found: C, 44.99; H, 5.51; N, 8.26.

4.4.8. Synthesis of 1,8- $C_{10}H_6(NSiMe_3)_2Bi(\mu-O)AlMeL$ (8)

n-Hexane (40 mL) was added to a mixture of LMeAl(OH) (0.33 g, 0.69 mmol) ($\text{L}=\text{CH}(\text{NAr}(\text{CMe}))_2$, $\text{Ar}=2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3$) and **1** (0.38 g, 0.69 mmol). The resulting solution was stirred for 12 h at room temperature. After concentration and keeping the solution at $-30\text{ }^{\circ}\text{C}$ in a freezer yellow crystals of **8** were isolated. Yield 0.47 g, (69 %); Mp $174\text{ }^{\circ}\text{C}$. ^1H NMR

(300.13 MHz, C₆D₆): δ 6.95- 7.24 (m, 12H), 4.68 (s, 1H), 3.06- 3.18 (m, 4H), 1.47 (s, 6H), 1.31 (d, 6H), 1.14 (d, 6H), 1.08 (d, 6H), 0.99 (d, 6H), 0.24 (s, 18H), -0.69 (s, 3H) ppm. ¹³C NMR (75.47 MHz, C₆D₆): δ 169, 146.6, 144.4, 139.9, 137.8, 129.6, 127, 126.2, 124.4, 124.3, 120.2, 116.6, 96.1, 28.6, 27.6, 26.2, 25.1, 24.9, 24.7, 23.2, 3.15, -8.4 ppm. ²⁹Si NMR (59.6 MHz, C₆D₆): δ 0.26 ppm. EI-MS [*m/z* (%): 984.5 (100) [M]⁺, 969 (36) [M-Me]⁺. Anal. Calcd for C₄₆H₆₈AlBiN₄OSi₂ (985.2): C, 56.08; H, 6.96; N, 5.69. Found: C, 55.88; H, 7.11; N, 5.78.

4.4.9. Synthesis of 1,8-C₁₀H₆(NSiMe₃)₂Bi(μ -O)GaMeL (**9**)

n-Hexane (40 mL) was added to a mixture of LMeGa(OH) (0.37 g, 0.71 mmol) and **1** (0.4 g, 0.71 mmol). The resulting solution was stirred for 12 h at room temperature. Yellow crystals of **9** were isolated from the mixture of *n*-hexane/toluene solution (3:1 v/v) at -30 °C in a freezer. Yield 0.39 g, (54 %); Mp 191 °C. ¹H NMR (300.13 MHz, C₆D₆): δ 6.96- 7.23 (m, 12H), 4.57 (s, 1H), 3.15- 3.23 (m, 4H), 1.57 (s, 6H), 1.28 (d, 6H), 1.14 (d, 6H), 1.09 (d, 6H), 1.02 (d, 6H), 0.28 (s, 18H), -0.56 (s, 3H) ppm. ¹³C NMR (125.7 MHz, C₆D₆): δ 167.7, 146.7, 144.7, 140.4, 137.8, 129.7, 127, 126.2, 124.4, 124.3, 120.2, 116.6, 96, 28.8, 27.8, 26.3, 25.2, 24.7, 24.6, 23.3, 3.17, -6.5 ppm. ²⁹Si NMR (99.3 MHz, C₆D₆): δ 0.52 ppm. Anal. Calcd for C₄₆H₆₈BiGaN₄OSi₂ (1027.9): C, 53.75; H, 6.67; N, 5.45. Found: C, 53.26; H, 6.71; N, 5.61.

4.4.10. Synthesis of 1,8-C₁₀H₆(NSiMe₃)₂Bi(μ -O)GeL (**10**)

n-Hexane (30 mL) was added to a mixture of LGe(OH) (0.32 g, 0.63 mmol) and **1** (0.35 g, 0.63 mmol). The resulting solution was stirred for 12 h at room temperature. After concentration and keeping the solution at -30 °C in a freezer yellow crystals of **10** were isolated. Yield 0.41 g, (63 %); Mp 251 °C. ¹H NMR (300.13 MHz, C₆D₆): δ 6.91-7.19 (m, 12H), 4.56 (s, 1H), 3.37 (sept, 2H), 3.14 (sept, 2H), 1.47 (s, 6H), 1.28 (d, 6H), 1.20 (d, 6H),

1.04 (d, 6H), 1.07 (d, 6H), 0.24 (s, 18H) ppm. ^{13}C NMR (75.47 MHz, C_6D_6): δ 162.6, 146.7, 145.2, 144, 139, 137.7, 130.1, 126.8, 126.4, 124.4, 120.5, 116.7, 95, 31.9, 28.7, 28.3, 26.8, 25.1, 24.8, 22.9, 22.6, 14.2, 3.0 ppm. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 0.99 ppm. EI-MS [m/z (%)]: 1016.4 (100) $[\text{M}]^+$, 599 (62) $[\text{M}-\text{CH}(\text{DippNCMe})_2]^+$. Anal. Calcd for $\text{C}_{48}\text{H}_{72}\text{BiGeN}_4\text{OSi}_2$ (1058.85, $10 \cdot \text{C}_7\text{H}_8$): C, 54.39; H, 6.79; N, 5.28. Found: C, 53.64; H, 6.79; N, 5.28.

4.4.11. Synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{ZrMeCp}^*_2$ (**11**)

n-Hexane (40 mL) was added to $\text{Cp}^*_2\text{MeZr}(\text{OH})$ (0.43 g, 1.1 mmol) and **1** (0.6 g, 1.1 mmol). The resulting solution was stirred for 12 h at room temperature, and then the solvent was partly removed and the solution kept at -30 °C in a freezer. Yellow crystals of **11** were isolated after 2 days. Yield 0.72 g, (73 %); Mp 206 °C. ^1H NMR (300.13 MHz, C_6D_6): δ 7.44-7.47 (m, 2H), 7.17-7.31 (m, 4H), 1.63 (s, 30H), 0.28 (s, 18H), -0.43 (s, 3H) ppm. ^{13}C NMR (75.47 MHz, C_6D_6): δ 147.2, 138.5, 130.4, 126.8, 120.4, 118.7, 117.2, 25.7, 11.3, 3.07 ppm. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 1.1 ppm. EI-MS [m/z (%)]: 524.2 (78) $[\text{M}-\text{Cp}^*_2\text{Zr}(\text{Me})\text{-H}]^+$, 494.1 (100) $[\text{M}-\text{Cp}^*_2\text{Zr}(\text{Me})\text{-OMe}]^+$. Anal. Calcd for $\text{C}_{37}\text{H}_{57}\text{BiN}_2\text{OSi}_2\text{Zr}$ (902.23): C, 49.25; H, 6.37; N, 3.10. Found: C, 48.78; H, 6.39; N, 3.19.

4.4.12. Synthesis of $1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}(\mu\text{-O})\text{HfClCp}^*_2$ (**12**)

To a solution of $\text{Cp}^*_2\text{HfCl}_2$ (0.445 g, 0.846 mmol) and $[\text{CN}(i\text{Pr})\text{C}_2\text{Me}_2\text{N}(i\text{Pr})]$ (:C, 0.15 g, 0.846 mmol) in toluene (60 mL) degassed and distilled water (15 μL , 0.846 mmol) was slowly added under vigorous stirring over a period of 30 min. The mixture was stirred for another 30 min and the formation of a white precipitate was observed. The resulting solid was filtered off and the filtrate was added to a flask containing **1** (0.47 g, 0.846 mmol) at -30 °C. The mixture was stirred for 1 h at this temperature and then was allowed to attain room temperature and stirring was continued for 12 h. All volatiles were removed *in vacuo* and the

4. Experimental Section

resulting solid was washed with *n*-hexane (20 mL) to give yellow crystalline **12**. The product was recrystallized from a toluene/*n*-hexane (2:1 v/v) mixture to yield yellow crystals at -30 °C. Yield 0.29 g (34 %); Mp 210 °C. ^1H NMR (300.13 MHz, C_6D_6): δ 7.44-7.47 (m, 2H), 7.22-7.34 (m, 4H), 1.73 (s, 30H), 0.3 (s, 18H) ppm. ^{13}C NMR (75.47 MHz, C_6D_6): δ 147.8, 138.5, 126.8, 120.3, 119.6, 116.8, 11.5, 2.9 ppm. ^{29}Si NMR (59.62 MHz, C_6D_6): δ 1.12 ppm. EI-MS [m/z (%)]: 1010.5 (43) $[\text{M}]^+$, 875 (100) $[\text{M-Cp}^*]^+$. Anal. Calcd for $\text{C}_{36}\text{H}_{54}\text{BiClHfN}_2\text{OSi}_2$ (1009.91): C, 42.77; H, 5.34; N, 2.77. Found: C, 42.40; H, 5.35; N, 2.80.

4.4.13. Synthesis of 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{BiCl}$ (**13**)

To a stirred solution of **1** (0.7g, 1.26 mmol) in *n*-hexane (30 mL) was added 0.16 mL (1.26 mmol) of ClSiMe_3 at -10 °C. The reaction mixture was warmed to room temperature and stirred over night. All volatiles were removed *in vacuo* and the resulting solid was washed with cold *n*-hexane (20 mL) to give red crystalline solid. The product was recrystallized from a toluene/*n*-hexane mixture (2:1) to yield red crystals of **13** at -30 °C. Yield: 0.47g (69%); Mp 163 °C. ^1H NMR (500.13 MHz, C_6D_6): δ 7.41 (d 2H), 7.28 (t, 2H), 7.16 (d 2H), 0.17(s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 146.53, 138, 131.7, 126.8, 121.1, 116.3, 2.65. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 4.6. Anal. Calcd for $\text{C}_{16}\text{H}_{24}\text{BiClN}_2\text{Si}_2$: C, 35.26; H, 4.44; N, 5.14. Found: C, 34.92; H, 4.21; N 5.09.

4.4.14. Synthesis of 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{BiC}\equiv\text{CPh}$ (**14**)

To a stirred solution of **1** (0.7g, 1.26 mmol) in *n*-hexane (30 mL) was added 0.14 mL (1.26 mmol) of phenyl acetylene at -10 °C. The reaction mixture was warmed to room temperature and stirred over night. Then the mixture was concentrated and stored at -30 °C in a freezer. Orange crystals of **14** were obtained after two days. Yield: 0.41g (54%); Mp 98

°C. ^1H NMR (500.13 MHz, C_6D_6): δ 7.48-7.51 (m, 2H), 7.24-7.31 (m, 3H), 6.85-6.89 (m, 2H), 6.74-6.77 (m, 3H), 0.23 (s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 150.1, 138, 134, 131.9, 126.4, 123.7, 120.6, 116.2, 111.9, 3.1. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 4.1. Anal. Calcd for $\text{C}_{24}\text{H}_{29}\text{BiN}_2\text{Si}_2$: C, 47.2; H, 4.79; N, 4.59. Found: C, 46.66; H, 4.80; N, 4.42.

4.4.15. Synthesis of 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{BiCp}^*(15)$

The *n*-hexane solution (30 mL) of **1** (0.28 g, 0.5 mmol) was added to a *n*-hexane solution (20 mL) of Cp^*H (0.07 g, 0.5 mmol) at room temperature. The reaction mixture was stirred over night, then concentrated and stored at -30 °C in a freezer to obtain orange crystals of **15**. Yield: 0.22g (68%); Mp 148 °C. ^1H NMR (500.13 MHz, C_6D_6): δ 7.23-7.41 (m, 6H), 1.86 (s, 315H), 0.27 (s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 148.8, 138.9, 130.3, 127.8, 126.6, 124.5, 120.3, 117.7, 10.4, 3.3. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 1.14. Anal. Calcd for $\text{C}_{26}\text{H}_{39}\text{BiN}_2\text{Si}_2$: C, 48.43; H, 6.10; N, 4.34. Found: C, 47.94; H, 5.98; N, 4.45.

4.4.16. Synthesis of 1,8- $\text{C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{BiMe}$ (16)

AlMe_3 (0.8 mL, 1.6 mmol, 2.0 M in *n*-hexane) was added to a *n*-hexane (50 mL) solution of **1** (0.88 g, 1.6 mmol) at -78 °C. The mixture was stirred for 1 h at this temperature and then was allowed to attain room temperature and stirring was continued for 12 h. The mixture was concentrated and stored at -30 °C in a freezer to give yellow microcrystalline solid of **17**. Yield: 0.31 g (37 %); Mp 109 °C. ^1H NMR (500.13 MHz, C_6D_6): δ 7.23-7.31 (m, 4H), 7.44-7.48 (m, 2H), 0.43 (s, 3H), 0.28 (s, 18H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 149.2, 138.4, 131.2, 126.2, 120.7, 117.9, 3.3, - 9.6. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 3.7. Anal. Calcd for $\text{C}_{17}\text{H}_{27}\text{BiN}_2\text{Si}_2$: C, 38.92; H, 5.19; N, 5.34. Found: C, 38.75; H, 5.11; N, 5.46.

4.4.17. Synthesis of 1,8-C₁₀H₆(NSiMe₃)₂BiOPh (17)

n-Hexane (40 mL) was added to the mixture of **1** (0.7g, 1.26 mmol) and phenol (0.12g, 1.26 mmol) at room temperature and stirred over night. After filtration the resulting solution concentrated and stored at -30 °C in a freezer to obtain a yellow crystals of **17**. Yield: 0.48g (64%); Mp 149 °C. ¹H NMR (500.13 MHz, C₆D₆): δ 7.41 (d, 2H), 7.24 (t, 2H), 6.94-7.02 (m, 4H), 6.65 (t, 1H), 6.29 (d, 2H), 0.11 (s, 18H). ¹³C NMR (125.8 MHz, C₆D₆): δ 159.2, 146.5, 138.2, 130.5, 126.7, 122.1, 120.8, 120.5, 117.1, 2.4. ²⁹Si NMR (59.6 MHz, C₆D₆): δ 3.11. Anal. Calcd for C₂₂H₂₉BiN₂OSi₂: C, 43.85; H, 4.85; N, 4.65. Found: C, 43.65; H, 4.91; N, 4.65.

4.4.18. Synthesis of 1,8-C₁₀H₆(NSiMe₃)₂BiNH₂ (18)

Dry ammonia gas was bubbled through the solution of **1** (0.6g, 1.1 mmol) in *n*-hexane (40 mL) for 15 min at room temperature. After that the solvent was removed in vacuum and the residue was extracted with cold *n*-pentane (30 mL). The solvent was removed under reduced pressure to yield 0.41g (71%) of compound **18** as a colorless oil. ¹H NMR (500.13 MHz, C₆D₆): δ 7.28-7.33 (m, 2H), 7.17-7.20 (m, 2H), 6.68-6.72 (m, 2H), 5.38 (s, 2H), 0.16 (s, 18H). ¹³C NMR (125.8 MHz, C₆D₆): δ 144.5, 138.1, 125.9, 121.1, 116.1, -0.05. ²⁹Si NMR (59.6 MHz, C₆D₆): δ 3.38.

4.4.19. Synthesis of [1,8-C₁₀H₆(NSiMe₃)₂Bi]₂S (19)

n-Hexane (50 mL) was added to the mixture of **1** (0.41g, 0.75 mmol) and sulfur (0.024g, 0.75 mmol) at room temperature and stirred for 2 days. After filtration, the resulting solution concentrated and stored at -30 °C in a freezer to obtain yellow crystals. Yield: 0.48g (61%); Mp 130 °C. ¹H NMR (500.13 MHz, C₆D₆): δ 7.11-7.44 (m, 10H), 6.68-6.72 (m, 2H), 0.18 (s,

36H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 149.3, 144.5, 138.2, 132.8, 125.9, 121.2, 120, 116.6, 116.1, 3.06. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 2.86.

4.4.20. Synthesis of $[1,8\text{-C}_{10}\text{H}_6(\text{NSiMe}_3)_2\text{Bi}]_2$ (**20**)

The *n*-hexane solution (30 mL) of **1** (1.0 g, 1.8 mmol) was added to a *n*-hexane solution (20 mL) of PhSiH_3 (0.2 g, 1.85 mmol) at $-30\text{ }^\circ\text{C}$. The reaction mixture was warmed to room temperature and stirred for additionally 12 h. Then the mixture was concentrated and stored at $-30\text{ }^\circ\text{C}$ in a freezer to obtain red crystals of **20**. Yield (1.1g 60%); Mp $245\text{ }^\circ\text{C}$; ^1H NMR (500.13 MHz, C_6D_6): δ 7.13-7.22 (m, 12H), 0.15 (s, 36H). ^{13}C NMR (125.8 MHz, C_6D_6): δ 155.7, 138.4, 137.1, 124.9, 120.4, 115.3, 3.7. ^{29}Si NMR (59.6 MHz, C_6D_6): δ 6.69. Anal. Calcd for $\text{C}_{32}\text{H}_{48}\text{Bi}_2\text{N}_4\text{Si}_4$: C, 37.72; H, 4.75; N, 5.50. Found: C, 37.66; H, 4.48; N, 5.58.

4.4.21. Synthesis of $\text{LiPh}_2\text{P}(\text{NSiMe}_3)_2$ (**22**)

To a solution of **21** (3.0 g, 8.33 mmol) in toluene (80 mL) at $-78\text{ }^\circ\text{C}$ was added *n*BuLi (2.5 M, 3.4 mL, 8.5 mmol). The mixture was warmed to room temperature and stirred overnight. Toluene was removed under reduced pressure and the product was recrystallized from a mixture of toluene and *n*-hexane. $[\text{LiPh}_2\text{P}(\text{NSiMe}_3)_2]$ formed colorless crystals. Yield (0.84 g, 83%), whose spectral data agreed with those reported in literature. ^1H NMR (200 MHz, C_6D_6 , 298 K): δ 0.12 (s, 18H, Si-Me), 7.61-7.80 (m, 4H, *o*-Ar), 6.98-7.07 (m, 6H, *p*-/*m*-Ar). ^{31}P NMR (121.50 MHz, C_6D_6 , 298 K): δ 7.4.

4.4.22. Synthesis of $\text{Ph}_2\text{P}(\text{NSiMe}_3)_2\text{AlCl}_2$ (**23**)

A toluene solution of freshly prepared **22** (1.0 g, 2.73 mmol) was added drop by drop to a cold suspension of freshly sublimed AlCl_3 (0.36 g, 2.73 mmol) in toluene at $-78\text{ }^\circ\text{C}$. The temperature was raised to $0\text{ }^\circ\text{C}$ and stirred for 1 h and stirring was continued for 3 h. The solution was allowed to attain room temperature and stirring was continued for 12 h. Then the

mixture was filtered, and the concentrated solution was stored in a freezer at -30 °C to obtain colorless crystals. Yield (0.76g 61%); Mp 135 °C (dec); ¹H NMR (200 MHz, C₆D₆, 298 K): δ -0.2 (s, 18H, SiCH₃), 7.65-7.81 (m, 4H, o-Ar), 6.95-7.07 (m, 6H, p-/m-Ar). ¹³C NMR (125.76 MHz, C₆D₆, 298 K): δ 1.91, 1.93, 132.3-127.4. ³¹P NMR (121.50 MHz, C₆D₆): δ 37.8.

4.4.23. Synthesis of Ph₂P(NSiMe₃)₂AlClMe (24)

AlMeCl₂ (1.0 M in *n*-hexane, 1.4 mL, 1.4 mmol) was added drop by drop to the solution of freshly prepared **22** (0.5 g, 1.36 mmol) in toluene (70 mL) at -78 °C. The mixture was stirred at -78 °C for 1 h and then the temperature was slowly raised to 0 °C and stirring was continued for 3 h. After that the solution was allowed to attain room temperature and stirring was continued for 12 h. Then the mixture was extracted with *n*-hexane (70 mL), and the concentrated solution was stored in a freezer at -30 °C to obtain colorless crystals. Yield (0.41g 68%); Mp 135 °C; ¹H NMR (200 MHz, C₆D₆, 298 K): δ -0.03 (s, 18H, Si-Me), 0.04 (s, 3H, Al-Me), 7.65-7.85 (m, 4H, o-Ar), 6.92-7.07 (m, 6H, p-/m-Ar). ¹³C NMR (125.76 MHz, C₆D₆, 298 K): δ -0.69, 1.71, 1.66, 131.6-129.2. ³¹P NMR (121.50 MHz, C₆D₆, 298 K): δ 33.2. ²⁹Si NMR (59.6 MHz, C₆D₆, 298 K): δ -0.31, -0.34. Anal. Calcd for C₁₉H₃₁AlClN₂PSi₂: C, 52.22; H, 7.15; N, 6.41. Found C, 52.54; H, 7.15; N, 6.21.

4.4.24. Synthesis of [Ph₂P(NSiMe₃)₂]₂AlH (25)

The *n*-hexane solution (30 mL) of **21** (1.0 g, 2.7 mmol) was added drop by drop to a cold suspension of freshly sublimed AlH₃·NMe₃ in THF (30 mL), (0.12 g, 1.39 mmol) at -78 °C. The mixture was stirred at -30 °C for 1 h and then the temperature was slowly raised to 0 °C and the stirring was continued for 3 h. Then the solution was allowed to attain room temperature under stirring for 12 h. Then the mixture was concentrated and stored in a freezer at -30 °C to obtain colorless crystals. Yield: (0.64g 61%); Mp 107 °C (dec); ¹H NMR (200

4. Experimental Section

MHz, CDCl₃, 298 K): δ -0.22 (s, 18H, Si-Me), -0.13 (s, 18H, Si-Me), 5.1 (very br, 1H, Al-H), 7.78-7.92 (m, 4H, *o*-Ar), 7.45-7.59 (m, 6H, *p*-/*m*-Ar). ¹³C NMR (125.76 MHz, C₆D₆, 298 K): δ 1.52, 1.56, 3.75, 3.80, 133.1-131.2. ³¹P NMR (121.50 MHz, C₆D₆, 298 K): δ 29.2, 34.6. ²⁹Si NMR (59.6 MHz, C₆D₆, 298 K): δ -4.15, -4.22. IR (cm⁻¹): ν (AlH) 1853. Anal. Calcd for C₃₇H₆₀AlN₄P₂Si₄: C, 58.31; H, 7.93; N, 7.35. Found C, 57.94; H, 7.83; N, 7.21.

4.4.25. Synthesis of Ph₂P(NSiMe₃)₂AlEt₂ (26)

AlEt₃ (2.2 mL, 2.2 mmol) was added to a *n*-hexane (50 mL) solution of **21** (0.8 g, 2.2 mmol) at -78 °C. The mixture was stirred for 1 h at this temperature and then was allowed to attain room temperature and stirring was continued for 12 h. The mixture was concentrated and stored in a freezer at -30 °C to give white microcrystalline solid. Yield: (0.86 g, 88 %); ¹H NMR (500 MHz, C₆D₆, 298 K): δ -0.05 (s, 18H, Si-Me), 0.45 (q, 4H, CH₂), 1.52 (t, 6H, CH₃), 7.71-7.81 (m, 4H, *o*-Ar), 6.97-7.11 (m, 6H, *p*-/*m*-Ar). ¹³C NMR (125.8 MHz, C₆D₆, 298 K): δ 1.99, 2.04, 10.4, 134.5-131.9. ³¹P NMR (121.50 MHz, C₆D₆, 298 K): δ 30.3. ²⁹Si NMR (59.6 MHz, C₆D₆, 298 K): δ -2.61, -2.54.

4.4.26. Synthesis of Ph₂P(NSiMe₃)₂Al(NMe₂)₂ (27)

n-Hexane (30 mL) solution of **21** (0.9 g, 2.5 mmol) was added to [Al(NMe₂)₃]₂ (0.4 g, 1.25 mmol) in *n*-hexane (20 mL), and then heated under reflux for 6 h. The mixture was cooled to room temperature, concentrated and stored in a freezer at -30 °C to obtain a colorless microcrystalline solid. Yield: (0.69 g, 58 %); Mp 156 °C; ¹H NMR (500 MHz, C₆D₆, 298 K): δ 0.024 (s, 18H, Si-Me), 3.06 (s, 12H, CH₃), 7.73-7.83 (m, 4H, *o*-Ar), 7.03-7.08 (m, 6H, *p*-/*m*-Ar). ¹³C NMR (125.8 MHz, C₆D₆, 298 K): δ 1.86, 1.92, 41.8, 134.3-132.0. ³¹P NMR (121.50 MHz, C₆D₆, 298 K): δ 28.6. ²⁹Si NMR (59.6 MHz, C₆D₆, 298 K): δ -2.67, -2.72.

4. Experimental Section

Anal. Calcd for $C_{22}H_{40}AlN_4PSi_2$: C, 55.66; H, 8.49; N, 11.80. Found C, 55.11; H, 8.63; N, 11.94.

5. Handling and Disposal of Solvents and Residual Waste

- 1 The recovered solvents were distilled or condensed into a cold-trap under vacuum and collected in halogen-free or halogen-containing solvent containers, and stored for disposal.
- 2 Used NMR solvents were classified into halogen-free and halogen-containing solvents and were disposed as halogen containing wastes, respectively.
- 3 Drying agents such as KOH, CaCl₂ and P₄O₁₀ were hydrolyzed and disposed as acid or base wastes.
- 4 Whenever possible, sodium metal used for drying solvents was collected for recycling.¹⁵⁷ The non-reusable sodium metal was carefully hydrolyzed in cold ethanol and poured into the base-bath used for cleaning glassware.
- 5 Ethanol and acetone used for cold-baths (with solid CO₂ or liquid N₂) were subsequently used for cleaning glassware.
- 6 The acid-bath used for cleaning glassware was neutralized with Na₂CO₃ and the resulting NaCl solution was washed-off in the communal water drainage.
- 7 The residue of the base-bath used for glassware cleaning was poured into container for base wastes.

Amounts of various types of disposable wastes generated during the work:

Heavy elements containing wastes 3 L

Halogen-containing solvent wastes 8 L

Halogen-free solvent wastes 40 L

Acid wastes 20 L

Base wastes 30 L

6. References

- (1) R. K. Grasselli, J. D. Burrington, *Adv. Catal.* **1981**, *30*, 133-163.
- (2) a) S. Shimada, O. Yamazaki, T. Tanaka, M. L. N. Rao, Y. Suzuki, M. Tanaka, *Angew. Chem.* **2003**, *115*, 1889-1892. *Angew. Chem., Int. Edit.* **2003**, *42*, 1845-1848.
b) G. O. Doak, L. D. Freedman, *Organometallic Compounds of Arsenic, Antimony, and Bismuth*; Wiley-Interscience: New York, **1970**. c) H. Gaspard-Illoughmane, C. Le Roux, *Eur. J. Org. Chem.* **2004**, 2517- 2532.
- (3) R. Dagani, *Chem. Eng. News* **1987**, *65*, 41-49.
- (4) M. Vehkamäki, T. Hatanpää, M. Ritala, M. Leskelä, *J. Mater. Chem.* **2004**, *14*, 3191-3197.
- (5) M. Vehkamäki, T. Hatanpää, M. Kemell, M. Ritala, M. Leskelä, *Chem. Mater.* **2006**, *18*, 3883-3888.
- (6) W. Clegg, N. A. Compton, R. J. Errington, G. A. Fisher, M. E. Green, D. C. R. Hockless, N. C. Norman, *Inorg. Chem.* **1991**, *30*, 4680-4682.
- (7) F. Ando, T. Hayashi, K. Ohashi, J. Koketsu, *J. Inorg. Nucl. Chem.* **1975**, *37*, 2011-2013.
- (8) S. C. James, N. C. Norman, A. G. Orpen, M. J. Quayle, U. Weckenmann, *J. Chem. Soc., Dalton Trans.* **1996**, 4159-4161.
- (9) U. Warringa, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, *Inorg. Chem.* **1994**, *33*, 4607-4608.
- (10) N. Burford, C. L. B. Macdonald, K. N. Robertson, T. S. Cameron, *Inorg. Chem.* **1996**, *35*, 4013-4016.
- (11) O. J. Scherer, P. Hornig, M. Schmidt, *J. Organomet. Chem.* **1966**, *6*, 259-264.
- (12) M. J. S. Gynane, A. Hudson, M. F. Lappert, P. P. Power, H. Goldwhite, *J. Chem. Soc. Dalton Trans.* **1980**, 2428-2433.

6. References

- (13) M. Veith, B. Bertsch, *Z. Anorg. Allg. Chem.* **1988**, 557, 7-22.
- (14) a) C. M. Kozak, P. Mountford, *Angew. Chem.* **2004**, 116, 1206-1209; *Angew. Chem., Int. Ed.* **2004**, 43, 1186-1189. b) R. Pryadun, D. Sukumaran, R. Bogadi, J. D. Atwood, *J. Am. Chem. Soc.* **2004**, 126, 12414-12420. c) Y. Ohki, M. D. Fryzuk, *Angew. Chem.* **2007**, 119, 3242-3245; *Angew. Chem., Int. Ed.* **2007**, 46, 3180-3183.
- (15) T. A. George, M. F. Lappert, *J. Organomet. Chem.* **1968**, 14, 327-337.
- (16) R. H. Cragg, M. F. Lappert, *J. Chem. Soc. A* **1966**, 82-85
- (17) P. Zhao, C. Krug, J. F. Hartwig, *J. Am. Chem. Soc.* **2005**, 127, 12066-12073.
- (18) E. Katayev, Y. Li, A. L. Odom, *Chem. Commun.* **2002**, 838-839.
- (19) C. G. Young, *Coord. Chem. Rev.* **1989**, 96, 89-251.
- (20) G. D. Lawrence, D. T. Sawyer, *Coord. Chem. Rev.* **1978**, 27, 173-193.
- (21) W. M. Coleman, L. T. Taylor, *Coord. Chem. Rev.* **1980**, 32, 1-31.
- (22) D. W. Stephan, *Coord. Chem. Rev.* **1989**, 95, 41-107.
- (23) U. Casellato, P. A. Vigato, *Coord. Chem. Rev.* **1977**, 23, 31-117.
- (24) K. Mochizuki, M. Tsutsumi, Y. Yamaji, *Inorg. Chim. Acta.* **1992**, 191, 35-37.
- (25) I. Murase, K. Hamada, S. Kida, *Inorg. Chim. Acta.* **1981**, 54, L171-L173.
- (26) J. P. Collman, J. E. Hutchison, M. A. Lopez, A. Tabard, R. Guilard, W. K. Seok, J. A. Ibers, M. L'Her, *J. Am. Chem. Soc.* **1992**, 114, 9869-9877.
- (27) K. D. Karlin, Y. Gultneh, J. P. Hutchinson, J. Zubieta, *J. Am. Chem. Soc.* **1982**, 104, 5240-5242.
- (28) J. -P. Costes, J. -F. Serra, F. Dahan, J. P. Laurent, *Inorg. Chem.* **1986**, 25, 2790-2795 and references cited therein.
- (29) N. Kitajima, T. Koda, S. Hasimoto, T. Kitagawa, Y. Moro-oka, *J. Chem. Soc., Chem. Commun.* **1988**, 151-152.

- (30) R. L. Lintvedt, M. D. Glick, B. K. Tomlonovic, D. P. Gavel, *Inorg. Chem.* **1976**, *15*, 1646-1654.
- (31) N. Kitajima, N. Tamura, M. Tanaka, Y. Moro-oka, Y. *Inorg. Chem.* **1992**, *31*, 3342-3343.
- (32) E. J. Larson, V. L. Pecoraro, *J. Am. Chem. Soc.* **1991**, *113*, 3810-3818.
- (33) R. Das, K. Nag, *Inorg. Chem.* **1991**, *30*, 2833-2835.
- (34) W. B. Tolman, S. Liu, J. G. Bentsen, S. J. Lippard, *J. Am. Chem. Soc.* **1991**, *113*, 152-164.
- (35) J. B. Vincent, J. C. Huffman, G. Christou, Q. Li, M. A. Nanny, D. N. Hendrickson, R. H. Fong, R. H. Fish, *J. Am. Chem. Soc.* **1988**, *110*, 6898-6900.
- (36) W. B. Tolman, A. Bino, S. J. Lippard, *J. Am. Chem. Soc.* **1989**, *111*, 8522-8523.
- (37) L. G. Hubert-Pfalzgraf, *Inorg. Chem. Comm.* **2003**, *6*, 102-120.
- (38) J. Gopalakrishnan, *Chem. Mater.* **1995**, *7*, 1265-1275.
- (39) T. D. Tilly, *J. Mol. Catal. A: Chemical* **2002**, *182-183*, 17-24.
- (40) P. Hodge, S. C. James, N. C. Norman, A. G. Orpen, *J. Chem. Soc., Dalton Trans.* **1998**, 4049-4054 and references therein.
- (41) S. Parola, R. Papiernik, L. G. Hubert-Pfalzgraf, C. Bois, *J. Chem. Soc., Dalton Trans.* **1998**, 737-739.
- (42) a) T. Ono, N. Ogata, R. L. Kuczkowski, *J. Catal.* **1998**, *175*, 185-193. b) R. K. Grasselli, J. D. Burrington, *Ind. Eng. Chem. Prod. Res. Dev.* **1984**, *23*, 393-404.
- (43) a) N. Zhang, Z. Ding, Y. Wu, M. Salomon, *IEEE Trans. Nucl. Science*, **1990**, *37*, 216-219. b) C. Li, T. Yoshino, *Appl. Opt.* **2002**, *41*, 5391-5397.
- (44) C. K. Lee, B. H. Bay, A. R. West, *J. Mater. Chem.* **1996**, *6*, 331-335.
- (45) H. J. Breunig, L. Königsmann, E. Lork, M. Nema, N. Philipp, C. Silvestru, A. Soran, R. A. Varga, R. Wagner, *Dalton Trans.* **2008**, 1831-1842.
- (46) G. Balazs, H. J. Breunig, E. Lork, *Organometallics* **2002**, *21*, 2584-2586.

- (47) A. J. Ashe, III; E. G. Ludwig, Jr, J. Oleksyszyn, *Organometallics* **1983**, *2*, 1859- 1866.
- (48) F. Calderazzo, A. Morvillo, G. Pelizzi, R. Poli, F. Ungari, *Inorg. Chem.* **1988**, *27*, 3730-3733.
- (49) S. Hoppe, K. H. Whitmire, *Organometallics* **1998**, *17*, 1347-1354.
- (50) L. Balazs, H. J. Breunig, E. Lork, C. Silvestru, *Eur. J. Inorg. Chem.* **2003**, 1361- 1365.
- (51) L. Balazs, H. J. Breunig, E. Lork, A. Soran, C. Silvestru, *Inorg. Chem.* **2006**, *45*, 2341-2346.
- (52) M. Wieber, I. Z. Sauer, *Z. Naturforsch.* **1987**, *42B*, 695-698.
- (53) a) L. Balazs, H. J. Breunig, *Coord. Chem. Rev.* **2004**, *248*, 603-621. b) H. J. Breunig, *Z. Anorg. Allg. Chem.* **2005**, *631*, 621-631.
- (54) H. Suzuki, Y. Matano, *Organobismuth Chemistry*, Elsevier, **2001**.
- (55) A. J. Ashe, III, J. W. Kampf, D. B. Puranik, S. M. Al-Taweel, *Organometallics* **1992**, *11*, 2743-2745.
- (56) H. J. Breunig, R. Rösler, E. Lork, *Angew. Chem.* **1998**, *110*, 3361-3363; *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 3175-3177.
- (57) D. C. Bradley, *Chem. Rev.* **1989**, *89*, 1317-1322.
- (58) See for example: G. A. Olah, S. Kobayashi, M. Tashiro, *J. Am. Chem. Soc.* **1972**, *94*, 7448-7461.
- (59) H. C. Brown, C. J. Shoaf, *J. Am. Chem. Soc.* **1964**, *86*, 1079-1085.
- (60) E. Wiberg, E. Amberger, *Hydrides of the Elements of Main Group I-IV*, Elsevier, New York, **1971**, Ch. 5.
- (61) A. Murphy, *Nature* **1991**, *350*, 223-225.
- (62) a) A. R. Barron, *Adv. Mater. Opti. Electron.* **1995**, *5*, 245-258. b) J. L. Atwood in *Coordination Chemistry of Aluminium* (Ed.: G. H. Robinson), VCH, New Yourk, **1993**, pp. 197-232. c) H. Sinn, W. Kaminsky, H. J. Vollmer, R. Woldt, *Angew.*

- Chem.* **1980**, *92*, 396-402; *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 390-392. d) H. Sinn, W. Kaminsky, *Adv. Organomet. Chem.* **1980**, *18*, 99-149.
- (63) R. G. Gordon, D. M. Hoffman, U. Riaz, *J. Mater. Res.* **1991**, *6*, 5-7.
- (64) R. G. Gordon, U. Riaz, D. M. Hoffman, *J. Mater. Res.* **1992**, *7*, 1679-1684.
- (65) Y. Takahashi, K. Yamashita, S. Motojima, K. Sugiyama, *Surf. Sci.* **1979**, *86*, 238-245.
- (66) K. L. Ho, K. F. Jensen, J. W. Hwang, W. L. Gladfelter, J. F. Evans, *J. Cryst. Growth* **1991**, *107*, 376-380.
- (67) K. L. Ho, D. C. Boyd, K. F. Jensen, S. A. Hanson, W. L. Gladfelter, J. F. Evans, *Mater. Res. Symp. Proc.* **1990**, *132*, 162.
- (68) R. K. Schulze, D. R. Mantell, W. L. Gladfelter, J. F. Evans, *J. Vac. Sci. Technol. A* **1988**, *6*, 2162-2163.
- (69) L. V. Interrante, W. Lee, M. McConnell, N. Lewis, E. Hall, *J. Electrochem. Soc.* **1989**, *136*, 472-478.
- (70) F. C. Sauls, L. V. Interrante, *Coord. Chem. Rev.* **1993**, *128*, 193-207.
- (71) K. Ziegler, F. Krupp, K. Zosel, *Angew. Chem.* **1955**, *67*, 425.
- (72) D. A. Neumayer, J. G. Ekerdt, *Chem. Mater.* **1996**, *8*, 9-25, and references cited therein.
- (73) D. C. Boyd, R. T. Haasch, P. R. Mantell, R. K. Schulze, J. F. Evans, W. L. Gladfelter, *Chem. Mater.* **1989**, *1*, 119-124.
- (74) A. Rabenau in: *Compounds Semiconductors* (Eds.: R. K. Willardson, H. L. Goerring), Reinhold Publishing Corp., New York, USA, **1962**, Vol. 1, Chapter 19, 174.
- (75) a) L.V. Interrante, L.E. Carpenter, C. Whitmarsch, W. Lee, G. A. Slack, *Mater. Res. Soc. Symp. Proc* **1986**, *73*, 359-366. b) *Chemistry of Aluminum, Gallium, Indium and Thallium* (Ed.: A. J. Downs), Blackie, Glasgow, **1993**. c) A. C. Jones, P. O'Brien in *CVD of Compound Semiconductors*, VCH, Weinheim, **1997**

- (76) a) G. H. Robinson, Ed., *Coordination Chemistry of Aluminum*, VCH Publishers; Weinheim, FRG, **1994**, pp.1-56; (b) G. H. Robinson, Ed., *Coordination Chemistry of Aluminum*, VCH Publishers; Weinheim, FRG, **1994**, pp.57-84.
- (77) a) L.V. Interrante, G.A. Sigel, M. Garbaskas, C. Hejna, G.A. Slack, *Inorg. Chem.* **1989**, 28, 252-257. b) W. Rockensüss, H. W. Roesky, *Adv. Mater.* **1993**, 5, 443-445.
- (78) D. C. Bradley, I. S. Harding, I. A. Maria, M. Motevalli, *J. Chem. Soc., Dalton Trans.* **1997**, 2969-2979.
- (79) K. J. Alford, K. Gosling, J. D. Smith, *J. Chem. Soc., Dalton Trans.* **1972**, 20, 2203-2208.
- (80) N. Wiberg, W. Baumeister, P. Zhan, *J. Organomet. Chem.* **1972**, 36, 267-276.
- (81) a) T. Akakibara, T. Hirabayashi, Y. Ishi, *J. Organomet. Chem.* **1972**, 46, 231-242.
- (82) C. Cui, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, *Organometallics* **1999**, 18, 5120-5123.
- (83) Z. Yang, X. Ma, R. B. Oswald, H. W. Roesky, M. Noltemeyer, *J. Am. Chem. Soc.* **2006**, 128, 12406-12407.
- (84) See for example: Z. Yang, H. Zhu, X. Ma, J. Chai, H. W. Roesky, C. He, H.-G. Schmidt, M. Noltemeyer, *Inorg. Chem.* **2006**, 45, 1823-1827.
- (85) J. D. Scollard, D. H. McConville, *J. Am. Chem. Soc.* **1996**, 118, 10008-10009.
- (86) B. Tsuie, D. C. Swenson, R. F. Jordan, J. L. Petersen, *Organometallics* **1997**, 16, 1392-1400.
- (87) L. Grocholl, V. Huch, L. Stahl, R. J. Staples, P. Steinhart, A. Johnson, *Inorg. Chem.* **1997**, 36, 4451-4457.
- (88) C. H. Galka, D. J. M. Trösch, I. Rüdener, L. H. Gade, I. Scowen, M. McPartlin, *Inorg. Chem.* **2000**, 39, 4615-4620
- (89) F. Ando, Y. Kohmura, J. Koketsu, *Bull. Chem. Soc. Jpn.* **1987**, 60, 1564-1566.

- (90) A. Jana, H. W. Roesky, C. Schulzke, A. Döring, *Angew. Chem.* **2009**, *121*, 1126-1129; *Angew. Chem., Int. Ed.* **2009**, *48*, 1106-1109.
- (91) U. N. Nehete, H. W. Roesky, V. Jancik, A. Pal, J. Magull, *Inorg. Chim. Acta* **2007**, *360*, 1248-1257.
- (92) T. A. Hanna, G. Keitany, C. Ibarra, R. D. Sommer, A. L. Rheingold, *Polyhedron* **2001**, *20*, 2451-2455.
- (93) O. M. Uy, J. Drowart, *Trans. Faraday Soc.* **1969**, *65*, 3221-3230.
- (94) N. C. Norman, *Chemistry of Arsenic, Antimony, and Bismuth*, Blackie Academic & Professional, London, **1998**.
- (95) S. Shimada, O. Yamazaki, T. Tanaka, S. Yohichi, M. Tanaka, *J. Organomet. Chem.* **2004**, *689*, 3012-3023.
- (96) G. G. Briand, N. Burford, M. D. Eelman, T. S. Cameron, K. N. Robertson, *Inorg. Chem.* **2003**, *42*, 3136-3141.
- (97) J. W. Pell, W. C. Davis, H. C. zur Loye, *Inorg. Chem.* **1996**, *35*, 5754-5755.
- (98) a) M. Mehring, *Coord. Chem. Rev.* **2007**, *251*, 974-1006. b) D. Meendoza-Espinosa, T. A. Hanna, *Inorg. Chem.* **2009**, *48*, 7452-7456. c) C. Knispel, C. Limberg, M. Mehring, *Organometallics* **2009**, *28*, 646-651. d) V. Stavila, J. H. Thurston, K. H. Whitmire, *Inorg. Chem.* **2009**, *48*, 6945-6951. e) B. Li, H. Zhang, L. Huynh, C. Diverchy, S. Hermans, M. Devillers, E. V. Dikarev, *Inorg. Chem.* **2009**, *48*, 6152-6158.
- (99) a) M. Hunger, C. Limberg, P. Kircher, *Angew. Chem.* **1999**, *111*, 1171-1174; *Angew. Chem., Int. Ed.* **1999**, *38*, 1105-1108. b) S. Roggan, C. Limberg, B. Zimmer, M. Brandt, *Angew. Chem.* **2004**, *116*, 2906-2910; *Angew. Chem., Int. Ed.* **2004**, *43*, 2846-2849.
- (100) J. H. Thurston, D. Trahan, T. Ould-Ely, K. H. Whitmire, *Inorg. Chem.* **2004**, *43*, 3299-3305.

- (101) a) P. M. Gurubasavaraj, H. W. Roesky, B. Nekoueishahraki, A. Pal, R. Herbst-Irmer, *Inorg. Chem.* **2008**, *47*, 5324-5331. b) S. K. Mandal, P. M. Gurubasavaraj, H. W. Roesky, R. B. Oswald, J. Magull, A. Ringe, *Inorg. Chem.* **2007**, *46*, 7594-7600. c) P. M. Gurubasavaraj, S. K. Mandal, H. W. Roesky, R. B. Oswald, A. Pal, M. Noltemeyer, *Inorg. Chem.* **2007**, *46*, 1056-1061. d) G. B. Nikiforov, H. W. Roesky, T. Schulz, D. Stalke, M. Witt, *Inorg. Chem.* **2008**, *47*, 6435-6443.
- (102) H. Gornitzka, D. Stalke, *Organometallics* **1994**, *13*, 4398-4405.
- (103) S. Singh, A. Pal, H. W. Roesky, R. Herbst-Irmer, *Eur. J. Inorg. Chem.* **2006**, 4029-4032.
- (104) L. W. Pineda, V. Jancik, H. W. Roesky, R. Herbst-Irmer, *Inorg. Chem.* **2005**, *44*, 3537-3540.
- (105) Y. Yang, H. W. Roesky, P. G. Jones, C.-W. So, Z. Zhang, R. Herbst-Irmer, H. Ye, *Inorg. Chem.* **2007**, *46*, 10860-10863.
- (106) G. Bai, S. Singh, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, *J. Am. Chem. Soc.* **2005**, *127*, 3449-3455.
- (107) B. Moraru, G. Kickelbick, U. Schubert, *Eur. J. Inorg. Chem.* **2001**, 1295-1301.
- (108) S. Parola, R. Papiernik, L. G. Hubert-Pfalzgraf, S. Jagner, M. Hakansson, *J. Chem. Soc., Dalton Trans.* **1997**, 4631-4635.
- (109) S. Roggan, C. Limberg, B. Ziemer, *Angew. Chem.* **2005**, *117*, 5393-5397; *Angew. Chem., Int. Ed.* **2005**, *44*, 5259- 5262.
- (110) K. C. H. Lange, T. M. Klapötke, In *The Chemistry of Organic Arsenic, Antimony, and Bismuth Compounds*; S. Patai, Ed.; Wiley: New York, 1994; Chapter 8.
- (111) K. Li, H. Xu, Z. Xu, M. Zeller and A. D. Hunter, *Inorg. Chem.* **2005**, *44*, 8855-8860.
- (112) H. Suzuki, T. Ikegami, Y. Matano, *Synthesis* **1997**, 249-267.

- (113) a) Z. Shen, J. Q. Zhang, H. X. Zou, M. M. Yang, *Tetrahedron Lett.* **1997**, 38, 2733. b) S. Repichet, C. Le Roux, J. Dubac, J. R. Desmurs, *Eur. J. Org. Chem.* **1998**, 2743-2746.
- (114) Y. Matano, H. Nomura, *Angew. Chem.* **2002**, 114, 3154-3157; *Angew. Chem., Int. Ed.* **2002**, 41, 3028-3031.
- (115) A. P. Soran, C. Silvestru, J. H. Breunig, G. Balazs, J. C. Green, *Organometallics* **2007**, 26, 1196-1203.
- (116) L. W. Pineda, V. Jancik, S. Nembenna, H. W. Roesky, *Z. Anorg. Allg. Chem.* **2007**, 633, 2205-2209.
- (117) J. Emsley, *Die Elemente*; Walter de Gruyter: Berlin, 1994.
- (118) H. Sitzmann, G. Wolmershäuser, *Z. Anorg. Allg. Chem.* **1999**, 625, 2103-2107.
- (119) M. Wieber, I. Sauer, *Z. Naturforsch.* **1984**, 39b, 887-889.
- (120) H. J. Breunig, D. Müller, *Z. Naturforsch.* **1986**, 41b, 1129-1132.
- (121) H. J. Breunig, K. H. Ebert, R. E. Schulz, M. Wieber, I. Sauer, *Z. Naturforsch.* **1995**, 50b, 735-744.
- (122) X.-W. Li, J. Lorberth, K. H. Ebert, W. Massa, S. Wocadlo, *J. Organomet. Chem.* **1998**, 560, 211-215.
- (123) H. J. Breunig, I. Ghesner, E. Lork, *J. Organomet. Chem.* **2002**, 664, 130-135.
- (124) M. Chovancova, R. Jambor, A. Ruzicka, R. Jirasko, I. Cisarova, L. Dostal, *Organometallics* **2009**, 28, 1934-1941.
- (125) F. Calderazzo, R. Poli, G. Pelizzi, *J. Chem. Soc., Dalton Trans.* **1984**, 2365.
- (126) O. Mundt, G. Becker, M. Rössler, C. Witthauer, *Z. Anorg. Allg. Chem.* **1983**, 506, 42-58.
- (127) W. Frank, V. Reiland, G. J. Reiss, *Angew. Chem.* **1998**, 110, 3154-3156; *Angew. Chem., Int. Ed. Engl.* **1998**, 37, 2983-2985.

- (128) See for further bindings modes: P. C. Junk, M. L. Cole, *Chem. Commun.* **2007**, 1579-1590.
- (129) M. Witt, H. W. Roesky, D. Stalke, F. Pauer, T. Henkel, G. M. Sheldrick, *J. Chem. Soc., Dalton Trans.* **1989**, 2173-2177.
- (130) A. Recknagel, M. Witt, F. T. Edelmann, *J. Organomet. Chem.* **1989**, 371, C40-C44.
- (131) H. Ackermann, O. Bock, U. Müller, K. Dehnicke, *Z. Anorg. Allg. Chem.* **2000**, 626, 1854-1856.
- (132) R. Boese, M. Düppmann, W. Kuchen, W. Peters, *Z. Anorg. Allg. Chem.* **1998**, 624, 837-845.
- (133) R. Vollmerhaus, R. Tomaszewski, P. Shao, N. J. Taylor, K. J. Wiacek, S. P. Lewis, A. Al-Humydi, S. Collins, *Organometallics* **2005**, 24, 494-507.
- (134) a) H. Schmidbaur, K. Schwirten, H.-H. Pickel, *Chem. Ber.* **1969**, 102, 564-567. b) W. Wolfsberger, W. Hager, *Z. Anorg. Allg. Chem.* **1976**, 425, 169-174. c) W. Wolfsberger, W. Hager, *Z. Anorg. Allg. Chem.* **1977**, 433, 241-254.
- (135) A. Steiner, D. Stalke, *Inorg. Chem.* **1993**, 32, 1977-1981.
- (136) R. Fleischer, D. Stalke, *Inorg. Chem.* **1997**, 36, 2413-2419.
- (137) E. Huheey, *Inorganic Chemistry*, 3rd edn., Harper and Row, New York, 1983.
- (138) H. R. Allcock, *Phosphorus-Nitrogen Compounds*, Academic Press, New York and London, 1972.
- (139) N. Kocher, D. Leusser, A. Murso, D. Stalke, *Chem. Eur. J.* **2004**, 10, 3622-3631.
- (140) J. F. Bickley, M. C. Copley, J. C. Jeffery, A. P. Leedham, C. A. Russell, D. Stalke, A. Steiner, T. Stey, S. Zacchini, *J. Chem. Soc., Dalton Trans.* **2004**, 989-995.
- (141) a) D. Leusser, J. Henn, N. Kocher, B. Engels, D. Stalke, *J. Am. Chem. Soc.* **2004**, 126, 1781-1793. b) N. Kocher, J. Henn, B. Gostevskii, D. Kost, I. Kalikhman, B. Engels, D. Stalke, *J. Am. Chem. Soc.* **2004**, 126, 5563-5568.

- (142) C. Cui, H. W. Roesky, M. Noltemeyer, M. F. Lappert, H.-G. Schmidt, H. Hao, *Organometallics* **1999**, *18*, 2256-2261.
- (143) a) H. Zhu, J. Chai, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, D. Vidovic, J. Magull, *Eur. J. Inorg. Chem.* **2003**, 3113-3119, and references therein. b) V. Jancik, M. M. Moya Cabrera, H. W. Roesky, R. Herbst-Irmer, D. Neculai, A. M. Neculai, M. Noltemeyer, H.-G. Schmidt, *Eur. J. Inorg. Chem.* **2004**, 3508-3512.
- (144) C. Cui, H. W. Roesky, H. Hao, H.-G. Schmidt, M. Noltemeyer, *Angew. Chem.* **2000**, *112*, 1885-1887; *Angew. Chem., Int. Ed.* **2000**, *39*, 1815-1817.
- (145) V. Jancik, Y. Peng, H. W. Roesky, J. Li, D. Neculai, A. M. Neculai, R. Herbst-Irmer, *J. Am. Chem. Soc.* **2003**, *125*, 1452-1453.
- (146) P. D. Godfrey, C. L. Raston, B. W. Skelton, V.-A. Tolhurst, A. H. White, *Chem. Commun.* **1997**, 2235-2236.
- (147) a) J. Müller, R. Schröder, R. Wang, *Eur. J. Inorg. Chem.* **2000**, 153-157. b) R.-C. Yu, C.-H. Hung, J.-H. Huang, H.-Y. Lee, J.-T. Chen, *Inorg. Chem.* **2002**, *41*, 6450-6455.
- (148) Y. Ying, T. Schultz, M. John, A. Ringe, H. W. Roesky, D. Stalke, J. Magull, Y. Hongqi, *Inorg. Chem.* **2008**, *47*, 2585-2592.
- (149) J. K. Ruff, M. F. Hawthorne, *J. Am. Chem. Soc.* **1961**, *83*, 535-538.
- (150) R. L. Stapleton, J. Chai, N. J. Taylor, S. Collins, *Organometallics* **2006**, *25*, 2514-2524.
- (151) M. Waggoner, M. M. Olmstead, P. P. Power, *Polyhedron* **1990**, *9*, 257-263.
- (152) C. H. Lee, Y. H. La, S. J. Park, J. W. Park, *Organometallics* **1998**, *17*, 3648-3655.
- (153) S. Singh, S. S. Kumar, V. Chandrasekhar, H.-J. Ahn, M. Biadene, H. W. Roesky, S. N. Hosmane, M. Noltemeyer, H.-G. Schmidt, *Angew. Chem.* **2004**, *116*, 5048-5051; *Angew. Chem., Int. Ed.* **2004**, *43*, 4940-4943.
- (154) S. Singh, V. Jancik, H. W. Roesky, R. Herbst-Irmer, *Inorg. Chem.* **2006**, *45*, 949-951.

6. References

- (155) L. W. Pineda, V. Jancik, H. W. Roesky, D. Neculai, A. M. Neculai, *Angew. Chem.* **2004**, *116*, 1443-1445; *Angew. Chem., Int. Ed.* **2004**, *43*, 1419- 1421.
- (156) P. M. Gurubasavaraj, H. W. Roesky, P. M. V. Sharma, R. B. Oswald, D. Dolle, A. Pal, *Organometallics* **2007**, *26*, 3346-3351.
- (157) B. Hübler-Blank, M. Witt, H. W. Roesky, *J. Chem. Educ.* **1993**, *70*, 408-409.

List of publications

- 1) *Angew. Chem.* **2009**, *121*, 4587-4590; *Angewandte Chemie International Edition*, **2009**, *48*, 4517–4520. “Addition of Dimethylaminobismuth to Aldehydes, Ketones, Alkenes, and Alkynes” **B. Nekoueishahraki**, S. Pillai Sarish, H. W. Roesky, D. Stern, C. Schulzke and D. Stalke.
- (2) *Inorganic Chemistry*, **2008**, *47*, 5324-5331. “From Unstable to Stable and Highly Active–Heterobimetallic Half-Metallocene Catalysts for Olefin Polymerization and Co-polymerization: Discussion on Structure/Reactivity Relationships” P. M. Gurubasavaraj, H. W. Roesky, **B. Nekoueishahraki**, S. K. Mandal, R. B. Oswald, A. Pal and R. Herbst-Irmer.
- (3) *Organometallics* **2009**, *28*, 5733-5738. “Synthesis and Structural Characterization of Heterobimetallic Bismuth Complexes with Main Group and Transition Metals” **B. Nekoueishahraki**, A. Jana, H. W. Roesky, D. Stern and D. Stalke.
- (4) *Inorganic Chemistry* **2009**, *48*, 9174-9179. “Synthesis and Structural Characterization of Aluminum Iminophosphonamide Complexes” **B. Nekoueishahraki**, H. W. Roesky, G. Schwab, D. Stern and D. Stalke.
- (5) *Organometallics* **2009**, *28*, 3763-3766. “Stable Compounds of Composition $LGe(II)R$ ($R = OH, PhO, C_6F_5O, PhCO_2$) Prepared by Nucleophilic Addition Reactions” A. Jana, **B. Nekoueishahraki**, H. W. Roesky and C. Schulzke.
- (6) *Inorganic Chemistry (Article)* (manuscript under preparation), “Organobismuth(III) and Dibismuthine Complexes Bearing N,N' -Disubstituted 1,8-Diaminonaphthalene Ligand: Synthesis, Structure and Reactivity” **Bijan Nekoueishahraki**, H. W. Roesky, Lallan Mishra, Daniel Stern, Dietmar Stalke and Carola Schulzke

Lebenslauf

Name	Bijan Nekoueishahraki	
Geboren	21. September 1974 in Shahrekord, Iran	
Staatsangehörigkeit	iranisch	
Schulbesuch	1980-1985	Primary School
	1985-1988	Secondary School
	1988 – 1992	High School
Studium	1993– 1997	B.Sc. Esfahan University, Iran
	2002 – 2005	M.Sc., Chamran University, Iran
Promotion	2006 Jan-present	Experimentelle Promotionsarbeit im Institut für Anorganische Chemie Göttingen unter der Leitung von Prof. Dr. Dr. h.c. mult. H. W. Roesky