AUS DER ABTEILUNG DERMATOLOGIE, VENEROLOGIE UND ALLERGOLOGIE
(PROF. DR. MED. M. P. SCHÖN)
IM ZENTRUM ARBEITS-, SOZIAL-, UMWELTMEDIZIN UND DERMATOLOGIE
DER MEDIZINISCHEN FAKULTÄT DER UNIVERSITÄT GÖTTINGEN

Platelet surface receptors and melanoma metastasis

INAUGURAL – DISSERTATION

zur Erlangung des Doktorgrades
der Medizinischen Fakultät der
Georg-August-Universität zu Göttingen

vorgelegt von

Luise Erpenbeck

aus

Berlin

Göttingen 2010

Dekan: Prof. Dr. med. Frömmel

I. Berichterstatter: Prof. Dr. med. Schön

II. Berichterstatter: Prof. Dr. med. Dobbelsstein

III. Berichterstatter: Prof. Dr. med., Dr. rer. nat. Crozier

Tag der mündlichen Prüfung: 18.5.2011

Contents

Abbreviations	II
1 Summary	1
2 Introduction	3
2.1 Melanoma: pathophysiology, epidemiology and prognosis	3
2.2 Platelets and platelet receptors in the process of hematogenous metastasis	4
2.3 The platelet receptor GPVI	9
2.3.1 GPVI and metastasis	9
2.3.2 Physiological role of GPVI	9
2.3.3 Structure of and signaling through GPVI	9
2.4 The platelet receptor GPIb-V-IX	12
2.4.1 GPIb-V-IX in metastasis	12
2.4.2 Physiological role of GPIb-V-IX	12
2.4.3 Structure of and signaling through GPIb-V-IX	13
2.5 Aim of this work	16
3 Materials and Methods	17
3.1 Antibodies	17
3.2 Cells and cell culture conditions	18
3.3 Mice	18
3.4 Flow cytometric assessment of GPIb α expression on platelets and B16BL6 melanoma cells	18
3.5 Immunohistochemistry	19
3.6 Aggregometry	19
3.7 Assessment of receptor occupancy	21
3.8 Experimental pulmonary metastasis	21
3.9 Analysis of metastatic areas in lung sections	23
3.10 Short term metastasis	23
3.11 Immunohistochemical staining against Mac-1	24
3.12 Apoptosis assay	25
3.13 Chemicals and laboratory equipment	26

4	Results	29
4.1	Antibody-mediated blockade of platelet GPIb α and its impact on TCIPA	29
4.1.1	Assessment of GPIb α expression on platelets	29
4.1.2	Immunohistochemical analysis of GPIb α expression in lung tissue	30
4.1.3	Tumor cell-induced platelet aggregation (TCIPA) under GPIb α blockade	31
4.1.4	Assessment of receptor occupancy	31
4.2	GPIb α and pulmonary metastasis	33
4.2.1	Experimental pulmonary metastasis after platelet depletion . . .	33
4.2.2	Pulmonary metastasis under GPIb α blockade	34
4.2.3	Critical time frame for GPIb α impact on metastasis	39
4.2.4	Short term metastasis	40
4.2.5	Mac-1-expressing immune cells after short-term metastasis . . .	45
4.2.6	Double blockade of GPIb α with Mac-1	45
4.2.7	Blockade of GPIb α and P-selectin in experimental melanoma metastasis	49
4.2.8	Impact of vWF on melanoma survival <i>in vitro</i>	51
4.3	Role of GPVI in metastasis	52
4.3.1	Experimental pulmonary metastasis under GPVI blockade . . .	52
5	Discussion	55
5.1	GPIb α is expressed only on platelets and does not alter tumor cell- induced platelet aggregation (TCIPA)	55
5.2	Blockade of GPIb α promotes experimental pulmonary metastasis in mice	56
5.3	Interaction of the integrin receptor Mac-1 with GPIb α is not responsible for the promotion of metastasis upon GPIb α blockade	60
5.4	GPIb α blockade does not lead to enhanced metastasis in P-selectin de- ficient mice	62
5.5	GPVI in experimental metastasis	66
5.6	Conclusion	66
6	Bibliography	68
	List of Figures	IV
	List of Tables	V
	Previously published results	VI

Abbreviations

ABP	Actin-binding protein
ADP	Adenosine diphosphate
BSA	Bovine serum albumin
CRP	Collagen-related peptide
ECM	Extracellular matrix
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme-linked Immunosorbent Assay
FACS	Fluorescence-activated cell sorting
FCS	Fetal calf serum
FITC	Fluorescein isothiocyanate
GFP	Green fluorescent protein
GP	Glycoprotein
H&E	Hematoxylin and eosin
HMWK	High molecular weight kininogen
HRP	Horseradish peroxidase
ITAM	Immunoreceptor tyrosine-based activation motif
LRR	Leucine-rich repeat
NK cell	Natural killer cell
PBS	Phosphate buffered saline
PRP	Platelet rich plasma
TCIPA	Tumor cell-induced platelet aggregation
TGF-beta	Transforming growth factor beta
UVR	Ultra Violet Radiation
VEGF	Vascular endothelial growth factor
vWF	von Willebrand Factor

1 Summary

Background and Design

In the last decades, accumulating experimental evidence has suggested a contribution of platelet receptors as well as factors of the blood coagulation system to hematogenous tumor metastasis of melanoma, a highly aggressive tumor arising from pigment cells of the skin. In fact, inhibition of either soluble factors involved in the coagulation cascade or blockade of platelet surface receptors which play a role in primary hemostasis have always lead to diminished metastasis in many experimental models. These results have raised expectations for novel cancer therapies which would be able to impede metastatic disease in high risk patients or in situations associated with an increased risk of tumor cell dissemination.

In this context, I have studied the role of the platelet surface receptor, GPIIb/IIIa, in experimental pulmonary metastasis. Experiments were aimed at unveiling the mechanisms underlying the involvement of this receptor in metastasis. Its role as a possible therapeutic target in the treatment of cancer was also evaluated.

Methods

I have tested the effect of GPIIb/IIIa inhibition by monoclonal monovalent anti- GPIIb/IIIa (p0p/B) Fab-fragments, which do not induce thrombocytopenia, on experimental pulmonary metastasis. Metastasis assays were performed in syngeneic C57BL/6 mice, the melanoma cells employed were B16F10 melanoma cells. To assess the early fate of melanoma cells under GPIIb/IIIa -blockade, B16F10 melanoma cells were stably transfected with GFP (Green fluorescent protein), which allowed tracking of tumor cells within the lung vasculature directly after injection. Retention of melanoma cells in the lungs was assessed by fluorescence microscopy and by flow cytometry. A possible stimulation or inhibition of the innate immune defense in these short-term metastasis assays was studied by immunohistochemical staining against the integrin receptor Mac-1 (CD11b/CD18). To assess a possible interference of anti-GPIIb/IIIa antibodies with tumor cell-induced platelet aggregation, aggregometry was performed using platelet rich plasma (PRP), which had previously been incubated with anti-GPIIb/IIIa, and B16F10

melanoma cells as inductor of aggregation. Finally, the effect of GPIIb α inhibition on experimental pulmonary metastasis was evaluated in mice lacking P-selectin.

Results

Surprisingly, GPIIb α inhibition led to a significant increase of pulmonary metastasis. Assessment of the early fate of circulating GFP-labeled B16F10 cells showed improved pulmonary arrest of the tumor cells within the lung vasculature shortly after GPIIb α inhibition. Tumor cell-induced platelet aggregation was not influenced by the addition of monoclonal, monovalent anti- GPIIb α . Interestingly, GPIIb α -blockade in mice deficient for P-selectin (a GPIIb α ligand) had no enhancing effect on metastasis.

Conclusion

This is the first experimental evidence showing that inhibition of a platelet receptor may not always impair tumor metastasis, but may indeed result in increased metastasis of a malignant tumor. The results of this work support the hypothesis that the extracellular domain of GPIIb α contributes to the control of metastatic tumor progression, in addition to its role in hemostasis. This novel function of GPIIb α is presumably involved in the promotion of the initial, P-selectin-dependent steps of metastasis, as GPIIb α -blockade in P-selectin deficient mice showed no enhancement of the number of metastatic foci.

2 Introduction

2.1 Melanoma: pathophysiology, epidemiology and prognosis

Melanoma is an aggressive, therapy-resistant malignancy of melanocytes. Once metastatic spread has occurred, the tumor is almost universally resistant to therapy.

Melanocytes originate from neural crest cells and later differentiate and migrate from the neural crest to several tissues during the early weeks of gestation. This includes the skin, where their function is best studied, but also the leptomeninges, the retina and mucosal tissues (Markovic et al. 2007a).

In the skin, melanocytes localize within the basal layer of the epidermis, where they are capable of producing and releasing melanin. Melanin absorbs ultraviolet radiation (UVR) and possesses an antioxidant potential which enables it to neutralize UVR-generated free radicals. Melanin is transferred to keratinocytes where melanin granules orient over the keratinocyte nucleus and thus protect them from UV damage to the DNA and help prevent skin cancers such as squamous cell and basal cell carcinoma. Unfortunately, melanocytes themselves are vulnerable to UVR and can undergo malignant transformation, resulting in a much more aggressive cancer type than those they protect the organism against. Risk factors to develop melanoma include a large number of melanocytic naevi, especially of dysplastic forms, a light skin type and a genetic disposition. Exposure to UVR is considered a potential trigger of melanoma, especially sun burns in childhood may lead to malignant transformation in later life (Markovic et al. 2007a).

The incidence of melanoma has been steadily increasing worldwide over the last decades, with the incidence in Germany having more than tripled since 1980 and patients becoming ever younger at the diagnosis of the disease. This may be due to a higher exposure to UVR, possibly caused by the depletion of the ozone layer as well as behavioral factors such as outdoor sports and sun tanning, but also by increased surveillance. Fortunately, no increase in mortality due to melanoma has been observed (thus cor-

roborating the latter notion). This might be explained by a better education of the population with respect to pigment cell lesions and, thus, a diagnosis of early forms of cutaneous melanoma which can be removed easily and are characterized by a high cure rate. Currently, 14,900 people in Germany are diagnosed with melanoma every year. In Germany, melanoma accounts for 4.1 % of all neoplasms in women and for 2.8 % of neoplasms in men, with women having an average age of 57 and men of 63 years at the onset of the disease. Given that melanoma can be detected and removed at early stages, the overall cure rates for melanoma are relatively high. The 5-year survival rate is 88 % for women and 84 % for men and melanoma is responsible for only 1 % of cancer-related deaths in both sexes (Batzler et al. 2008).

At the other end of the spectrum, prognosis of patients with metastasized melanoma remains exceedingly poor. When the tumor has reached stage IV, meaning that the tumor has formed distant metastases in skin, subcutis, lymph nodes or distant organs like the lung or other visceral organs, the median survival is less than 9 months and 5-year survival probability shrinks to less than 5 %, notwithstanding numerous trials using chemotherapy, immunotherapy or other regimes. To date, only 2 substances have been approved by the Food and Drug Administration (FDA) for treatment of metastasized melanoma: Dacarbazine and IL-2, both of which show moderate response rates and neither of which prolongs overall survival in a significant way (Markovic et al. 2007b). Several new substances and novel combinations of established chemotherapeutics are being tested to improve tumor response rates and patient survival. However, it is clear that future research must be aimed at unveiling the processes underlying metastasis to facilitate the development of agents that will specifically inhibit those steps that turn a tumor disease which can be relatively successfully treated into a malignancy with an infaust prognosis.

2.2 Platelets and platelet receptors in the process of hematogenous metastasis

In most patients suffering from cancer, metastasis remains the main cause of death. As mentioned above, melanoma is an especially impressive example of this notion, as metastasized melanoma is almost universally resistant to chemotherapy. The process of hematogenous metastasis is a highly complex cascade of events, including the detachment of cells from the primary tumor, the transport of the metastatic cell along the blood stream, the retention in the vasculature of the metastatic site and the proliferation in the distant tissue. Upon entering the blood stream, the tumor cells encounter a highly hostile environment in which they must thwart the immune surveillance of

the host and resist high mechanical tensions in the blood flow. Astonishingly, however, these first steps of metastasis, namely the extravasation into the blood stream, arrest in the capillary bed of the secondary organ and extravasation at the metastatic site, are completed very efficiently, as can be shown in animal models. For example, in an experimental model where B16F1 murine melanoma cells were injected into the mesenteric vein of C57BL/6 mice, over 87 % of the cells were arrested in the liver and had survived 90 minutes after injection (Luzzi et al. 1998). The limiting steps in the development of clinically detectable metastases seem to be the formation of micrometastases, subsequent vascularization and growth. Yet this not only indicates that the success determining steps in metastasis occur after extravasation, but also that tumor cells have developed very efficient mechanisms to protect themselves while exposed to the blood components of the host. Nevertheless, the passage through the blood of the tumor cells could constitute an important selection step as cells which cannot protect themselves against the immune system or are less resistant to shear stress within the vasculature would be more easily destroyed and would never even have a chance to initiate the formation of micrometastases (Pantel and Brakenhoff 2004).

During the early phase of hematogenous metastasis, platelets appear to play a crucial role. As early as 1865, the first indication for an involvement of platelets in metastasis was described by A. Trousseau who recognized the relation between thromboembolism and cancer (Trousseau 1865). In fact, the association of cancer and thrombosis has a prevalence rate of as much as 10 % – 20 % (Otten et al. 2004). In 1986, the ability of tumor cells to induce platelet aggregation *in vitro*, a process termed tumor cell-induced platelet aggregation (TCIPA), was discovered by Gasic et al (Gasic et al. 1968) who could also show that the aggregation caused by the tumor cells directly correlates to their metastatic potential *in vivo*. It is thought that platelet aggregation provides several advantages for tumor cells implicated in the process of metastasis (see figure 2.1).

It is possible that a protective thrombus surrounding the cancer cells shields them from the host immune system (Nieswandt et al. 1999; Palumbo et al. 2005). A number of studies support the concept that platelets can limit the ability of natural killer cells to lyse tumor cells *in vitro* and *in vivo*, an observation which is reversed after NK cell depletion (Nieswandt et al. 1999; Palumbo et al. 2005). Very recent studies have shown that platelet-derived transforming growth factor beta (TGF-beta), secreted upon platelet activation e.g. by tumor cells, down-regulates the activating immunoreceptor NKG2D on NK cells (Kopp et al. 2009). Furthermore, it is conceivable that platelets facilitate the interactions of tumor cells with the endothelium of the vasculature at the metastatic site (Ludwig et al. 2004) and thus promote the arrest of the tumor cells

Fig. 2.1: Proposed advantages of tumor cell interactions with platelets.

Interaction of tumor cells with platelets within the blood stream may provide the tumor cells with many advantages. These could be protection from the immune system, arrest of the tumor cells within the vasculature of the metastatic organ through facilitated interaction with endothelial cells or through size restriction because of the surrounding thrombus. Platelets can also release proangiogenic and growth promoting factors such as VEGF and Angiopoietin-1.

within the vasculature (Kemp et al. 1971), possibly through the formation of tumor cell-platelet emboli. Finally, growth factors can be secreted by platelets which may also support tumor survival and growth, such as platelet derived growth factor, vascular endothelial growth factor (VEGF) and Angiopoietin-1 (Kepner and Lipton 1981; Mohle et al. 1997; Nierodzik and Karparkin 2006). Taking into account these diverse promotional effects of platelets in metastasis, it is not surprising that platelet depletion in *in vivo* models of metastasis always resulted in an abrogation of cancer cell dissemination (Gasic et al. 1968; Gasic 1984).

Molecules such as thrombin (Nierodzik et al. 1991), fibrinogen (Biggerstaff et al. 1999; Palumbo et al. 2000), von Willebrandt factor (vWF, Terraube et al. 2007), tissue factor or lysophosphatidic acid (Camerer et al. 2004) were all found to affect metastasis in such a way that their depletion or functional blockade severely diminished the metastatic

potential of tumor cells.

The platelet receptors involved in coagulation appear to be equally important. Under normal physiological conditions, these receptors allow a rapid recruitment of platelets to the site of vascular injury where they interact with components of the subendothelial extracellular matrix (ECM) . This interaction leads to the production of soluble factors that strengthen platelet adhesion, subsequent receptor activation and further recruitment of platelets, all of which finally result in the formation of a stable thrombus (Jackson et al. 2003; Andrews et al. 2004).

In the metastatic cascade, the same platelet receptors become indispensable for the metastasizing tumor cell. A prominent example is the integrin receptor $\alpha_{IIb}\beta_3$ (GPIIb/IIIa), a molecule which becomes activated upon platelet stimulation and then leads to platelet aggregation through binding of fibrinogen, fibronectin and vWF (Hynes 2002; Shattil and Newman 2004). This integrin has long been known to support metastasis *in vitro* and *in vivo* (Bastida et al. 1987; Grossi et al. 1987; Karparkin et al. 1988). A similarly important role in metastasis has been established for the thrombin PAR-receptors, which, among other things, physiologically initiate platelet shape change, upregulation of surface P-selectin expression, integrin activation as well as release of platelet fibrinogen and vWF (Davey and Luscher 1967; Hamberg et al. 1975; Henn et al. 1998). While thrombin stimulation of platelets increases their adhesion to tumor cells *in vitro* (Nierodzik et al. 1991), mice whose platelets fail to respond to thrombin show a clear reduction in metastasis *in vivo* (Camerer et al. 2004). Similar observations providing evidence for a participation of platelet receptors in metastasis also exist for the ADP-receptors P2Y(1) and P2Y(12) (Davey and Luscher 1967; Alonso-Escolano et al. 2004) and for P-selectin (Kim et al. 1998; Ludwig et al. 2004).

An overview of platelet receptors implicated in metastasis, their binding partners as well as their potential interactions with tumor cells can be seen in figure 2.2. Of note, some platelet receptors such as the integrin receptors or PAR-receptors can also be expressed on tumor cells and increase their metastatic potential (Trikha et al. 1998; Trikha et al. 2002; Melnikova et al. 2008).

Fig. 2.2: Platelet receptors involved in hematogenous tumor metastasis. Schematic representation of platelet surface molecules whose primary functions contribute to hemostasis and coagulation and which have been found to be involved in hematogenous metastasis. Generation of genetically engineered mice lacking these receptors has, in most cases, led to an attenuation of metastasis. Some platelet receptors are also expressed and utilized by the tumor cells where they contribute to the metastatic potential. Black arrows indicate binding of the respective receptor with the corresponding ligand and dashed lines represent other (not specified) interactions.

2.3 The platelet receptor GPVI

In contrast to the above mentioned findings, the roles of the GPIb-V-IX-complex and GPVI, two platelet receptors essential for the initiation of platelet interactions with the ECM, remain very poorly investigated as far as their role in metastasis is concerned.

2.3.1 GPVI and metastasis

In the case of GPVI, no involvement in metastasis has hitherto been shown.

2.3.2 Physiological role of GPVI

The lack of a known role of GPVI in metastasis would seem surprising, considering that recent publications have identified GPVI as one of the principal collagen receptors in platelets (Nieswandt et al. 2001a; Massberg et al. 2003; Nieswandt and Watson 2003). In general, collagen plays a major role in the initial recruitment of platelets to the injured vessel wall, especially under conditions of medium or high shear rates that may be found in arteries and damaged vessels. Binding of collagen to the glycoprotein receptor GPVI is by itself insufficient for stable adhesion. However, this interaction is one of the most potent inducers of "inside-out signaling", causing the integrin receptors $\alpha_2\beta_1$ and $\alpha_{IIb}\beta_3$ to undergo conversion to the high affinity conformation. Interaction of the integrin receptor $\alpha_2\beta_1$ with collagen as well as interaction of the integrin $\alpha_{IIb}\beta_3$ with vWF then lead to firm adhesion of platelets to the ECM. This stable binding is reinforced by the release of soluble mediators like thrombin, thromboxane A2 and ADP.

2.3.3 Structure of and signaling through GPVI

The structure of GPVI was reported by Clemetson et al. in 1999 following the cloning of the receptor and identifying it as a member of the Ig receptor superfamily. Its extracellular chain carries two Ig-C2-like domains formed by disulfide bridges, linked to a mucin-like region that contains several sites for O-linked glycosylation. An arginine group of the transmembrane region is required for the association with the FcR γ -chain through a salt bridge (Clemetson et al. 1999). The cytosolic tail of the GPVI-chain contains a proline-rich domain that binds to the Src homology 3 (SH3) domain of the Src family tyrosine kinases, Fyn and Lyn (Suzuki-Inoue et al. 2002). The FcR γ -chain is a disulfide-linked homodimer and contains two conserved sequences known as immunoreceptor tyrosine-based activation motif (ITAM-motifs, see figure 2.3).

Activation of GPVI by collagen is believed to induce cross-linking of collagen receptors which then brings the Src kinases in close proximity of two ITAM motifs localized on

Fig. 2.3: The platelet receptor GPVI. Schematic representation of the GPVI receptor associated to the FcR γ -chain. GPVI consists of 2 Ig-C2-like domains linked to a mucin-like region with several sites for O-linked glycosylation. The transmembrane domain has an arginine residue which is necessary for the association with the FcR γ -chain through a salt bridge. The cytosolic chain of the FcR γ -chain contains two ITAM-motifs each (adapted from Nieswandt and Watson 2003, page 452).

the FcR γ -chain. The signaling cascade that is thus initiated and which also includes the activation of another tyrosine kinase, Syk, eventually leads to tyrosine phosphorylation of a number of downstream proteins including linker of activated T cells (LAT) and phospholipase C γ 2 (PLC γ 2; Ezumi et al. 1998).

2.4 The platelet receptor GPIb-V-IX

2.4.1 GPIb-V-IX in metastasis

The putative role of GPIb-V-IX, though investigated by several research groups, has not been clearly identified, either. In fact, many results in this field are quite contradictory, depending strongly on tumor cell line and experimental setting. While some research groups have described an involvement of GPIb α , the largest subunit of the GPIb-V-IX complex (see figure 2.4), in TCIPA (Bastida et al. 1987; Grossi et al. 1987), no influence on this process could be observed by others (Karparkin et al. 1988). *In vivo*, an abrogation of metastasis was seen in GPIb α -deficient mice (Jain et al. 2007). However, these experiments bear a considerable challenge as mice lacking GPIb α suffer from a severe bleeding phenotype together with giant platelets which resembles the human Bernard Soulier syndrome (Kato et al. 2004) (see below). Even smaller modifications of GPIb α -expression might lead to aberrant membrane development through the interference with megakaryocyte maturation (Kanaji et al. 2002; Poujol et al. 2002). Considering that functional platelets are a prerequisite for successful metastasis, models that alter platelet membrane integrity or platelet count should be critically evaluated.

2.4.2 Physiological role of GPIb-V-IX

Under physiological conditions, interaction of GPIb-V-IX with the subendothelial von Willebrand factor (vWF) is essential for primary hemostasis. Upon damage of the vessel wall, von Willebrand factor (vWF), circulating in the blood stream, binds to exposed components of the subendothelial tissue and via its interaction with GPIb-V-IX initiates platelet adhesion to the subendothelial matrix. Especially under conditions of high shear which can be found in arterioles and small arteries, this mechanism appears to be important.

Similarly to GPVI, ligand binding of GPIb-V-IX by itself is not sufficient to convey stable platelet adhesion. However, signaling through the receptor leads to a series of events that finally enables thrombus formation. It exerts its role in hemostasis mainly through interaction with the A1 domain of vWF, a large glycoprotein stored in platelet α -granules and Weibel-Palade bodies of endothelial cells, from which it can be released upon activation. While dimeric forms of vWF circulate in the plasma, carrying coagulation factor VIII, multimeric forms are immobilized in the ECM, allowing rapid and reversible platelet recruitment upon exposure of the subendothelial matrix. While this interaction alone is not sufficient to induce stable platelet adhesion, it consequently results in integrin-mediated platelet adhesion and activation. While interaction with vWF is presumably the main function of GPIb-V-IX, there are a number of additional extracellular ligands GPIb-V-IX can bind to. These ligands include thrombin, clotting

factors XI and XII, high molecular weight kininogen (HMWK), P-selectin, the leukocyte integrin Mac-1 and thrombospondin-1. The binding sites for all these ligands are located in the extracellular N-terminal domain of GPIb α .

Considering the central role of the GPIb-V-IX-complex in primary hemostasis, further investigations of the influence of the GPIb-V-IX-complex on metastasis would seem justified and necessary.

2.4.3 Structure of and signaling through GPIb-V-IX

GPIb-V-IX is a complex of glycoproteins of the leucine-rich repeat (LRR) family, constitutively expressed on the platelet membrane. The four transmembrane proteins that compose the receptor complex, GPIb α , GPIb β , GPIX and GPV, are displayed in a ratio of 2:2:2:1. While GPIb α and GPIb β are linked by a disulfide-bond, GPV and GPIX are non-covalently associated to the complex.

The N-terminal domain of GPIb α , delimited by two conserved disulphide loop structures and composed by 8 LRRs, is the major ligand-binding region of GPIb-IX-V, containing partially overlapping binding sites for vWF, the leukocyte integrin Mac-1, thrombin, P-selectin, clotting factors XI and XII, thrombospondin-1 as well as HMWK. (Romo et al. 1999; Simon et al. 2000; Andrews et al. 2003). It is followed by a region of enriched negatively charged residues, including aspartic and glutamic acids as well as three sulfated tyrosines, a rather long mucin-like macroglycopeptide domain which is highly glycosylated, a single transmembrane region and the cytoplasmic tail. This cytoplasmic tail contains binding sites for proteins of the cytoskeleton such as the actin-binding protein (ABP), thus conveying an anchorage to the actin filament network of the platelet. It also mediates binding to intracellular signaling molecules such as 14-3-3 ζ (Du et al. 1996).

GPIb β and GPIX both contain only one LRR with conserved flanking sequences. The cytoplasmic tail of GPIb β has been found to interact with 14-3-3 ζ as well as with calmodulin (Wardell et al. 1989). While association of GPIb-V-IX with 14-3-3 ζ seems to be important for $\alpha_{IIb}\beta_3$ activation and for GPIb-V-IX binding to the PI-3K (Munday et al. 2000), the role of calmodulin binding to GPIb-V-IX remains altogether unclear. GPV has been shown to strengthen the interactions of vWF and GPIb-V-IX under conditions of high shear. Furthermore, it has the capability to bind thrombin as well as collagen (Fredrickson et al. 1998; Moog et al. 2001). It contains 15 LRRs in its extracellular domain, and its cytoplasmic tail again carries binding regions for 14-3-3 ζ and calmodulin. Activation of GPIb-V-IX is a complex process which finally leads to $\alpha_{IIb}\beta_3$ activation and stable aggregation. Platelet activating events are dependent on the stimulation of tyrosine kinases, including Syk, FAK, and Pyk2 and the activation of

Fig. 2.4: The platelet receptor complex GPIb-V-IX. The complex GPIb-V-IX consists of four different transmembrane polypeptides: GPIb α , GPIb β , GPIX, and GPV in a ratio of 2:2:2:1. GPIb α is covalently linked to GPIb β by a disulphide group, GPIX and GPV are non-covalently associated with the surrounding proteins. Leucine-rich repeats of all four proteins are represented by grey circles. The red minus sign represents the region of negatively charged residues of GPIb α . Carbohydrate moieties are depicted as black lines. The cytoplasmic domains of the single subunits interact with a number of proteins including calmodulin (in yellow, as indicated), the 14-3-3 ζ protein (grey semi-circle) and the cytoskeletal actin-binding protein (in yellow, as indicated), allowing an anchorage to the actin filament-network (adapted from Andrews et al. 2003, page 1173).

PLC, PLA2, and PI-3K. The recruitment of some of these effectors requires the phosphorylation of the ITAM motifs of the Fc receptor $\text{Fc}\gamma\text{RIIA}$ and $\text{FcR}\gamma$ -chains, made possible by the close proximity of these receptors to GPIb-V-IX in lipid rafts on the platelet surface. Cytoskeletal proteins also appear to be engaged in the formation of this signaling complex. Activation of different protein kinases as well as production of second messengers finally lead to the above mentioned integrin activation.

2.5 Aim of this work

The present work aims at investigating the roles of GPVI and GPIb-V-IX, more specifically GPIb α , in the process of hematogenous metastasis, considering the central position both receptors occupy in the initiation of primary hemostasis. Of particular interest were the determination of the *in vivo* relevance of both receptors in a model of experimental metastasis and the elucidation of possible mechanisms underlying any observations made in this model. Furthermore, it was evaluated whether the obtained results could lead to novel therapeutic targets in cancer treatment.

3 Materials and Methods

3.1 Antibodies

Hybridomas producing the following monoclonal antibodies were cultured in our lab: Monoclonal anti-CD11b M1/70 (Mac-1, α M-integrin) (ATCC, Rockville, MD).

Antibodies were purified using a protein G column (Hi Trap Protein G HP, GE Healthcare, München, Germany). In short, hybridoma supernatant was collected. The Protein G column was washed with 100 ml PBS (pH 7.3, PAA Laboratories GmbH, Cölbe, Germany), cleaned with 50 ml 1 M acetic acid (pH 2.5) and washed again with 200 ml PBS, using a Harvard Pump 11 Plus Advanced Single Syringe with Dual RS-232 (Harvard Apparatus, Holliston, USA).

Hybridoma supernatant was then perfused over the column, fallthrough was collected in a sterile flask. The column was then washed with binding buffer - 20 mM Na_2HPO_4 , 20 mM NaH_2PO_4 (both Roth, Karlsruhe, Germany), pH 7. Protein was eluted by perfusing the column with elution buffer - 0.1 M glycine-hydrochloride, pH 2.7 (Sigma Aldrich, Taufkirchen, Germany). The eluate was caught in Eppendorf tubes equipped with 50 μl of 1 M Tris(hydroxymethyl)aminomethane-buffer (Tris-buffer, Sigma Aldrich), pH 3. The column was equilibrated with 100 ml PBS, washed with 50 ml 20 % ethanol and stored in 20 % ethanol. Determination of protein concentration in the eluate was performed using a Bradford protein assay (Bio-Rad Laboratories GmbH, München, Germany). Antibody was then pooled and dialyzed against PBS for at least 4h using Slide-A-Lyzer-Cassettes (Pierce Protein Research Products, Thermo Fisher Scientific Inc., Rockford, USA), and sterile filtered. Finally, the protein concentration was measured again using the Bradford protein assay.

The following monoclonal antibodies, including the respective fusion proteins with fluorescein isothiocyanate (FITC) or Fab-fragments were generously provided by Prof. B. Nieswandt (Nieswandt et al. 2000a): p0p/B (directed against the N-terminal 45-kDa portion of the GPIb receptor), p0p1 (directed at the membrane-anchored part of the GPIb α receptor which remains after proteolytic cleavage of GPIb α), p0p3 and p0p4 (with binding sites located on the N-terminal 45-kD portion of the GPIb α receptor, Bergmeier et al. 2000), JAQ1 (anti-GPVI; Nieswandt et al. 2000b), JON/A (directed

against the high affinity conformation of GPIIb/IIIa; Bergmeier et al. 2002). Rat IgG2b control monoclonal antibody (ATCC) was also provided by Prof. Nieswandt.

3.2 Cells and cell culture conditions

B16F10 murine melanoma cells (Hart 1979) were used for flow cytometric evaluation of GPIb α expression on melanoma cells, all long-term experimental metastasis assays as well as for aggregometry and apoptosis assays. These melanoma cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, PAA Laboratories GmbH), high glucose, with 10% fetal calf serum (FCS), 2 mM L-glutamine, 100 μ U/ml streptomycin and 100 U/ml penicillin (both Gibco, Invitrogen GmbH Karlsruhe, Germany). The cells were kept in 75 cm² cell culture flasks (Greiner Bio-One GmbH, Frickenhausen, Germany) in an incubator at 37°C with 5 % CO₂.

3.3 Mice

C57BL/6 wildtype mice as well as P-selectin knockout mice on a C57BL/6 genetic background were obtained from Charles River (Sulzfeld, Germany). Mice were kept in a specific pathogen-free environment on a 12 hours light-dark cycle at the animal facility of the Rudolf Virchow Center. Mice were between 6 and 9 weeks old when used for the experiments, all groups were sex and age-matched. Animal experiments were approved by the Regierung von Unterfranken.

3.4 Flow cytometric assessment of GPIb α expression on platelets and B16BL6 melanoma cells

To investigate whether B16F10 melanoma cells expressed GPIb α , subconfluent grown melanoma cells were detached from the surface of a T75-flask using a cell scraper (Greiner Bio-One GmbH), in order to avoid interference of trypsin with surface molecules. Cells were washed in phosphate buffered saline (PBS, PAA Laboratories) 2 times and spun down at 1300 rpm for 3 min. 1×10^6 B16F10 cells were then suspended in 50 μ l PBS and stained with 0,5 μ g p0p/B-FITC for 30 min at 4°C. Cells were washed again at 1300 rpm for 3 min and suspended in 1 ml FACS-buffer (1 g BSA 0,1 g NaN₃ in 100 ml PBS). Mean fluorescence was measured in a FACScalibur (BD Biosciences, Heidelberg, Germany). As control, platelet surface expression of GPIb α was evaluated by flow cytometry. For generation of platelet-rich-plasma (PRP), C57BL/6 mice were bled under ether anesthesia from the retroorbital plexus using short glass capillaries (Brand GmbH & Co.KG, Wertheim, Germany). Approximately 700 μ l of blood were collected

in a 1,5 ml Eppendorf tube which contained 300 μ l heparin (Liquemin N25000, Roche Diagnostics GmbH, Penzberg, Germany) at 20 U/ml. The blood was spun at 1800 rpm for 5 min. The resulting supernatant was centrifuged again at 900 rpm for 6 min. 50 μ l of PRP was incubated with 0.5 μ p0p/B-FITC for 10 min at room temperature (incubation of platelets on ice was avoided to prevent clustering of GPIb on the platelet surface). Platelets were centrifuged again at 900 rpm for 6 min and then resuspended in 1 ml FACS-buffer. Receptor surface presentation was then measured by flow cytometry.

3.5 Immunohistochemistry

For immunohistochemistry, 5 μ m cryostat-cut sections were stained by the ABC-immunoperoxidase method (Vector Labs, Burlingame, USA). Briefly, air-dried sections were fixed in acetone for 20 seconds at 4°C and then incubated with buffer containing 1 % normal goat serum and 1 % normal rabbit serum for 20 min at room temperature in a moist chamber. Sections were then incubated with 5 μ g/ml of the primary antibody at room temperature in a moist chamber for 45 min, followed by 0.3 % H₂O₂ (Roth) in PBS for 20 min to block endogenous peroxidase. After washing in PBS twice, the slides were incubated with biotinylated secondary antibody, according to the primary antibody used, for 30 min at room temperature, again in the moist chamber. After washing twice, streptavidin-labeled with horseradish peroxidase (HRP, Dako, Hamburg, Germany) was added to the slides for 3 min. In the case of p0p1 and p0p/B, the primary antibodies were already HRP labeled, making an incubation with a secondary antibody unnecessary. After careful washing in PBS, the slides were submerged in 3-amino-9-ethylcarbazole substrate solution in 0.1 M acetate buffer (pH 5.2). Color development was stopped by 10 % formalin (pH 5.2) for 10 min. Subsequently, slides were counterstained with hematoxylin (Merck KGaA, Darmstadt, Germany) and LiCO₃ (Sigma Aldrich) and embedded with immumount mounting medium (Thermo Fisher Scientific Inc.).

3.6 Aggregometry

Aggregometry is a standardized photometric method which allows determination of platelet aggregation upon the addition of different agents. As shown in figure 3.1, light transmission through a cuvette in which platelet-rich-plasma (PRP) is stirred by a small magnetic stir bar can be displayed as proportional to platelet aggregation.

PRP was generated as described above (chapter 3.4). After generation, PRP was kept at 37°C in an incubator to maintain normal platelet function and avoid clustering of

Fig. 3.1: Measurement of platelet aggregation by aggregometry. Platelet aggregation can be displayed as proportional to light transmission through a cuvette containing platelet rich plasma (PRP) which is being stirred by a small magnetic stir bar.

platelet surface receptors. Platelets were incubated for a maximum of 30 min. Directly before use in the aggregometer, platelets were incubated for 10 min at room temperature with p0p/B or p0p1 Fab fragments at 50 $\mu\text{g}/\text{ml}$ each (see chapter 3.1 on page 17) or with the selective P2Y(12) inhibitor AR-C69931MX at 10 μM , which was a generous gift from ASTRA Charnwood (Loughborough Leics, England). Subconfluent B16F10 cells were detached and washed as described above (chapter 3.4) and resuspended in Tyrode's buffer (137 mmol/l NaCl, 2 mmol/l KCl, 12 mol/l NaHCO_3 , 0.3 mmol/l NaH_2PO_4 , 5.5 mmol/l glucose, 5 mmol/l HEPES, pH 7.3, 0.35% bovine serum albumin) at 1×10^3 cells/ μl . Aggregometry was performed in an ATRACT 4 S Analyzer (Rolf Greiner BioChemica GmbH, Flacht, Germany) with 50 μl of PRP and 110 μl Tyrode's buffer per cuvette (LABiTec GmbH, Ahrensburg, Germany).

After 30 seconds of stirring of the diluted PRP without any inductor of platelet aggregation, 50 μl of tumor cell suspension or ADP (adenosine 5'-diphosphate) at a final concentration of 10 μM were added to induce aggregation. Light transmission was monitored for 20 min. Aggregation curves were calculated and displayed using the ATRACT 4 software and Excel (Microsoft Office, Microsoft Deutschland GmbH, Unterschleißheim, Germany). Aggregation was calculated as percent of average aggregation induced by B16F10 alone.

3.7 Assessment of receptor occupancy

To determine the amount of anti-GPIb α Fab-fragments needed for optimum receptor blockade *in vivo*, C57BL/6 mice were injected with 5 μ g, 10 μ g, 20 μ g, 35 μ g, 50 μ g or 100 μ g of p0p/B (anti-GPIb α) Fab-fragments or with 100 μ g control rat IgG Fab-fragments ($n = 2$ mice for each concentration). Two hours later, mice were bled from the retroorbital plexus under ether anesthesia, as described before. 50 μ l of heparinized blood were incubated with 0.5 μ g of p0p/B-FITC for 10 min at room temperature. After incubation, blood was diluted up to 1 ml with PBS and fluorescence intensity was measured by flow cytometry. Receptor occupancy was computed as shown in the following equation:

$$RO = \left(1 - \frac{F(m)}{F(0)}\right) \times 100$$

RO: Receptor occupancy

F(m): Fluorescence intensity for a given amount m of unlabeled antibody p0p/B

3.8 Experimental pulmonary metastasis

Antibodies to deplete platelets or to block the platelet receptors of interest were applied to the mice as follows: For platelet depletion, mice were injected intraperitoneally with 100 μ g of a mixture of p0p3 and p0p4, two monoclonal rat-anti-mouse antibodies directed against the N-terminal 45-kD portion of GPIb α (chapter 3.1, page 17) in 200 μ l PBS. Control mice received the same amount (100 μ g), of monoclonal rat IgG in 200 μ l PBS. Intraperitoneal antibody application was performed 1 day prior to the injection of melanoma cells.

For GPIb α -inhibition, mice were injected with 50 μ g of p0p/B Fab-fragments into the retroorbital venous plexus. Control mice received the according amount of monoclonal rat IgG Fab fragments. In most experiments, GPIb α blockade was carried out 2 h before the injection of melanoma cells. In all experiments using p0p/B fab fragments to inhibit GPIb α , successful blockade of GPIb α was controlled directly after tumor cell injection: A droplet of mouse blood from the lateral tail vein was caught in a 500 μ l Eppendorf tube containing 300 μ l of 20 U/ml heparin. 650 μ l of PBS were added to the anticoagulated blood and 50 μ l of blood diluted in this way were then stained with p0p/B-FITC as described in chapter 3.7. The blood of mice that had received p0p/B Fab fragments was compared to that of mice which had received isotype control antibodies. For those experiments where GPIb α -blockade was carried out at different time points in relation to melanoma cell injection, 2.5×10^5 B16F10 in 150 μ l PBS were applied 2 h before, concomitantly with, or 2 h after intravenous tumor cell inoculation.

GPVI blockade was achieved by intraperitoneal injection of 200 μg of JAQ1 (anti-GPVI) divalent monoclonal antibody in 100 μl PBS five days prior to the injection of B16F10 melanoma cells. As control, mice were injected with the same amount of rat IgG antibody, also intraperitoneally. Receptor blockade was controlled by flow cytometry on the day of tumor cell injection as described previously for the control of GPIIb α -blockade, using monoclonal JAQ1-FITC.

For blockade of GPIIb/IIIa, mice were injected retroorbitally with 100 μg of JON/A (anti-GPIIb/IIIa) Fab-fragments, 2 h prior to tumor cell application. Control antibodies were 100 μg of monoclonal rat IgG Fab-fragments. For control of receptor blockade, a droplet of mouse blood was stained as described for the control of GPIIb α blockade, using monoclonal JON/A-FITC antibodies. Blockade in comparison to the control group that had received only rat IgG control Fab-fragments was assessed by flow cytometry.

Blockade of Mac-1 was achieved by intraperitoneal injection of 200 μg M1/70 monoclonal antibody 2 hours before tumor cell inoculation. As control, mice received 200 μg of rat IgG. Again, receptor blockade was controlled by flow cytometry on the day of tumor cell injection, using M1/70-FITC monoclonal antibody.

Experimental pulmonary metastasis was performed using B16F10 melanoma cells and C57BL/6 mice: Subconfluent B16F10 that had been cultured in a T75-flask were washed with PBS (Gibco, Invitrogen GmbH) and detached by a 0.25 % trypsin-, 0.02 EDTA solution (Sigma Aldrich).

The cells were washed twice by suspension in DMEM containing 10 % FCS and centrifugation at 1300 rpm for 3 min. The melanoma cells were then resuspended in PBS at 1.66×10^6 cells/ml and kept on ice until shortly before injection, when they were gently resuspended and warmed to body temperature. 150 μl of the cell suspension, containing 2.5×10^5 cells were injected per mouse through the lateral tail vein. After injection, the mice were returned to their cages. After 10 days, the mice were killed under ether anesthesia by cervical dislocation, the thorax was opened and the lungs were removed *in toto* and rinsed with PBS. After the lungs had been freed of adjacent tissue or remains of larger vessels, the weight of each lung was measured, a picture of each lung was taken using a Ricoh Caplio R8 Digital camera, and the numbers of surface metastases on each side of all lobes were counted using a loupe and a cell counter. After that, the lungs were dissected into single lung lobes. The right pulmonary lobes were conserved in paraformaldehyde (4% in PBS) for paraffin embedding and hematoxylin and eosin (H&E) staining, and the left pulmonary lobes were snap frozen in liquid nitrogen for immunohistochemical analyses (Schön et al. 2008).

3.9 Analysis of metastatic areas in lung sections

For histological analysis, 5 μm sections of paraffin-embedded lungs were prepared using a LEICA CM3050 S microtome (Leica Microsystems, Wetzlar, Germany) microtome. Hematoxylin and eosin (H&E) staining of paraffin-embedded lung sections was performed according to standard protocols. Microscopic pictures were taken with a Zeiss Axioskop 2 microscope (Jena, Germany) at a 40x magnification. Computer-based morphometric analysis of the area covered by tumor masses within the lung sections in comparison to the whole histological section was performed using the MetaMorph (Molecular Devices, Downing town, PA) and Image J (<http://rsbweb.nih.gov/ij/>) softwares.

3.10 Short term metastasis

B16F10 melanoma cells were stably transfected with green fluorescent protein (GFP) according to an optimized protocol for B16F10 (Amaxa Köln, Germany) using 1.5×10^6 cells, 2 μl pmaxFP-Green-C and 100 μl of Nucleofactor solution V (Amaxa): In short, cells were trypsinized, counted, centrifuged at 90 g and resuspended in Cell Line Nucleofactor Solution V at 1.5×10^6 cells / 100 μl for 15 min. 2 μg pmaxFP-Green-C were added, the suspension transferred to an amaxa-certified cuvette, and the appropriate Nucleofactor program, P-020, was run according to the manual. Afterwards, cells were transferred to 6-well plates and selected for gene expression starting 2 days after nucleotransfection by culturing in cell culture medium that contained 1 mg/ml selection supplement geneticin-148 (Sigma Aldrich).

Transfected melanoma cells could easily be distinguished from non-transfected cells by flow cytometry. To these means, a single cell suspension of GFP-transfected melanoma cells was prepared by trypsinization of subconfluent melanoma cells. Cells were suspended in FACS-buffer (1 g BSA, 0.1 g NaN_3 in 100 ml PBS), and fluorescence was assessed using the FACScalibur (BD Biosciences). GFP expression was measured in the FITC channel at 520 nm. To evaluate whether a discrimination between normal, non-transfected B16F10 melanoma cells and GFP-expressing transfectants was possible by flow cytometry, transfected and non-transfected cells were mixed in one single cell suspension, and their mean fluorescence intensity in the FITC-channel was evaluated. As it was very easy to differentiate between transfected and non-transfected cells, transfected cells with the highest expression of GFP were selected using a FACS Diva cell sorter (BD Biosciences). and subcloned by this method three times. The subclone with the highest expression of GFP after this selection process was used for all subsequent short-term metastasis assays.

For short-term metastasis experiments, GFP-labeled B16F10 melanoma cells were detached and washed as described in chapter 3.8. Concentration of melanoma cells was set to 2×10^6 cells in 150 μ l PBS. 6-week-old C57BL/6 mice which were sex-, and weight-matched either received 50 μ g p0p/B Fab-fragments or 50 μ g rat control Fab-fragments 2 h prior to tumor cell injection. 2×10^6 GFP-labeled melanoma cells were then injected under ether anesthesia into the retroorbital venous complex. Control mice had also been injected with either p0p/B Fab-fragments or isotype control antibodies, and received only 150 μ l of PBS instead of tumor cells. Mice were killed either 30 min, 2 h or 5 h after injection of the tumor cells. The control mice that had received only PBS were killed together with the first mouse that had received tumor cells. The lungs were excised and prepared as described in chapter 3.8. The lung lobes were separated from each other, and the right lung lobes were snap-frozen in liquid nitrogen for histological analysis. From the left lung lobes, a single cell suspension was prepared for flow cytometric analysis. Towards this end, the left lung lobes were chopped into small pieces and incubated in 1 mg/ml liberase solution, an enzyme blend containing collagenase isoforms as well as thermolysin (Roche Diagnostics GmbH) at 37°C for 90 min for enzymatic dissociation of the lung tissue complex. The tissue was then minced in Hank's Buffered Salt Solution (HBSS) containing 0.1 % bovine serum albumin (BSA) and passed through a 40 μ m cell strainer (BD Biosciences). The suspension was washed once in HBSS (0.1 % BSA) in a 15 μ l Falcon tube (BD Biosciences) and spun down at 1600 rpm for 5 min. For hypotonic lysis of erythrocytes, lung cells were resuspended in 2 ml of ice-cold distilled water while vortexing. 2 ml of 0.8 % NaCl solution were added immediately afterward. The Falcon tubes were filled up with HBSS (0.1 % BSA), passed through a cell strainer (Greiner Bio-One GmbH) and centrifuged for 5 min at 1600 rpm. The cell pellet was resuspended in 1 ml FACS buffer. 4 μ l of propidium iodide solution (BD Biosciences) were added to discern viable tumor cells. Viable GFP-positive B16F10 cells were quantitated by flow cytometry, GFP-positive cells were counted per 100,000 lung cells. In a complementary analysis, 5 μ m cryostat-cut sections of the right lung lobes were assessed by fluorescence microscopy (Axioskop, Zeiss). GFP-labeled B16F10 cells were counted in at least 13 microscopic fields per experimental condition. Sequential phase contrast and fluorescence microscopy allowed topographic analysis of tumor cells within the lung tissue.

3.11 Immunohistochemical staining against Mac-1

Immunohistochemical analysis of leukocyte recruitment into lung tissue after short-term metastasis was performed according to the protocol in chapter 3.5 19 by staining against Mac-1 in cryostat-cut lung sections from short-term metastasis experiments.

3.12 Apoptosis assay

To assess whether vWF had a direct apoptotic influence on B16F10 melanoma cells, an apoptosis assay was performed using the Roche Cell Death Detection ELISA (Roche Diagnostics GmbH). For preparation of the cells, 1×10^5 B16F10 were cultivated in a 6-well BD Falcon plate (BD Biosciences). After 10 h, cells were exposed to purified human plasma-derived vWF (American Diagnostica, Pfungstadt, Deutschland) either at 5 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$ or 20 $\mu\text{g/ml}$ in DMEM, high glucose, with 10% fetal calf serum (FCS), 2 mM L-glutamine, 100 $\mu\text{g/ml}$ streptomycin and 100 U/ml penicillin, with 10 $\mu\text{g/ml}$ vWF being the physiological concentration that is to be expected in mouse blood (Terraube et al. 2007). For negative control, cells were incubated with fresh medium without any supplementary factors. For positive control, cells were treated with staurosporin (Roche Diagnostics GmbH) at 37.5 ng/ml final concentration, also in DMEM containing 4.5 g/ml glucose. Cells were incubated for 16 h, and the apoptosis assay was performed the next day according to the manual. In short, cells were lysed and, by centrifugation at 10,000 $\times$ g for 10 min, a clear supernatant was prepared. Equal amounts of supernatant were added to microtiter plates that had been coated with anti-histone antibodies the previous day, and incubated for 90 min at room temperature. The anti-histone Cell Death Detection ELISA detected histone-bound DNA fragments. The substrate reaction, resulting in a green color, was evaluated by a plate reader at 405 nm. For each condition, three substrate reactions were measured and the average strength of the substrate reaction representing the magnitude of apoptosis is displayed in (%), compared to the value of untreated cells.

3.13 Chemicals and laboratory equipment

Table 3.1: *Chemicals*

Chemical	Company
Acetate buffer solution, pH 5.2	Sigma Aldrich, Taufkirchen, Germany
Acetone	Merck KGaA, Darmstadt, Germany
Adenosine 5'-Diphosphate (ADP)	Sigma Aldrich, Taufkirchen, Germany
Bovine serum albumin (BSA)	Roth, Karlsruhe, Germany
Diethyl ether	Merck KGaA, Darmstadt, Germany
Disodium hydrogen phosphate (Na_2HPO_4)	Roth, Karlsruhe, Germany
Dulbecco's PBS	PAA Laboratories GmbH, Cölbe, Germany
Geneticin	Gibco, Invitrogen GmbH Karlsruhe, Germany
Glucose	Roth, Karlsruhe, Germany
Glycine	Roth, Karlsruhe, Germany
Dulbecco's modified eagle medium (DMEM)	Gibco, Invitrogen GmbH Karlsruhe, Germany
Fetal calf serum (FCS)	Sigma Aldrich, Taufkirchen, Germany
Formaldehyde	Th. Geyer GmbH, Renningen, Germany
Glycine-hydrochloride	Sigma Aldrich, Taufkirchen, Germany
Hank's balanced salt solution (HBSS)	Sigma Aldrich, Taufkirchen, Germany
Hematoxylin solution	Merck KGaA, Darmstadt, Germany
Heparin (Liquemin, N25000)	Roche Diagnostics GmbH, Penzberg, Germany

Table 3.1: Chemicals

Chemical	Company
Hydrogen peroxide	Roth, Karlsruhe, Germany
Potassium chloride	Roth, Karlsruhe, Germany
Liberase-Solution	Roche Diagnostics GmbH, Penzberg, Germany
Lithium carbonate (LiCO_3)	Sigma Aldrich, Taufkirchen, Germany
Methanol	Merck KGaA, Darmstadt, Germany
Monosodium phosphate (NaH_2PO_4)	Roth, Karlsruhe, Germany
Normal goat serum	PAA Laboratories GmbH, Cölbe, Germany
Normal rabbit serum	PAA Laboratories GmbH, Cölbe, Germany
Paraformaldehyde	Roth, Karlsruhe, Germany
Penicillin	Gibco, Invitrogen GmbH Karlsruhe, Germany
Sodium azide (NaN_3)	Sigma Aldrich, Taufkirchen, Germany
Sodium carbonate (NaHCO_3)	Roth, Karlsruhe, Germany
Staurosporin	Sigma Aldrich, Taufkirchen, Germany
Tris(hydroxymethyl)aminomethane-buffer (Tris-buffer)	Sigma Aldrich, Taufkirchen, Germany
Trypsin-EDTA	Gibco, Invitrogen GmbH Karlsruhe, Germany

Table 3.2: *Laboratory equipment*

Laboratory equipment	Company
APACT 4 S Analyzer	Rolf Greiner BioChemica GmbH, Flacht, Germany
Centrifuges	Eppendorf, Hamburg, Germany
ElisaReader FluorStar Optima	BMG Labtechnology, Offenburg, Germany
FACS Calibur	BD Biosciences, Heidelberg, Germany
FACS Diva	BD Biosciences, Heidelberg, Germany
Harvard Pump 11 Plus Advanced Single Syringe with Dual RS-232	Harvard Apparatus, Holliston, USA
Microtome LEICA CM 3050 S	Zeiss, Jena, Germany
Zeiss Axioskop2	Leica Mikrosysteme Vertrieb GmbH, Wetzlar, Germany

4 Results

4.1 Antibody-mediated blockade of platelet GPIb α and its impact on TCIPA

In this section, several *in vitro* experiments such as the assessment of GPIb α expression on platelets, on melanoma cells or in normal lung tissue as well as aggregometry under GPIb α blockade were performed to provide a foundation for later *in vivo* metastasis experiments.

4.1.1 Assessment of GPIb α expression on platelets

Fig. 4.1: Assessment of GPIb α expression on platelets and B16F10 melanoma cells. 50 μ l of platelet rich plasma or 10^6 B16F10 melanoma cells in 50 μ l PBS were stained with 0.5 μ g of monovalent p0p/B (anti-GPIb α)-FITC and analyzed by flow cytometry. Mean fluorescence intensities of stained platelets and melanoma cells (unshaded histograms) are displayed in comparison with unstained platelets and unstained melanoma cells (grey shaded histograms).

To ascertain that only platelets but not B16F10 melanoma cells expressed GPIb α and

to thus exclude any direct interference of the antibodies we used with the B16F10 melanoma cells, a flow cytometric assay was performed. Reactivity of isolated murine platelets in platelet rich plasma (PRP) and of melanoma cells with p0p/B-FITC mAb (anti GPIb α) was analyzed, as shown in figure 4.1. While the shift of the mean fluorescence intensity in the flow cytometric assay indicated a strong expression of GPIb α on the platelet surface, B16F10 melanoma cells showed no reactivity with p0p/B-FITC.

4.1.2 Immunohistochemical analysis of GPIb α expression in lung tissue

Fig. 4.2: Immunohistochemical staining of GPIb α on lung sections. 5 μ m cryostat-cut lung sections of untreated C57BL/6 mice were stained by the ABC-immunoperoxidase method, using p0p/B- and p0p1 HRP-conjugated monoclonal antibodies directed against GPIb α . Intravascular platelets show reactivity, whereas lung tissue is negative. The scale bar corresponds to 10 μ m.

To exclude an influence of the GPIb α -blockade on endothelial cells and other cells present in the lung tissue and to extend a previous report dealing with this question (Bergmeier et al. 2000), an immunohistochemical staining against GPIb α was performed on lung sections of untreated wildtype mice using two different HRP - conjugated monoclonal antibodies (p0p/B and p0p1). While platelets localized within the blood vasculature clearly showed reactivity with the employed antibodies, no other cells within the lung sections could be visualized in this manner (see figure 4.2).

4.1.3 Tumor cell-induced platelet aggregation (TCIPA) under GPIb α blockade

As mentioned in the introduction, the ability of tumor cells to induce platelet aggregation seems to be of crucial importance to their survival within the blood stream and to the ensuing establishment of metastatic foci (chapter 2.2). Therefore, it appeared necessary to test whether the B16F10 melanoma cells used could also induce this aggregation and whether blockade of GPIb α would interfere with this process. Therefore platelet aggregation was tested after B16F10 melanoma cell addition using aggregometry, a photometrical method to quantify the formation of platelet aggregates after the addition of B16F10 melanoma cells in a standardized manner. Platelet rich plasma was either incubated with monoclonal monovalent p0p/B antibody or monoclonal monovalent p0p1 antibody, as this antibody also specifically binds GPIb α but is not functionally inhibitory (Bergmeier et al. 2000). The specific inhibitor of the ADP receptor P2Y(12), AR-C69931MX, was used as positive control for the inhibition of TCIPA, as this process is known to depend strongly upon ADP (Alonso-Escolano et al. 2004). As expected, B16F10 melanoma cells induced a profound platelet aggregation (see figure 4.1).

While AR-C69931MX reduced this aggregation to 13 % (± 7.73 %) as compared to TCIPA using B16F10 cells only as inductor ($n = 4$ independent experiments in the B16F10 alone group and $n = 3$ experiments for inhibition with AR-C69931MX, $P = 0.00017$), neither p0p/B nor p0p1 led to any alteration of the aggregation induced by the melanoma cells. After incubation with p0p/B Fab fragments, B16F10 melanoma cells still induced 96 % of the aggregation achieved by melanoma cells alone (± 13.07 %; $n = 4$ experiments, $P = 0.66$ compared to the untreated controls). Incubation of platelets with p0p1 Fab fragments led to an aggregation of 107 % as compared to the untreated controls (± 5.8 %; $n = 3$ experiments, $P = 0.70$). In conclusion, this functional assay does not yield any indication of a positive or negative influence of monoclonal monovalent p0p/B antibody on tumor cell-induced platelet aggregation.

4.1.4 Assessment of receptor occupancy

In order to titrate the minimal amount of monoclonal monovalent p0p/B antibody necessary to achieve saturating receptor binding *in vivo* and thus functional blockade of GPIb α , a flow cytometric analysis was employed to evaluate the necessary antibody concentrations.

While mice had received either 10 μg , 20 μg , 35 μg or 50 μg retroorbitally, it could be shown that a concentration of 20 μg was enough to saturate GPIb α receptors,

Fig. 4.3: TCIPA under GPIb α blockade. Platelet rich plasma was incubated with p0p/B (anti-GPIb α) Fab fragments or p0p1 Fab fragments at 50 μ g/ml each or with the selective P2Y(12) inhibitor AR-C69931MX at 10 μ M. After 30 seconds, 5×10^4 B16F10 melanoma cells were added to the cuvette, and aggregation of platelets was monitored for 20 minutes. The figure shows characteristic graphs for each condition, as indicated.

Fig. 4.4: Determination of optimum antibody concentrations for GPIIb/IIIa binding. To determine optimum antibody concentrations for nearly complete receptor blockade, C57BL/6 mice were injected with the indicated amounts of monovalent p0p/B (anti-GPIIb/IIIa) Fab fragments. $n = 2$ per antibody concentration. 50 μl of full blood per mouse were then drawn from the retro-orbital plexus and diluted with 300 μl heparin in PBS. Diluted blood was then incubated with 0.5 μg of p0p/B-FITC. GPIIb/IIIa blockade was subsequently assessed by flow cytometry. The graph shows the average receptor blockade in percent, $\pm$ SD.

leading to a receptor occupancy of 95 %. Therefore, in all consecutive experiments that required *in vivo* GPIIb/IIIa blockade, 50 μg of p0p/B Fab fragments were used to ensure near complete receptor blockade.

4.2 GPIIb/IIIa and pulmonary metastasis

The next series of experiments was designed to determine whether specific inhibition of GPIIb/IIIa, without induction of thrombocytopenia, could alter metastasis formation in a murine *in vivo* model of pulmonary metastasis. Subsequent experiments were aimed at unraveling the mechanisms underlying the influence GPIIb/IIIa showed on the metastatic potential of melanoma cells.

4.2.1 Experimental pulmonary metastasis after platelet depletion

Even though the dependency of the metastatic process upon platelets had been shown previously by many groups, it was necessary to verify this relationship in the experimental system used here, that is, using B16F10 melanoma cells and C57BL/6 mice. Platelets were therefore depleted using a mixture of two monoclonal divalent antibodies,

p0p3/p0p4, which selectively bind the N-terminal 45 kD-portion of the GPIIb α receptor and thus induce rapid and irreversible aggregate formation and subsequent platelet depletion (Bergmeier et al. 2000). Ten days after injection of B16F10 melanoma cells, the group of mice that had received the platelet-depleting antibody mix showed a significantly reduced number of metastatic foci as compared to the IgG control group (see figure 4.5).

Fig. 4.5: Experimental pulmonary metastasis after platelet depletion. Experimental pulmonary metastasis was performed in C57BL/6 wt mice by i.v. injection of 2.5×10^5 B16F10 melanoma cells. Lungs were extracted 10 days later and metastatic foci counted. Platelets had been depleted one day previously by intraperitoneal injection of 100 μ g of the divalent anti-GPIIb α antibody mix p0p3/4. Control mice had received divalent rat IgG ($n = 7$ per group). $\pm$ SD. P values are indicated.

While the mice from the control group presented an average number of 319.1 ± 182.2 metastases per lung surface, in the group where platelets had been depleted this number was reduced to 17.9 ± 9.3 metastatic foci ($n = 7$ mice in both groups, $P < 0.001$).

4.2.2 Pulmonary metastasis under GPIIb α blockade

The following experiment was directed at assessing the role of GPIIb α in pulmonary metastasis. As a positive control to demonstrate that metastasis can be altered by blockade of a platelet surface receptor, a group of mice was treated with 100 μ g monoclonal JON/A Fab-fragments ($n = 8$ mice), directed against the integrin receptor

Fig. 4.6: Experimental pulmonary metastasis under blockade of GPIIb/IIIa. Experimental pulmonary metastasis was performed in C57BL/6 mice by i.v. injection of 2.5×10^5 B16F10 melanoma cells, 2 hours after i.v. injection of either 100 μ g of monovalent rat IgG Fab fragments, 100 μ g of monovalent JON/A (anti-GPIIb/IIIa) or 50 μ g p0p/B (anti-GPIIb), as indicated. $n = 7$ mice in the p0p/B group, $n = 8$ mice in both the control group and the JON/A group. Lungs were removed 10 days later and surface metastases were counted. Three representative pictures from each group are shown. The graph shows the average number of metastases per group, $\pm$ SD. P values are indicated.

GPIIb/IIIa (α IIB β 3 integrin) which has been shown to play an important role in metastatic dissemination of different tumor cell lines (Karpatkin et al. 1988), although its effect on metastasis of B16F10 has not been shown so far. A second group of control mice ($n = 8$) received 100 μ g of rat IgG isotype control. Finally, GPIb α was blocked in a third group ($n = 7$) by retroorbital injection of 50 μ g of monoclonal monovalent p0p/B (anti-GPIb α). As expected, blockade of GPIIb/IIIa led to a significant reduction of the number of metastatic foci in the lung tissue as could be observed 10 days after injection of B16F10 melanoma cells. While in the isotype control group the average number of metastases per lung surface was 359.4 ± 200.7 , GPIIb/IIIa-blockade led to a significantly attenuated number of metastases of 181.5 ± 75.4 per lung ($P = 0.045$) (see figure 4.6). Surprisingly, blockade of GPIb α did not lead to a decline in metastasis but instead to a strong increase of metastatic foci within the lungs in comparison to the control mice. The average number of metastases in the group that had received p0p/B Fab-fragments was 899.4 ± 358.2 compared to the isotype control ($P = 0.0047$). This metastasis-promoting effect of GPIb α inhibition was verified by two consecutive, independent experiments and showed similar results (not shown).

Fig. 4.7: Histological analysis of lung sections after experimental pulmonary metastasis under $GPIIb\alpha$ blockade. As described in figure 4.6, experimental pulmonary metastasis was performed in C57BL/6 mice by i.v. injection of 2.5×10^5 B16F10 melanoma cells after blockade of $GPIIb/IIIa$ or $GPIIb\alpha$, as indicated. Lungs were removed after 10 days, and $5 \mu\text{m}$ thick lung sections were stained with hematoxylin-eosin. The percentage of the lung section covered with tumor cells was evaluated morphometrically (figure 4.8). Representative lung sections from all three groups are shown. Arrows point to tumor masses. Scale bar = $500 \mu\text{m}$.

Fig. 4.8: Morphometric analysis of lung sections after experimental pulmonary metastasis under GPIb α blockade. As shown in figure 4.7, experimental pulmonary metastasis was performed in C57BL/6 mice by i.v. injection of 2.5×10^5 B16F10 melanoma cells after blockade of GPIIb/IIIa or GPIb α , as indicated. Lungs were removed after 10 days, 5 μ m thick lung sections were stained with hemotoxylin-eosin, and morphometric evaluation of the percentage of the lung section covered with tumor cells was performed. The graph shows the average metastatic area in lung sections from all three groups, $\pm$ SD. P values are indicated, $n = 7$ mice in the p0p/B group, $n = 8$ mice in both the control group and the JON/A group.

Finally, the strong differences between mice of the control group, the JON/A group and the GPIb α group could also be visualized in paraffin sections of the lung tissue after a hematoxylin-eosin-staining had been performed. Tumor cell aggregates, visible as strongly basophilic conglomerates of cells, are not only more numerous in the group that received p0p/B Fab fragments, but also larger in size (figure 4.7). Objective computer-based morphometric evaluation of lung sections from the control group, the group that had received JON/A Fab (anti-GPIIb/IIIa) fragments or p0p/B (anti-GPIb α) revealed a significant difference in tumor masses in relation to the total histological area per section in the three experimental groups. In the control group, the average sectional

area covered by tumor cells was $7.9 \pm 2.1\%$ of the total lung section. In the JON/A group, the average value was $2.0 \pm 1.2\%$ ($P < 0.001$). In strong contrast to this, the average metastatic area in the p0p/B (anti-GPIb α) group was $17.3 \pm 6.3\%$ ($P < 0.001$ compared with that of the control group, $n = 7$ mice in the p0p/B group, $n = 8$ mice in both the control group and the JON/A group).

4.2.3 Critical time frame for GPIb α impact on metastasis

Fig. 4.9: Time frame for the influence of GPIb α blockade in pulmonary metastasis. Experimental pulmonary metastasis was performed in C57BL/6 mice by the i.v. injection of 2.5×10^5 B16F10 melanoma cells. GPIb α was blocked with 50 μ g monoclonal monovalent p0p/B antibody at different time points, either 2 hours prior to, together with or 2 hours after tumor cell inoculation. Lungs were removed after 10 days, and metastatic foci on the lung surfaces were quantitated. Three representative pictures from each group are shown. The graph depicts the average number of metastases per group, $\pm$ SD. P values are indicated, $n = 4$ mice in all groups.

Platelet receptors are likely to influence the initial steps of metastasis, that is, the survival of the tumor cells within the blood stream as well as adhesion to endothelial cells, rather than tumor cell growth or local progression (chapter 2.2). Therefore, it seemed reasonable to assume that GPIIb/IIIa-blockade would only play a role in our experimental system of pulmonary metastasis within a very narrow time frame in relation to the injection of the tumor cells. To test this hypothesis, C57BL/6 mice were injected with 50 μ g monoclonal, monovalent p0p/B antibody at different time points, either 2 hours prior to, together with or 2 hours after tumor cell inoculation. Evaluation of pulmonary metastases 10 days later revealed that blockade of GPIIb/IIIa was only effective before or together with tumor cell injection. GPIIb/IIIa-blockade 2 hours prior to tumor cell injection resulted in a significant increase of metastases compared to the rat IgG control group from 66.3 ± 11.9 metastases/lung to 166.0 ± 62.0 metastases/lung ($P = 0.044$). Simultaneous GPIIb/IIIa blockade and tumor cell inoculation also led to an increase from 54.5 ± 33.2 metastases/lung in the control group that received rat IgG Fab fragments together with the tumor cells to 194.0 ± 61.7 metastases/lung ($n = 4$ mice in all groups, $P = 0.036$). In contrast to these results, later inhibition of GPIIb/IIIa did not result in an elevated number of metastases. With 55.4 ± 23.66 metastases/lung, mice that had received p0p/B Fab-fragments 2 hours after tumor cell inoculation showed an average number of metastases very similar to that of the group which had received rat IgG 2 hours after injection, which was 63.6 ± 26.2 metastases/lung ($n = 5$ mice in both groups, $P = 0.86$). So while GPIIb/IIIa has a pronounced influence on certain phases of the metastatic process, this influence seems to be limited to early steps in the metastatic cascade.

4.2.4 Short term metastasis

To further elucidate the early effects of GPIIb/IIIa blockade, B16F10 were stably transfected with GFP (green fluorescent protein) to allow the detection of us single cells within the lung tissue directly after tumor cell inoculation.

The transfected melanoma cells could be easily distinguished from non-transfected cells by flow cytometry (figure 4.10 allowing selection of the subclone with the highest GFP expression. GFP expression could be evaluated by flow cytometry by measuring the mean fluorescence intensity in the FITC-channel at 519 nm and there was hardly any overlap between the non-transfected cells and the GFP positive cells.

Pulmonary metastasis was assessed using the cell clone with the highest level of GFP-expression so as to allow easy detection of melanoma cells by flow cytometry and by fluorescence microscopy in lung sections.

Again, mice received 50 μ g of p0p/B monoclonal, monovalent antibody or of monovalent rat IgG isotype control. Tumor cell inoculation was performed by retroorbital

Fig. 4.10: *Flow cytometric evaluation of GFP expression after stable transfection of melanoma cells.* A single cell suspension of GFP-positive tumor cells together with non-transfected melanoma cells was prepared and the mean fluorescence intensity in the FITC-channel determined to distinguish GFP-positive from GFP-negative cells.

injection of 2×10^6 GFP-positive melanoma cells. Mice were killed either 30 min, 2 hours or 5 hours after tumor cell inoculation or without any previous injection of tumor cells to allow control for autofluorescent structures ($n = 3$ mice per condition). The respective number of tumor cells at each time point was evaluated by two complementary methods: flow cytometric assessment of GFP-positive tumor cells within the lung tissue (see figure 4.11) or microscopic quantification of tumor cells in cryostat-cut lung sections (shown in figure 4.12).

Fig. 4.11: Flow cytometric evaluation of short term-metastasis under blockade of GPIIbα. For short term metastasis, 2×10^6 B16F10 melanoma cells stably transfected with green fluorescent protein (GFP) were injected into the retroorbital plexus of C57BL/6 mice that had received either 50 μ g of monovalent p0p/B (anti-GPIIbα) monoclonal antibody or 50 μ g of monovalent rat IgG 2 hours earlier. Mice were killed at different time points (30 minutes, 2 hours, or 5 hours after melanoma cell injection). A single cell suspension was prepared from the left lung lobes and the number of GFP-positive tumor cells in relation to the total number of lung cells was determined by flow cytometry. Three individual experiments are shown under #1 to #3 as baseline variations made it difficult to compare the three experiments.

Fig. 4.12: Short term metastasis under GPIIb/IIIa blockade. For short term metastasis, 2×10^6 B16F10 melanoma cells stably transfected with green fluorescent protein (GFP) were injected into the retroorbital plexus of C57BL/6 mice that had received either 50 μ g of monovalent p0p/B (anti-GPIIb/IIIa) monoclonal antibody or 50 μ g of monovalent rat IgG 2 hours earlier. Mice were killed at different time points (30 minutes, 2 hours, or 5 hours after melanoma cell injection). Representative cryostat-cut lung sections (merged fluorescence and phase contrast images) from the control group and the group that had received p0p/B Fab fragments are depicted, as indicated. Scale bar = 20 μ m. The graph displays the average number of GFP-labeled tumor cells per microscopic field, $\pm$ SD, $n = 3$ per group, P-values are indicated.

In accordance with previous findings, microscopic assessment of GFP-labeled tumor cells revealed a significantly elevated number of tumor cells very early after tumor cell injection in the group where GPIIb α had been blocked previously as compared to the mice that had received only isotype control antibodies. As early as 30 minutes after tumor cell inoculation, mice that had received p0p/B Fab fragments showed 18.7 ± 0.2 tumor cells per microscopic field (10 x magnification) while the control mice merely presented 14.9 ± 1.2 ($P = 0.007$). At 2 hours after tumor cell inoculation, this difference was still very pronounced, as in the lung-samples of control mice an average of 10.6 ± 1.9 tumor cells per microscopic field was determined, while the respective value in the p0p/B-group was 14.0 ± 0.5 ($P < 0.045$). Finally, 5 hours after tumor cell transplantation, analysis of cryostat-cut-lung sections of control mice yielded 7.1 ± 2.8 GFP-labeled melanoma cells per microscopic field in contrast to an average of 10.1 ± 3.5 melanoma cells per microscopic field under GPIIb α -blockade ($P = 0.3$). This difference is not statistically significant anymore which seems to be mainly do to the fact that baseline values between the three experiments varied considerably. When comparing the numbers of GFP-positive cells present after 5h in the lungs of control animals and animals that had received p0p/B antibody in the individual experiments, the pro-metastatic effect of GPIIb α inhibition becomes visible: In the first experiment, 5 hours after tumor cell injection the mouse that had received anti-GPIIb α antibody showed an average of 6.9 tumor cells per microscopic field vs. 4.2 cells in the lung of the mouse that had received control antibody. In the second experiment, p0p/B-antibody injection produced a tumor cell number per microscopic field of 9.5 vs 7.1 in the control mouse. Finally, in the last experiment, the mouse that had received anti-GPIIb α had an average of 13.9 GFP-labeled melanoma cells in its lung, as compared to 9.9 in the control mouse.

Flow cytometric evaluation of the number of GFP-positive tumor cells in relation to the total number of cells within the lung tissue generally showed similar results. However, this method proved to be technically challenging, as treatment of the mice before tumor cell inoculation, preparation of single-cell suspensions as well as flow cytometric assessment all had to be performed within a very tight time frame on one single day, not allowing the analysis of a great number of mice in one experiment. Therefore, one mouse per day and per condition was analyzed, and experiments were carried out three times. Unfortunately, day-to-day variations regarding the baseline of GFP-positive cells in the lung tissue in this specific approach made it difficult to compare values obtained on different days. Therefore, the three conducted experiments are displayed individually in figure 4.11, all three of them consistently showing an increase in the number of B16F10 melanoma cells after GPIIb α blockade, and thus confirming the results of the fluorescence microscopy approach.

4.2.5 Mac-1-expressing immune cells after short-term metastasis

In order to unravel the mechanisms underlying the observed effects upon blockade of GPIb α , important binding factors of GPIb α were focused on. The integrin Mac-1 ($\alpha_M\beta_2$, CD11b/CD18, CR3) belongs to the family of β_2 integrin receptors and is expressed on leukocytes, particularly NK cells, monocytes and polymorphonuclear neutrophils. It is not only an important complement receptor recognizing the iC3b fragment (Todd 1996), essential for neutrophil function (Li 1999) and, thus, implicated in the recruitment of neutrophils in inflammatory processes (Asaduzzaman et al. 2009). It has also been shown to interact with the GPIb-V-IX complex and thus mediate platelet-leukocyte interactions (Ehlers et al. 2003), a process important for the biological response to vascular injury (Wang et al. 2005).

Therefore, it was conceivable that Mac-1 mediated a certain anti-metastatic protection through its interactions with GPIb α on activated platelets surrounding the tumor cells. Recognition of platelet-tumor cell emboli and subsequent elimination by leukocytes seemed a possible explanation for the observed effects after GPIb α -blockade. It was hypothesized that an abrogation of these interactions would lead to an inhibition of immune surveillance and, thus, to a promotion of metastasis. Presumably, under GPIb α blockade, reduced recruitment of Mac-1 positive cells in the lung tissue after tumor cell injection would be an indicator of an altered immune response. Therefore, the lungs from the short-term metastasis experiments were stained for Mac-1 positive cells by immunohistochemistry, using the M1/70 antibody. In both groups, the one that had received p0p/B Fab-fragments as well as in the isotype control group, injection of melanoma cells produced a strong inflammatory reaction, compared to the mice that had received only antibody and no tumor cells. Immunohistochemical staining of Mac-1 positive cells did not yield any visible differences between the group where GPIb α had been blocked compared to the isotype control group, as shown in figure 4.13. Thus, blockade of GPIb α did not affect the tumor cell-induced pulmonary recruitment of Mac-1 expressing cells of the innate immune system.

4.2.6 Double blockade of GPIb α with Mac-1

Therefore, another experimental approach to evaluate the importance of Mac-1-GPIb α -interaction was used. For this approach, GPIb α and Mac-1 were blocked by monoclonal antibodies in experimental pulmonary metastasis. Mice received either monoclonal, monovalent p0p/B antibody (anti-GPIb α), monoclonal, divalent M1/70, both p0p/B and M1/70 or rat IgG as isotype control.

Fig. 4.13: Pulmonary expression of Mac-1 after short-term metastasis. 5 μ m cryostat-cut sections from short-term metastasis experiments were stained by the ABC-immunoperoxidase method using the M1/70 monoclonal antibody against Mac-1 (CD11b). Mice had received 50 μ g of monovalent monoclonal p0p/B (anti-GPIIb α) antibody or 50 μ g of monovalent rat IgG 2 hours before inoculation of 2×10^6 B16F10 melanoma cells and were killed either 30 minutes, 3 hours or 5 hours after tumor cell inoculation. Characteristic pictures of lung sections of all three time-points under blockade of GPIIb α or of the control group are depicted, as indicated. Scale bar = 100 μ m.

Fig. 4.14: Experimental pulmonary metastasis under double blockade of GPIIb/IIIa with Mac-1. For experimental pulmonary metastasis, C57BL/6 mice were injected with 2.5×10^5 B16F10 melanoma cells. For functional blockade of GPIIb/IIIa, mice had received 50 μg monoclonal, monovalent p0p/B antibody 2 hours prior tumor cell inoculation. For Mac-1 blockade, intraperitoneal injection of 200 μg of M1/70 monoclonal, divalent antibody was performed 2 hours before tumor cell inoculation. For double blockade of GPIIb/IIIa and Mac-1, mice received p0p/B as well as M1/70. For control, rat IgG monoclonal antibody was used, as indicated. Lungs were removed after 10 days, and metastatic foci on the lung surfaces were quantified (figure 4.15). Three representative pictures from each group are shown.

Fig. 4.15: Quantitative evaluation of experimental pulmonary metastasis under double blockade of GPIb α with Mac-1. As described in figure 4.14, experimental pulmonary metastasis with B16F10 melanoma cells was performed in C57BL/6 mice under functional blockade of GPIb α , Mac-1 or, double blockade of both GPIb α and Mac-1. Lungs were removed after 10 days, and metastatic foci on the lung surfaces were quantified. The graph depicts the average number of metastases per group, $\pm$ SD. P values are indicated, $n = 8$ mice in the p0p/B group, the control group, and the double blockade group, $n = 7$ mice in the M1/70 group.

As shown in figures 4.14 and 4.15, mice lacking functionally active GPIb α after p0p/B-Fab injection showed a significant increase in metastasis. An average of 408.0 ± 139.7 metastases/lung surface were counted compared to the rat IgG control group with 118.0 ± 43.11 metastases/lung surface ($n = 8$ mice in both groups, $P = 0.00023$). The group in which Mac-1 had been functionally blocked, showed a similar increase of metastases up to 439.29 ± 256.69 metastases/lung surface ($n = 7$ mice, $P = 0.0023$ compared to the control group). Double blockade, however, led to an even more impressive increase in the number of metastatic colonies. In this final group, 914.63 ± 446.35 metastases/lung surface could be counted ($n = 8$ mice, $P = 0.00019$ compared to the control group). These observations were confirmed in a consecutive, independent experiment yielding very similar results (not shown).

4.2.7 Blockade of GPIb α and P-selectin in experimental melanoma metastasis

As the results obtained from of Mac-1-blockade alone or together with GPIb α inhibition showed the importance of Mac-1 during metastasis but did not yield a satisfactory explanation for the effects seen after GPIb α blockade, attention was put on another important binding partner of GPIb α : P-selectin. P-selectin is a C-type selectin which recognizes sialylated, fucosylated carbohydrates, specifically glycans containing a terminal Sialyl Lewis^x (sLe^x), in a calcium-dependent fashion. The molecule is expressed on activated endothelial cells as well as on platelets with its major counterreceptor, PSGL-1, being present mainly on circulating leukocytes. In addition, P-selectin is an important counterreceptor for GPIb α as well (Romo et al. 1999). It is known to play a role in hematogenous metastasis, as inhibition of P-selectin by heparin severely diminishes metastasis (Borsig 2007). The importance of P-selectin in the process of hematogenous tumor dissemination can also be seen in genetically engineered mice lacking P-selectin (Ludwig et al. 2004).

To test whether there was a functional link between the observed enhancement of metastasis upon GPIb α blockade and the interaction of the GPIb-V-IX complex with P-selectin, GPIb α was inhibited using 50 μ g of p0p/b Fab fragments 2 hours prior to tumor cell inoculation in P-selectin deficient C57BL/6 mice and, as control, in wild-type C57BL/6 mice. Corresponding groups for each genotype received only 50 μ g rat IgG Fab fragments. As expected, P-selectin deficient mice showed a markedly reduced metastatic dissemination compared with the wild-type group. While the wild-type group showed an average of 80.2 ± 61.66 metastases/lung surface after treatment with rat IgG Fab fragments, P-selectin null mice only presented an average of 43.0 ± 14.9 metastases, as is shown in figure 4.16.

Fig. 4.16: Experimental pulmonary metastasis under GPIIb α blockade in P-selectin deficient mice. Experimental pulmonary metastasis was assessed in C57BL/6 wt mice or P-selectin deficient mice on a C57BL/6 background, as indicated, after i.v. injection of 2.5×10^5 B16F10 melanoma cells, 2 hours before injection 50 μ g p0p/B (anti-GPIIb α) or rat IgG control. Lungs were removed 10 days later and surface metastases counted. $n = 5$ in the wt groups, $n = 4$ in the P-selectin knockout groups. The graph shows the average number of metastases per group, $\pm$ SD. P values are indicated.

Upon blockade of GPIIb/IIIa, the number of metastases increased to 260.8 ± 159.8 metastases/lung in the wild-type group ($n = 5$ mice in both groups, $P = 0.046$). Interestingly, GPIIb/IIIa blockade in P-selectin deficient mice did not lead to an enhanced metastasis, as mice in this group only had 38.5 ± 5.8 metastases/lung surface ($n = 4$ mice in both groups, $P = 0.57$). Repetition of this experiment led to similar results (data not shown). Thus, the metastasis-promoting effect of GPIIb/IIIa blockade was completely abrogated in P-selectin deficient mice.

4.2.8 Impact of vWF on melanoma survival *in vitro*

Fig. 4.17: Influence of vWF on apoptosis of B16F10 melanoma cells. 1×10^5 B16F10 melanoma cells were incubated with the indicated amounts of vWF for 16 hours. Apoptosis was measured using an anti-histone ELISA. Substrate reaction is displayed as the percentage of substrate reaction of the untreated cells that were used as control. Treatment with staurosporin served as positive control.

Given that another research group had recently discussed a possible pro-apoptotic effect of vWF on melanoma cells (Terraube et al. 2007), the hypothesis was formulated

that under normal conditions vWF might accumulate around tumor cells through interaction with GPIb α on platelets that had previously been activated by the tumor cells. This regional accumulation would then lead to an apoptotic response of the metastasizing cells, providing a natural anti-tumoral defense mechanism. Blockade of GPIb α by monovalent antibodies would be expected to impede this mechanism, resulting in an increase of metastases in this model of experimental pulmonary metastasis. To test this hypothesis, it was necessary to confirm the pro-apoptotic effect of vWF on tumor cells that had been described as a very preliminary result by the other group. Melanoma cells were incubated with vWF and performed an anti-histone ELISA the following day. However, no apoptotic activity could be detected, not even when melanoma cells had been incubated with twice the physiological concentration of vWF, namely 20 $\mu\text{g}/\text{ml}$. The value of the photometrically measured substrate reaction of untreated cells was used as a reference value and set to 100%. Staurosporin was used as a positive control for the induction of apoptosis, and compared to untreated cells it induced a substrate reaction of $442 \% \pm 152,91 \%$ ($P = 0.019$). In contrast to this, the average substrate reaction of the cells treated with 5 $\mu\text{g}/\text{ml}$ vWF was $102.0 \% \pm 3.7 \%$. The cells treated with 10 $\mu\text{g}/\text{ml}$ showed a reaction of $99 \% \pm 13.5 \%$ and, finally, under 20 $\mu\text{g}/\text{ml}$ vWF a substrate reaction of $94 \% \pm 7.1 \%$ compared to the untreated cells could be measured ($P > 0.5$ in all three groups). Therefore, as it seemed unlikely that vWF could provide any kind of protection against melanoma metastasis, this hypothesis was not pursued any further.

4.3 Role of GPVI in metastasis

4.3.1 Experimental pulmonary metastasis under GPVI blockade

After having seen rather astonishing results when performing experimental pulmonary metastasis assays under GPIb α blockade, inhibiting of another important platelet adhesion receptor, the collagen receptor GPVI, was the next aim. 200 μg of JAQ1 (anti-GPVI) divalent monoclonal antibody in 100 μl PBS were used five days prior to the injection of B16F10 melanoma cells to block GPVI. Control mice received the corresponding amounts of divalent rat IgG. Treatment with JAQ1 has previously been shown to induce reversible thrombocytopenia for up to 72 hours. This effect is presumably not Fc dependent, as Fab fragments of this antibody led to the same result. After approximately 72 hours, platelet counts return to normal numbers and after 5 days, that is, on the day of the injection of melanoma cells, no thrombocytopenia can be expected any more (Nieswandt et al. 2001b; Schulte et al. 2006). A depletion in the sense of an irreversible removal of the GPVI-receptor as well as an abolishment of platelet response to collagen and CRPs (known to be a strong GPVI-specific platelet

agonist) of at least two weeks has been shown to be achieved by treatment with JAQ1 (Nieswandt et al. 2001b).

Fig. 4.18: Experimental pulmonary metastasis under blockade of GPVI. Experimental pulmonary metastasis was induced in C57BL/6 wt mice by i.v. injection of 2.5×10^5 B16F10 melanoma cells. Lungs were extracted 10 days later and metastatic foci were counted. 5 days before tumor cell injection, mice had received 200 μ g of JAQ1 (anti-GPVI) divalent monoclonal antibody intraperitoneally to block GPVI. Control mice received the corresponding amount of rat IgG monoclonal antibody. Representative pictures of lungs are depicted, the graph shows the average number of metastases per group, $\pm$ SD. P values are indicated, $n = 8$ mice each group.

In this assay of experimental pulmonary metastasis, mice again received 2.5×10^5 melanoma cells and lungs were harvested 10 days later. Unlike the results found in case of GPIIb/IIIa blockade, GPVI blockade did not influence melanoma metastasis in any visible way in this assay, as can be seen in figure 4.18. While the control mice showed an average of 359.4 ± 214.5 metastases/lung surface, the mice that had received JAQ1 antibody presented an average of 410.7 ± 162.5 metastases/lung ($n = 8$ mice in both groups, $P = 0.598$). Similar results were obtained when the assay was repeated (data not shown).

Therefore, no further experiments to test the *in vivo* influence of GPVI on pulmonary melanoma metastasis were conducted.

5 Discussion

5.1 GPIb α is expressed only on platelets and does not alter tumor cell-induced platelet aggregation (TCIPA)

The first part of this work characterized the expression of GPIb α on platelets, tumor cells, lung endothelium and other cells within the lung tissue and evaluated the role that GPIb α played in tumor cell-induced platelet aggregation. As the aim of this work was to investigate the influence of platelet GPIb α on tumor metastasis, it was necessary to assure that the interaction of the p0p/B monoclonal antibodies used was limited to platelets. It had to be excluded that the antibody influenced any cells that could be relevant in the *in vivo* system such as the melanoma cells themselves or cells within the lung tissue. In line with previous observations of other research groups (Bergmeier et al. 2000), the presence of GPIb α could be detected neither on melanoma cells nor on lung epithelial cells when using either flow cytometric evaluation of GPIb α expression on melanoma cells or immunohistochemical staining of lung tissue with two different monoclonal antibodies directed against GPIb α . Therefore, it is unlikely that inhibition of GPIb α with monoclonal, monovalent antibodies would, apart from platelets, directly influence relevant cells such as the melanoma cells, lung endothelium or lung stroma cells.

The role of GPIb α blockade in tumor cell-induced platelet aggregation was evaluated next, as this *in vitro* phenomenon appears to be, to a certain extent, an indicator of the *in vivo* metastatic potential of tumor cells (Karparkin et al. 1988). It therefore seemed important to test whether B16F10 melanoma cells could induce platelet aggregation and to elucidate whether p0p/B had any influence on this aggregation. An increase of TCIPA or a diminished aggregation under GPIb α blockade would have been a predictor of a respective promotion or an inhibition of metastasis *in vivo*. *In vitro* assays to determine an influence of GPIb α on TCIPA have hitherto been very contradictory, strongly depending on the experimental setting and the tumor cell line used. While some groups could not detect any diminished aggregation upon GPIb α blockade (Karparkin et al. 1988), early experiments performed by other research groups indicate a role of GPIb α

in at least some phases of TCIPA (Bastida et al. 1987; Grossi et al. 1987). The experimental setting, using B16F10 melanoma cells, the blood of C57BL6 mice and p0p/B monoclonal, monovalent antibodies, corroborated a strong induction of platelet aggregation by the B16F10 melanoma cells, but did not bring forth an impact of GPIb α inhibition on TCIPA (see figure 4.3). Therefore, any influence of p0p/B antibodies on melanoma metastasis *in vivo* could, according to these results, not originate from a direct alteration of the potential of B16F10 to induce TCIPA.

5.2 Blockade of GPIb α promotes experimental pulmonary metastasis in mice

The next part of this experimental work was directed at unveiling the role of GPIb α in the metastatic cascade *in vivo*. Therefore a syngeneic model of experimental pulmonary metastasis was used, in which C57BL/6 mice were injected with B16F10 melanoma cells through the lateral tail vein.

For several reasons, this model can only mimic the natural metastatic process in a simplified way. Under physiological conditions, primary tumors can release small numbers of cells that have acquired the ability to metastasize at an early stage of the disease.

This model of experimental metastasis does not represent this process as in this *in vivo* model high numbers of tumor cells are injected into the mouse. The shedding of up to millions of tumor cells into the blood stream is an event which takes place only when the primary tumor has already reached a very advanced stage (Pantel and Brakenhoff 2004; Sahai 2007). Ideally, therapies targeted at the disruption of the metastatic cascade should be aimed at intercepting the early stages of tumor cell dissemination rather than the more advanced stages in this process which can be mimicked by models of experimental metastasis. This is clearly a drawback of experimental models. Yet, compared to spontaneous metastasis models, where orthotopic tumors metastasize over time in a more natural fashion, injection of a large number of these tumor cells has the advantage of resulting in an easier quantification of tumor cell metastases, thus making a detection of possibly subtle changes in the metastatic potential of the tumor cells upon treatment possible.

Another disadvantage of experimental pulmonary metastasis models might be that cancer cell dissemination is likely to be a continuous process and not a singular event like our tumor cell injection. However, the injection of a determined number of tumor cells

at one well defined time point has the advantage of allowing a better understanding of the kinetics of metastasis in relation to any anti- or prometastatic treatment conducted before or after tumor cell inoculation. Given that one aim of this study was to find out at which point in the process of metastasis GPIIb α would play a role, it was crucial to time the injection of melanoma cells and treatment with the antibody. Furthermore, an orthotopic model of melanoma dissemination would have made a continuous therapy with our antibody necessary, as there would have been no way of knowing when metastasis would take place and when the receptors have to be blocked.

As the p0p/B antibody is a rat antibody against a murine epitope, such treatment would have led to an immune response, which would have rendered any evaluation of specific, antibody-generated effects imponderable. Overall, the model chosen here is reproducible and validated, which is not the case for natural tumors.

Generally, the impact of platelet receptors on metastasis has long been recognized and different platelet proteins constitute attractive therapeutic targets to diminish metastasis (Erpenbeck and Schön 2010).

An interesting candidate for this therapeutic approach could be, for example, the platelet integrin receptor $\alpha_{IIb}\beta_3$ (GPIIb/IIIa), a protein with a long career as a metastasis-related molecule, as explained in chapter 2.2. As could be corroborated in this work, experimental pulmonary metastasis under blockade of GPIIb/IIIa leads to a severely reduced formation of pulmonary melanoma colonies (see figure 4.6 on page 35).

This receptor blockade could, in the future, be therapeutically relevant as GPIIb/IIIa inhibitors exist and have been approved to decrease the incidence of ischemic complications in patients undergoing percutaneous coronary artery intervention (De Luca et al. 2009). These new therapeutics include abciximab (a Fab fragment of a chimeric human-mouse antibody), eptifibatide (a cyclic heptapeptide), and tirofiban (a tyrosine-derived nonpeptide molecule). It is conceivable that these or similar inhibitory drugs could be used in the future to protect patients in advanced stages of cancer or even in early stages, when micrometastases might already form. Of course, other inhibitors of platelet receptors, which are important in the process of thrombus formation, might be expected to be similarly promising drugs. Therefore, unveiling of the importance of GPIIb α in this cascade seemed a promising starting point for the development of new therapeutic anti-metastatic strategies.

For this reason, GPIIb α was inhibited, again using monovalent monoclonal antibodies against the extracellular binding region of the receptor, functionally inhibiting the receptor without inducing thrombocytopenia. It is important to emphasize that this treatment did not reduce platelet counts, as thrombocytopenia is induced by virtually

all so-called targeted therapies proposed as antitumoral treatments, and may, thus, interfere with metastasis in a rather nonspecific manner. Surprisingly, a considerable increase in the number of metastatic colonies which had formed 10 days after performing the experimental metastasis assay was detected (see figure 4.6).

This directly contradicts several previous results of other groups which showed a supportive effect of GPIb α in pulmonary metastasis. Jain et al. had provided evidence for a pro-metastatic effect of GPIb α in GPIb α -deficient mice, showing that a functional absence of GPIb-V-IX correlates with a 15-fold reduction in the number of lung metastatic foci using B16F10.1 melanoma cells in a GPIb α -knockout mouse (Jain et al. 2007). However, it must be considered that mice lacking functional GPIb α present a phenotype comparable to the human Bernard Soulier syndrome (Caen et al. 1976; Kato et al. 2004). These mice do not only have giant and abnormal platelets with an aberrant membrane function, they also suffer from thrombocytopenia, leading to a severe bleeding disorder. In fact, it has been shown that GPIb α is essential for the normal membrane development of megakaryocytes (Poujol et al. 2002). It is, therefore, difficult to estimate whether an increase of metastasis in these knockout mice can be attributed to the specific abrogation of GPIb α functions or whether the general platelet dysfunction leads to this impairment, as functional platelets are a prerequisite for a successful metastatic cascade. Even transgenic models that present relatively minor genetic modifications of the GPIb α receptor cannot be trusted to have a completely normal platelet count or a normal membrane development, as has been shown by the experimental work of Kanaji et al., where a mouse model with a GPIb α receptor was used in which most of the extracytoplasmic sequence of GPIb α had been replaced by the human interleukin-4 receptor (Kanaji et al. 2002). Even though the severe makrothrombocytopenia seen in GPIb α null mice could be somewhat ameliorated, platelet counts did not return to normal. Therefore, models using any sort of genetic alteration of GPIb α provide important data concerning the function of GPIb α in thrombus formation and in other physiological processes. However, especially in the context of metastasis, they must be complemented by alternative experimental settings, among them the blockade of GPIb α by monoclonal antibodies that have been proven not to induce thrombocytopenia.

To determine the timeframe in which GPIb α is important for tumor cell dissemination, several experiments were conducted where antibody-mediated blockade of GPIb α was carried out either before, together with or after tumor cell inoculation (see figure 4.9). Interestingly, receptor blockade only influenced metastasis when the antibody was applied either before or, at the latest, together with the tumor cells. An application two hours later had no effect whatsoever. This implies that GPIb α plays a role in the initial arrest and adhesion of the tumor cells rather than in later steps such as extravasation

and tumor growth. The influence of GPIb α shows similar dynamics as the impact of other platelet- or endothelial cell receptors known to be important for the primary interaction of tumor cells with the vasculature, such as P-selectin (Ludwig et al. 2004).

As the experimental system of experimental pulmonary metastasis has the disadvantage of allowing only an endpoint analysis of pulmonary metastases that have developed after several days, a modified assay of pulmonary metastasis was designed to evaluate the immediate effects of GPIb α blockade on hematogenous dissemination. To this end, B16F10 melanoma cells were stably transfected with a GFP variant (pmaxFP-Green-C) so that their fate in the lungs could be traced both by fluorescence microscopy and by flow cytometry. When GFP-positive melanoma cells were injected into the lateral tail vein of mice that had either received p0p/B monovalent antibody or rat IgG control antibody, it was possible to analyze tumor cell retainment in the lungs after 30 minutes, 2 hours or 5 hours. By counting the GFP-positive cells in the lungs, using a fluorescence microscope, significant differences between the groups that had received p0p/B or the control antibody could be detected. In fact, as early as 30 minutes after tumor cell inoculation, blockade of GPIb α led to a clear promotion of tumor cell accumulation in the lungs. While the total number of cells that had been arrested in the lungs decreased as more time passed from tumor cell inoculation to the moment the mice were sacrificed, the effect produced by p0p/B was still visible at later time points, even though less pronounced. At 5 hours, no statistical significance between the two groups could be observed anymore. However, this might have been due to large interindividual differences of the basic values between the groups of different experiments. When comparing the values from single experiments, there was always a pronounced difference between the group that had received p0p/B antibody Fab fragments and the group that had been given only isotype control.

This confirms the previous finding that GPIb α is likely to be important in the primary steps of tumor cell dissemination, that is, the initial adhesion to the vasculature of the lung. The observation that the number of tumor cells which are retained in the lung decreases rapidly over the first 5 hours might seem rather unexpected at first. However, assays using different methods to label the tumor cells in models of experimental metastasis could show a comparably fast decline in the number of arrested tumor cells (Terraube et al. 2006). Possibly, this is due to mechanical destruction of the tumor cells within the lung vasculature by shear forces or to a lack of stimulatory signals that would drive the tumor cell to extravasate and finally form micrometastases. As has been shown in many models, the primary retention of tumor cells in the metastatic organ occurs mainly by size restriction, a primary mechanical process (Luzzi et al. 1998; Chambers et al. 2001). This would explain the large numbers of tumor cells that become retained in the lung of the mice 30 minutes after injection. However, once the

cells have traveled to an organ, their further survival also depends on the regulation of the molecular interactions between cancer cells and the environment of the new organ (Chambers et al. 2002). In a model like the pulmonary metastasis assay, where a great number of cells is injected into an organism that has not “adapted” to tumor cell burden, it appears possible that a lack of these interactions causes the poor survival of the tumor cells.

Confirming these findings obtained by fluorescence microscopy, elevated numbers of tumor cells were also detected when GFP-positive cells in the lungs were measured in relation to the total number of cells by flow cytometry. Unfortunately, this method yielded several difficulties as it turned out to be rather difficult to compare the numbers of retained tumor cells in mice of different experiments, that is, of different days. This could be due to differences between the tumor cell batches used. For example, it is conceivable that the tumor cells were not in the same phase of the cell cycle, when they were used, even though experimental procedures were standardized as far as possible. As other parts of these same lungs could be easily compared with each other by fluorescence microscopy, regardless of which day they were harvested, it seems more likely that the very long and laborious experimental procedure of preparing a single cell suspension from the lungs caused a certain variation of the basic level of tumor cells that could be detected within the lungs. However, as experiments were carried out three times, it was possible to compare these groups with one another in three independent experiments. Even though the baseline of tumor cell dissemination was different between the experiments, all three experiments (which are displayed separately in figure 4.11) point towards the same result: GPIb α blockade increases initial tumor cell accumulation in the lungs as early as 30 minutes after inoculation.

5.3 Interaction of the integrin receptor Mac-1 with GPIb α is not responsible for the promotion of metastasis upon GPIb α blockade

In order to find a mechanistic explanation for the effect seen after GPIb α blockade, attention was focused on the leukocyte integrin, Mac-1, a binding partner for GPIb α (Simon et al. 2000; Ehlers et al. 2003) expressed by neutrophils, monocytes and NK cells (Arnaout 1990; Li 1999). Apart from GPIb α , Mac-1 recognizes a multitude of different protein and non-protein ligands such as, among many others, fibronectin, laminin, collagen, fibrinogen and vitronectin, ICAM-1 as well as coagulation factor X, the complement pathway product C3bi and non-protein ligands such as LPS, beta-glycans or heparin. These diverse interactions explain the importance of Mac-1 in a multitude of

processes which include leukocyte adhesion, chemotaxis, migration, phagocytosis and cytotoxicity (Li 1999). Recent investigations have shown an important role for Mac-1 in *in vivo* antitumor activity, specifically in the antibody-mediated, FcR-dependent cytotoxicity which requires the action of phagocytes, specifically neutrophils (van Spriel et al. 2003). In a model of pulmonary metastasis which used B16F10 melanoma cells that were intravenously injected into C57BL/6 mice, migration of neutrophils and binding to tumor cells was not abrogated in mice that lacked the Mac-1 receptor. However, upon treatment of the mice with TA99 monoclonal antibodies directed against the melanoma cells, Mac-1 null mice showed a lack of response compared to wildtype mice, where a protection from melanoma metastasis could be achieved by antibody-treatment (van Spriel et al. 2003).

As GPIb α clearly played a role in the early phases of tumor cell dissemination, an involvement of GPIb α in FcR mediated antitumoral protection seemed unlikely. Nevertheless, it was possible that Mac-1 is involved in additional natural defense mechanisms together with GPIb α , especially as other sets of cells belonging to the innate immunity, such as NK cells, also express this receptor and are known to be important in the initial defense against melanoma cells, a function from which platelets directly protect disseminating tumor cells in the blood stream (Nieswandt et al. 1999). Therefore, the recruitment of Mac-1 positive cells was evaluated under blockade of GPIb α after injection of melanoma cells into wildtype mice. Following the injection of the melanoma cells, a marked inflammatory response could be seen in the lungs as early as 30 minutes after inoculation. However, in line with the results obtained by van Spriel et al., no difference was observed between those mice that had received only isotype control and those that had been treated with p0p/B mAb prior to tumor cell injection (see figure 4.13).

In a complementary approach, Mac-1 was blocked in a long-term metastasis experiment either solitarily or together with GPIb α to test for synergistic or complementary effects between these two receptors. Contrary to results published by van Spriel et al., a statistically significant difference in the tumor colony formation of mice that had received anti Mac-1 antibody and those that had received only isotype control was perceived. Tumor defense seemed to be seriously impaired when Mac-1 was inhibited (see figure 4.14).

Why other groups saw no difference in long term metastasis in Mac-1 deficient mice as compared to knockout mice as long as mice were not treated with a specific immunological treatment against melanoma epitopes, cannot be satisfactorily explained at the moment (van Spriel et al. 2003). It must be considered, however, that van Spriel et al. focused mainly on the behavior of neutrophils in the neighborhood of tumor cells

and saw no difference in the behavior of those cells that expressed functional Mac-1 receptors and those that completely lacked this molecule.

The experiments conducted here, using monoclonal anti Mac-1 antibody that had been injected intraperitoneally into wildtype mice, may not have reached all Mac-1 positive cells in every compartment of the organism and may therefore have affected processes such as neutrophil migration towards tumor cells, binding to the metastatic cells or their phagocytosis or cell lysis in a different way than complete Mac-1 deficiency. Furthermore, even though antibodies have long half-lives (Zalevsky et al. 2010) and could therefore influence physiological processes for several days, there would still be a certain decline in antibody concentrations in the mice and this might again lead to different results than in Mac-1 deficient mice.

In accordance to previous experiments, GPIb α blockade again led to a significant increase in metastasis in a similar dimension as Mac-1 blockade (4.14). However, when both receptors were blocked in a joint experimental approach, an additive effect that led to an even higher number of lung metastases after 10 days was observed. This does not rule out an interaction between the two molecules in the context of tumor defense. It does, however, make the concept unlikely that the interaction between GPIb α and Mac-1 is the exclusive and decisive factor in the effects seen after GPIb α blockade.

So while this experimental approach further confirms the importance of GPIb α and establishes an important role for Mac-1 in melanoma metastasis, an interaction of both receptors in terms of an anti-tumoral defense mechanism does not provide a satisfactory explanation for the here achieved results, as the effects of GPIb α and Mac-1 blockade appear to be additive rather than complementary.

5.4 GPIb α blockade does not lead to enhanced metastasis in P-selectin deficient mice

The hypothesis that P-selectin, another important binding partner of GPIb α (see chapter 2.4.2 on page 13), was involved in the metastasis promoting effects seen after GPIb α inhibition was tested next. Generally, P-selectin, together with E-selectin and L-selectin, belongs to the C-type lectins, which recognize sialylated, fucosylated carbohydrates, specifically glycans containing a terminal sialyl Lewis^x (sLe^x), often in a calcium-dependent fashion. Selectins mediate transient leukocyte or platelet contacts with vascular endothelial cells (Vestweber and Blanks 1999).

For platelets that roll on activated endothelium this behavior was shown to be both P-selectin and GPIb α dependent. Interestingly, contrary to the interaction of P-selectin with its major ligand, P-selectin glycoprotein ligand-1 (PSGL-1), this interaction was

not calcium dependent (Romo et al. 1999). However, P-selectin is not only crucial for these platelet-endothelial-interactions but also mediates the binding of platelets to tumor cells via sialylated fucosylated glycans displayed on mucin and non-mucin structures on tumor cells (Mannori et al. 1995; Borsig et al. 2002). Inhibition of P-selectin abrogates the coating of tumor cells by platelets *in vivo* (Borsig et al. 2001) and platelets from P-selectin-deficient mice fail to adhere to tumor cells *in vitro* and *in vivo* (Kim et al. 1998; Ludwig et al. 2004). Inhibition of P-selectin *in vivo* or experimental metastasis assays in P-selectin knockout mice show a severely reduced colonization of the lungs of these mice, as compared to wildtype mice (Kim et al. 1998). Interestingly though, the influence of P-selectin on metastasis is not limited to platelet tumor cell interactions.

Several melanoma cell lines appear to interact with P-selectin displayed on vascular endothelial cells and this interaction seems to be of importance to the metastatic cascade. This can be shown by performing experimental melanoma metastasis assays in P-selectin-deficient mice and in bone marrow transplants that selectively lack the platelet P-selectin while displaying intact P-selectin on the vascular endothelium. In complete P-selectin knockouts, metastasis is much more severely compromised than in the mice lacking only platelet P-selectin, which shows that endothelial P-selectin accounts in part for the role of P-selectin in metastasis. The observation that human melanoma cell lines may directly interact with the vascular endothelium in a P-selectin dependent manner corroborates this hypothesis (Ludwig et al. 2004). It is of great interest that heparin has been shown to effectively block P-selectin interactions by mimicking carbohydrate ligands which constitute the natural binding motifs of P-selectin (Stevenson et al. 2007). This discovery may bear great therapeutic potential for future cancer treatments (although clinical observations were somewhat sobering so far). Because of its importance in the process of metastasis, its role as a ligand of GPIIb/IIIa and also because of the accessibility of novel therapeutic approaches in anti-cancer therapy linked to P-selectin, this molecule was interesting to study in the context of the results obtained by blocking GPIIb/IIIa in experimental melanoma metastasis.

To reveal a possible link between the effects observed after GPIIb/IIIa blockade and P-selectin, the experimental pulmonary metastasis assay was repeated in P-selectin deficient mice with a C57BL/6 genetic background, using wild type C57BL/6 mice as controls. Again, the results obtained in the wild type group confirmed the previous findings concerning the blockade of GPIIb/IIIa and the subsequent elevation of the number of metastases. In the P-selectin knockout group, metastasis was severely diminished, as was to be expected considering the above mentioned role of P-selectin in metastasis. Remarkably though, the number of metastases did not change significantly upon

blockade of GPIb α in this same group. This infers that there might be a mechanistic link between GPIb α , its importance in metastasis and its role as a binding partner of P-selectin. It might at first seem strange that the inhibition of an interaction of two molecules which are both important for platelet function would lead to an increase in metastasis. Considering the importance of most other platelet surface proteins (see figure 2.2), one would expect the opposite effect, namely an inhibition of metastasis. However, the aggregometry experiments in this study have shown that blockade of GPIb α does not interfere with tumor cell-induced platelet aggregation. Therefore, a different phenomenon than the simple interference with platelet aggregation appears to be responsible for any effects GPIb α inhibition could have on metastasis.

Fig. 5.1: Proposed effect of GPIb α inhibition on tumor-platelet-endothelial and tumor cell-endothelial interactions. Under physiological conditions, platelets expressing the GPIb-X-IV complex can interact with activated endothelium expressing P-selectin. After blockade of GPIb α , platelet interaction with endothelial cells is impeded and tumor cells expressing sialylated fucosylated glycans can directly interact with the endothelium in the metastatic organ which would lead to tumor cell adhesion and subsequent extravasation.

The following hypothesis can be proposed: A decrease in the interactions between platelet GPIb α with endothelial P-selectin could lead to an increased availability of endothelial P-selectin for the tumor cells which could then directly interact with the endothelium and adhere more efficiently to the vessel walls in the metastatic organ. This hypothesis is supported by the above mentioned observation that endothelial P-selectin clearly contributes to the formation of metastases (Ludwig et al. 2004). One could propose that the rolling of tumor cells on activated endothelium, a P-selectin

dependent process, could outweigh the advantages of an indirect mediation of contacts between tumor cells *via* surrounding platelet thrombi, as illustrated in figure 5.1.

Another, different approach to explain both the influence of GPIb α on melanoma metastasis as well as the lack of response in P-selectin deficient mice results from the observation that vWF knockout mice present a very similar phenotype in experimental melanoma metastasis (Terraube et al. 2007). As mentioned above, GPIb α interacts with vWF mainly under conditions of very high shear. An arrest of melanoma cells under these conditions could possibly constitute high mechanical stress for these cells. If this interaction between GPIb α and vWF was blocked, melanoma cells would be able to move on to regions of lesser shear, as illustrated in figure 5.2.

Fig. 5.2: Proposed effect of GPIb α inhibition on tumor cell arrest under conditions of higher or lower shear. Activated platelets surrounding the tumor cell could interact with vWF under conditions of high shear, leading to an arrest of tumor cells under mechanically unfavorable conditions. After blockade of GPIb α , platelet interaction with vWF is impeded and tumor cells could move on to smaller vessels with lower shear, where an adhesion and subsequent extravasation might be easier.

A recent report had proposed a direct interaction between vWF and the melanoma cells as in some preliminary experiments they had observed an induction of cellular death by vWF without any influence on cell proliferation or invasion of the melanoma cells (Terraube et al. 2007). To confirm this data, an anti-histone ELISA was conducted to detect death of melanoma cells induced directly by vWF, using up to twice the physiological concentration. However, no differences between the untreated cells and those treated with vWF was seen, so that a direct apoptotic influence of vWF on B16F10 melanoma

cells appears highly unlikely. Of note, however, is the slightly different timeframe in which vWF-deficiency appears to become important in the experimental metastasis. While a clear effect of GPIb α -inhibition was seen as early as 30 minutes after tumor cell injection in this study, Terraube et al. found no differences in the vWF deficient and the wildtype control mice until 4 hours after injection. It remains to be investigated whether this difference can be explained by different experimental procedures or whether the elevation of the number of pulmonary metastases in both cases is a mere coincidence and has no functional connection.

5.5 GPVI in experimental metastasis

GPVI constitutes an important molecule in the process of platelet activation in its role of one of the principal collagen receptors in platelet function (Nieswandt et al. 2001a, see chapter 2.3 on page 10). Very little, however, is known concerning its possible role in the metastatic cascade. It therefore seemed a logical step to investigate the importance of GPVI in the experimental system, using an assay of experimental melanoma metastasis in C57BL/6 mice and B16F10 melanoma cells. However, no alteration of the behavior of the metastasizing melanoma cells was seen. There was neither a conceivable change in the aspect of grown metastases nor a significant change in the number of metastatic foci (see figure 4.18).

Of course, it cannot be formally excluded from these experiments that a collagen receptor with such a central role in platelet adhesion and aggregation may also play a role in metastasis. It might nevertheless be interesting, to further investigate the behavior of GPVI in the context of metastasis, using different experimental methods.

5.6 Conclusion

In conclusion, the findings of this work demonstrate for the first time that a platelet surface receptor important for platelet adhesion and subsequent thrombus formation may also have an anti-metastatic function in the sense that its blockade in metastasis leads to an increase of experimental metastasis. This is especially important in light of the fact that over the years, the inhibition of many factors and receptors linked to platelet aggregation has been discussed intensively with the purpose of finding new tumor targets and ways to inhibit metastasis. It must be considered that, as shown here, not all platelet receptors can be seen as adequate targets in tumor therapy. On the contrary, especially when talking about GPIb α , this must be debated very carefully,

as possibly the contrary effect of the desirable attenuation of metastatic potential of the tumor entity could ensue. It will be necessary to further investigate this subject by incorporating different tumor cell lines in the studies and to use complementary, possibly spontaneous, models of tumor cell metastasis to test the influence of GPIIb α .

6 Bibliography

Alonso-Escolano D, Strongin AY, Chung AW, Deryugina EI, Radomski MW (2004): Membrane type-1 matrix metalloproteinase stimulates tumour cell-induced platelet aggregation: role of receptor glycoproteins. *Br J Pharmacol* 141, 241-252

Andrews RK, Gardiner EE, Shen Y, Whisstock JC, Berndt MC (2003): Glycoprotein Ib-IX-V. *Int J Biochem Cell Biol* 35, 1170-1174

Andrews RK, Gardiner EE, Shen Y, Berndt MC (2004): Platelet interactions in thrombosis. *IUBMB Life* 56, 13-18

Arnaout MA (1990): Structure and function of the leukocyte adhesion molecules CD11/CD18. *Blood* 75, 1037-1050

Asaduzzaman M, Lavasani S, Rahman M, Zhang S, Braun OO, Jeppsson B, Thorlacius H (2009): Platelets support pulmonary recruitment of neutrophils in abdominal sepsis. *Crit Care Med* 37, 1389-1396

Bastida E, Almirall L, Ordinas A (1987): Tumor-cell-induced platelet aggregation is a glycoprotein-dependent and lipoxygenase-associated process. *Int J Cancer* 39, 760-763

Batzler WU, Giersiepen K, Hentschel S, Husmann G, Kaatsch P, Katalinic A, Kischke J, Kraywinkel K, Meyer M, Stabenow R, Stegmaier C: Krebs in Deutschland 2003-2004. Häufigkeiten und Trends. 6. überarbeitete Auflage. Robert Koch-Institut (Hrsg) und die Gesellschaft der epidemiologischen Krebsregister in Deutschland e. V. (Hrsg). Berlin 2008

Bergmeier W, Rackebrandt K, Schroder W, Zirngibl H, Nieswandt B (2000): Structural and functional characterization of the mouse von Willebrand factor receptor GPIb-IX with novel monoclonal antibodies. *Blood* 95, 886-893

Bergmeier W, Schulte V, Brockhoff G, Bier U, Zirngibl H, Nieswandt B (2002): Flow cytometric detection of activated mouse integrin $\alpha IIb\beta 3$ with a novel monoclonal antibody. *Cytometry* 48, 80-86

Biggerstaff JP, Seth N, Amirkhosravi A, Amaya M, Fogarty S, Meyer TV, Siddiqui F, Francis JL (1999): Soluble fibrin augments platelet/tumor cell adherence in vitro and in vivo, and enhances experimental metastasis. *Clin Exp Metastasis* 17, 723-730

Borsig L (2007): Antimetastatic activities of modified heparins: selectin inhibition by

heparin attenuates metastasis. *Semin Thromb Hemost* 33, 540-546

Borsig L, Wong R, Feramisco J, Nadeau DR, Varki NM, Varki A (2001): Heparin and cancer revisited: mechanistic connections involving platelets, P-selectin, carcinoma mucins, and tumor metastasis. *Proc Natl Acad Sci U S A* 98, 3352-3357

Borsig L, Wong R, Hynes RO, Varki NM, Varki A (2002): Synergistic effects of L- and P-selectin in facilitating tumor metastasis can involve non-mucin ligands and implicate leukocytes as enhancers of metastasis. *Proc Natl Acad Sci U S A* 99, 2193-2198

Caen JP, Nurden AT, Jeanneau C, Michel H, Tobelem G, Levy-Toledano S, Sultan Y, Valensi F, Bernard J (1976): Bernard-Soulier syndrome: a new platelet glycoprotein abnormality. Its relationship with platelet adhesion to subendothelium and with the factor VIII von Willebrand protein. *J Lab Clin Med* 87, 586-596

Camerer E, Qazi AA, Duong DN, Cornelissen I, Advincula R, Coughlin SR (2004): Platelets, protease-activated receptors, and fibrinogen in hematogenous metastasis. *Blood* 104, 397-401

Chambers AF, Naumov GN, Varghese HJ, Nadkarni KV, MacDonald IC, Groom AC (2001): Critical steps in hematogenous metastasis: an overview. *Surg Oncol Clin N Am* 10, 243-255, vii

Chambers AF, Groom AC, MacDonald IC (2002): Dissemination and growth of cancer cells in metastatic sites. *Nat Rev Cancer* 2, 563-572

Clemetson JM, Polgar J, Magnenat E, Wells TN, Clemetson KJ (1999): The platelet collagen receptor glycoprotein VI is a member of the immunoglobulin superfamily closely related to Fc α R and the natural killer receptors. *J Biol Chem* 274, 29019-29024

Davey MG, Luscher EF (1967): Actions of thrombin and other coagulant and proteolytic enzymes on blood platelets. *Nature* 216, 857-858

De Luca G, Navarese E, Marino P (2009): Risk profile and benefits from Gp IIb-IIIa inhibitors among patients with ST-segment elevation myocardial infarction treated with primary angioplasty: a meta-regression analysis of randomized trials. *Eur Heart J* 30, 2705-2713

Du X, Fox JE, Pei S (1996): Identification of a binding sequence for the 14-3-3 protein within the cytoplasmic domain of the adhesion receptor, platelet glycoprotein Ib alpha. *J Biol Chem* 271, 7362-7367

Ehlers R, Ustinov V, Chen Z, Zhang X, Rao R, Luscinskas FW, Lopez J, Plow E, Simon DI (2003): Targeting platelet-leukocyte interactions: identification of the integrin Mac-1 binding site for the platelet counter receptor glycoprotein Ib α . *J Exp Med* 198, 1077-1088

- Erpenbeck L, Schön MP (2010): Deadly allies: the fatal interplay between platelets and metastasizing cancer cells. *Blood* 115, 3427-3436
- Erpenbeck L, Nieswandt B, Schön M, Pozgajova M, Schön MP Inhibition of platelet GPIb alpha and promotion of melanoma metastasis (2009): *J Invest Dermatol* 130, 576-586
- Ezumi Y, Shindoh K, Tsuji M, Takayama H (1998): Physical and functional association of the Src family kinases Fyn and Lyn with the collagen receptor glycoprotein VI-Fc receptor gamma chain complex on human platelets. *J Exp Med* 188, 267-276
- Fredrickson BJ, Dong JF, McIntire LV, Lopez JA (1998): Shear-dependent rolling on von Willebrand factor of mammalian cells expressing the platelet glycoprotein Ib-IX-V complex. *Blood* 92, 3684-3693
- Gasic GJ (1984): Role of plasma, platelets, and endothelial cells in tumor metastasis. *Cancer Metastasis Rev* 3, 99-114
- Gasic GJ, Gasic TB, Stewart CC (1968): Antimetastatic effects associated with platelet reduction. *Proc Natl Acad Sci U S A* 61, 46-52
- Grossi IM, Fitzgerald LA, Kendall A, Taylor JD, Sloane BF, Honn KV (1987): Inhibition of human tumor cell induced platelet aggregation by antibodies to platelet glycoproteins Ib and IIb/IIIa. *Proc Soc Exp Biol Med* 186, 378-383
- Hamberg M, Svensson J, Samuelsson B (1975): Thromboxanes: a new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc Natl Acad Sci U S A* 72, 2994-2998
- Hart IR (1979): The selection and characterization of an invasive variant of the B16 melanoma. *Am J Pathol* 97, 587-600
- Henn V, Slupsky JR, Gräfe M, Anagnostopoulos I, Förster R, Müller-Berghaus G, Kroczeck RA (1998): CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* 391, 591-594
- Hynes RO (2002): Integrins: bidirectional, allosteric signaling machines. *Cell* 110, 673-687
- Jackson SP, Nesbitt WS, Kulkarni S (2003): Signaling events underlying thrombus formation. *J Thromb Haemost* 1, 1602-1612
- Jain S, Zuka M, Liu J, Russell S, Dent J, Guerrero JA, Forsyth J, Maruszak B, Gärtner TK, Felding-Habermann B et al. (2007): Platelet glycoprotein Ib alpha supports experimental lung metastasis. *Proc Natl Acad Sci U S A* 104, 9024-9028
- Kanaji T, Russell S, Ware J (2002): Amelioration of the macrothrombocytopenia associated with the murine Bernard-Soulier syndrome. *Blood* 100, 2102-2107

Karpatkin S, Pearlstein E, Ambrogio C, Collier BS (1988): Role of adhesive proteins in platelet tumor interaction in vitro and metastasis formation in vivo. *J Clin Invest* 81, 1012-1019

Kato K, Martinez C, Russell S, Nurden P, Nurden A, Fiering S, Ware J (2004): Genetic deletion of mouse platelet glycoprotein Ibbeta produces a Bernard-Soulier phenotype with increased alpha-granule size. *Blood* 104, 2339-2344

Kemp RB, Jones BM, Groschel-Stewart U (1971): Aggregative behaviour of embryonic chick cells in the presence of antibodies directed against actomyosins. *J Cell Sci* 9, 103-122

Kepner N, Lipton A (1981): A mitogenic factor for transformed fibroblasts from human platelets. *Cancer Res* 41, 430-432

Kim YJ, Borsig L, Varki NM, Varki A (1998): P-selectin deficiency attenuates tumor growth and metastasis. *Proc Natl Acad Sci U S A* 95, 9325-9330

Kopp HG, Placke T, Salih HR (2009): Platelet-derived transforming growth factor-beta down-regulates NKG2D thereby inhibiting natural killer cell antitumor reactivity. *Cancer Res* 69, 7775-7783

Li Z (1999): The alphaMbeta2 integrin and its role in neutrophil function. *Cell Res* 9, 171-178

Ludwig RJ, Boehme B, Podda M, Henschler R, Jäger E, Tandi C, Boehncke WH, Zollner TM, Kaufmann R, Gille J (2004): Endothelial P-selectin as a target of heparin action in experimental melanoma lung metastasis. *Cancer Res* 64, 2743-2750

Luzzi KJ, MacDonald IC, Schmidt EE, Kerkvliet N, Morris VL, Chambers AF, Groom AC (1998): Multistep nature of metastatic inefficiency: dormancy of solitary cells after successful extravasation and limited survival of early micrometastases. *Am J Pathol* 153, 865-873

Mannori G, Crottet P, Cecconi O, Hanasaki K, Aruffo A, Nelson RM, Varki A, Bevilacqua MP (1995): Differential colon cancer cell adhesion to E-, P-, and L-selectin: role of mucin-type glycoproteins. *Cancer Res* 55, 4425-4431

Markovic SN, Erickson LA, Rao RD, Weenig RH, Pockaj BA, Bardia A, Vachon CM, Schild SE, McWilliams RR, Hand JL et al. (2007a): Malignant melanoma in the 21st century, part 1: epidemiology, risk factors, screening, prevention, and diagnosis. *Mayo Clin Proc* 82, 364-380

Markovic SN, Erickson LA, Rao RD, Weenig RH, Pockaj BA, Bardia A, Vachon CM, Schild SE, McWilliams RR, Hand JL et al. (2007b): Malignant melanoma in the 21st century, part 2: staging, prognosis, and treatment. *Mayo Clin Proc* 82, 490-513

- Massberg S, Gawaz M, Gruner S, Schulte V, Konrad I, Zohlnhofer D, Heinzmann U, Nieswandt B (2003): A crucial role of glycoprotein VI for platelet recruitment to the injured arterial wall in vivo. *J Exp Med* 197, 41-49
- Melnikova VO, Villares GJ, Bar-Eli M (2008): Emerging Roles of PAR-1 and PAFR in Melanoma Metastasis. *Cancer Microenviron* 1, 103-111
- Mohle R, Green D, Moore MA, Nachman RL, Rafii S (1997): Constitutive production and thrombin-induced release of vascular endothelial growth factor by human megakaryocytes and platelets. *Proc Natl Acad Sci U S A* 94, 663-668
- Moog S, Mangin P, Lenain N, Strassel C, Ravanat C, Schuhler S, Freund M, Santer M, Kahn M, Nieswandt B et al. (2001): Platelet glycoprotein V binds to collagen and participates in platelet adhesion and aggregation. *Blood* 98, 1038-1046
- Munday AD, Berndt MC, Mitchell CA (2000): Phosphoinositide 3-kinase forms a complex with platelet membrane glycoprotein Ib-IX-V complex and 14-3-3zeta. *Blood* 96, 577-584
- Nierodzik ML, Karparkin S (2006): Thrombin induces tumor growth, metastasis, and angiogenesis: Evidence for a thrombin-regulated dormant tumor phenotype. *Cancer Cell* 10, 355-362
- Nierodzik ML, Plotkin A, Kajumo F, Karparkin S (1991): Thrombin stimulates tumor-platelet adhesion in vitro and metastasis in vivo. *J Clin Invest* 87, 229-236
- Nieswandt B, Watson SP (2003): Platelet-collagen interaction: is GPVI the central receptor? *Blood* 102, 449-461
- Nieswandt B, Hafner M, Echtenacher B, Männel DN (1999): Lysis of tumor cells by natural killer cells in mice is impeded by platelets. *Cancer Res* 59, 1295-1300
- Nieswandt B, Bergmeier W, Rackebrandt K, Gessner JE, Zirngibl H (2000a): Identification of critical antigen-specific mechanisms in the development of immune thrombocytopenic purpura in mice. *Blood* 96, 2520-2527
- Nieswandt B, Bergmeier W, Schulte V, Rackebrandt K, Gessner JE, Zirngibl H (2000b): Expression and function of the mouse collagen receptor glycoprotein VI is strictly dependent on its association with the FcRgamma chain. *J Biol Chem* 275, 23998-24002
- Nieswandt B, Brakebusch C, Bergmeier W, Schulte V, Bouvard D, Mokhtari-Nejad R, Lindhout T, Heemskerk JW, Zirngibl H, Fässler R (2001a): Glycoprotein VI but not alpha2beta1 integrin is essential for platelet interaction with collagen. *Embo J* 20, 2120-2130
- Nieswandt B, Schulte V, Bergmeier W, Mokhtari-Nejad R, Rackebrandt K, Cazenave JP, Ohlmann P, Gachet C, Zirngibl H (2001b): Long-term antithrombotic protection by in vivo depletion of platelet glycoprotein VI in mice. *J Exp Med* 193, 459-469

- Otten HM, Mathijssen J, ten Cate H, Soesan M, Inghels M, Richel DJ, Prins MH (2004): Symptomatic venous thromboembolism in cancer patients treated with chemotherapy: an underestimated phenomenon. *Arch Intern Med* 164, 190-194
- Palumbo JS, Kombrinck KW, Drew AF, Grimes TS, Kiser JH, Degen JL, Bugge TH (2000): Fibrinogen is an important determinant of the metastatic potential of circulating tumor cells. *Blood* 96, 3302-3309
- Palumbo JS, Talmage KE, Massari JV, La Jeunesse CM, Flick MJ, Kombrinck KW, Jirouskova M, Degen JL (2005): Platelets and fibrin(ogen) increase metastatic potential by impeding natural killer cell-mediated elimination of tumor cells. *Blood* 105, 178-185
- Pantel K, Brakenhoff RH (2004): Dissecting the metastatic cascade. *Nat Rev Cancer* 4, 448-456
- Poujol C, Ware J, Nieswandt B, Nurden AT, Nurden P (2002): Absence of GPIb α is responsible for aberrant membrane development during megakaryocyte maturation: ultrastructural study using a transgenic model. *Exp Hematol* 30, 352-360
- Romo GM, Dong JF, Schade AJ, Gardiner EE, Kansas GS, Li CQ, McIntire LV, Berndt MC, Lopez JA (1999): The glycoprotein Ib-IX-V complex is a platelet counter-receptor for P-selectin. *J Exp Med* 190, 803-814
- Schön M, Wienrich BG, Kneitz S, Sennefelder H, Amschler K, Vohringer V, Weber O, Stiewe T, Ziegelbauer K, Schön MP (2008): KINK-1, a novel small-molecule inhibitor of IKK β , and the susceptibility of melanoma cells to antitumoral treatment. *J Natl Cancer Inst* 100, 862-875
- Schulte V, Reusch HP, Pozgajova M, Varga-Szabo D, Gachet C, Nieswandt B (2006): Two-phase antithrombotic protection after anti-glycoprotein VI treatment in mice. *Arterioscler Thromb Vasc Biol* 26, 1640-1647
- Shattil SJ, Newman PJ (2004): Integrins: dynamic scaffolds for adhesion and signaling in platelets. *Blood* 104, 1606-1615
- Simon DI, Chen Z, Xu H, Li CQ, Dong J, McIntire LV, Ballantyne CM, Zhang L, Furman MI, Berndt MC et al. (2000): Platelet glycoprotein Ib α is a counterreceptor for the leukocyte integrin Mac-1 (CD11b/CD18). *J Exp Med* 192, 193-204
- Stevenson JL, Varki A, Borsig L (2007): Heparin attenuates metastasis mainly due to inhibition of P- and L-selectin, but non-anticoagulant heparins can have additional effects. *Thromb Res* 120 Suppl 2, S107-111
- Suzuki-Inoue K, Tulasne D, Shen Y, Bori-Sanz T, Inoue O, Jung SM, Moroi M, Andrews RK, Berndt MC, Watson SP (2002): Association of Fyn and Lyn with the proline-rich domain of glycoprotein VI regulates intracellular signaling. *J Biol Chem* 277, 21561-21566

Terraube V, Pendu R, Baruch D, Gebbink MF, Meyer D, Lenting PJ, Denis CV (2006): Increased metastatic potential of tumor cells in von Willebrand factor-deficient mice. *J Thromb Haemost* 4, 519-526

Terraube V, Marx I, Denis CV (2007): Role of von Willebrand factor in tumor metastasis. *Thromb Res* 120 Suppl 2, S64-70

Todd RF, 3rd (1996): The continuing saga of complement receptor type 3 (CR3). *J Clin Invest* 98, 1-2

Trikha M, Raso E, Cai Y, Fazakas Z, Paku S, Porter AT, Timar J, Honn KV (1998): Role of α IIb β 3 integrin in prostate cancer metastasis. *Prostate* 35, 185-192

Trikha M, Zhou Z, Timar J, Raso E, Kennel M, Emmell E, Nakada MT (2002): Multiple roles for platelet GPIIb/IIIa and α v β 3 integrins in tumor growth, angiogenesis, and metastasis. *Cancer Res* 62, 2824-2833

Trousseau A (1865): Phlegmasia alba dolens. *Clinique Medicale de l'Hotel-Dieu de Paris*. 2nd ed. Paris, France: JB Baillière & Fils 1865, 654-712.

van Spriël AB, van Ojik HH, Bakker A, Jansen MJ, van de Winkel JG (2003): Mac-1 (CD11b/CD18) is crucial for effective Fc receptor-mediated immunity to melanoma. *Blood* 101, 253-258

Vestweber D, Blanks JE (1999): Mechanisms that regulate the function of the selectins and their ligands. *Physiol Rev* 79, 181-213

Wang Y, Sakuma M, Chen Z, Ustinov V, Shi C, Croce K, Zago AC, Lopez J, Andre P, Plow E et al. (2005): Leukocyte engagement of platelet glycoprotein Ib α via the integrin Mac-1 is critical for the biological response to vascular injury. *Circulation* 112, 2993-3000

Wardell MR, Reynolds CC, Berndt MC, Wallace RW, Fox JE (1989): Platelet glycoprotein Ib beta is phosphorylated on serine 166 by cyclic AMP-dependent protein kinase. *J Biol Chem* 264, 15656-15661

Yang J, Furie BC, Furie B (1999): The biology of P-selectin glycoprotein ligand-1: its role as a selectin counterreceptor in leukocyte-endothelial and leukocyte-platelet interaction. *Thromb Haemost* 81, 1-7

Zalevsky J, Chamberlain AK, Horton HM, Karki S, Leung IW, Sproule TJ, Lazar GA, Roopenian DC, Desjarlais JR (2010): Enhanced antibody half-life improves in vivo activity. *Nat Biotechnol* 28, 157-159

List of Figures

2.1	Proposed advantages of tumor cell interactions with platelets	6
2.2	Platelet-tumor cell interactions	8
2.3	The platelet receptor GPVI	10
2.4	The platelet receptor complex GPIb-V-IX	14
3.1	Principles of aggregometry	20
4.1	GPIb α expression on platelets and B16F10 melanoma cells	29
4.2	GPIb α on lung epithelium and on platelets	30
4.3	TCIPA under GPIb α blockade	32
4.4	Determination of optimum antibody concentrations for full GPIb α blockade	33
4.5	Experimental pulmonary metastasis after platelet depletion	34
4.6	Experimental pulmonary metastasis under GPIb α blockade	35
4.7	Histological analysis of lung sections after experimental pulmonary metastasis under GPIb α blockade	37
4.8	Morphometric analysis of lung sections after experimental pulmonary metastasis under GPIb α blockade	38
4.9	Determination of a timeframe for the influence of GPIb α blockade in pulmonary metastasis	39
4.10	Flow cytometric evaluation of GFP expression after stable transfection of melanoma cells	41
4.11	Flow cytometric evaluation of short term-metastasis under blockade of GPIb α	42
4.12	Short term metastasis under GPIb α blockade	43
4.13	Pulmonary expression of Mac-1 after short-term metastasis	46
4.14	Experimental pulmonary metastasis under double blockade of GPIb α with Mac-1	47
4.15	Quantitative evaluation of experimental pulmonary metastasis under double blockade of GPIb α with Mac-1	48
4.16	Experimental pulmonary metastasis under GPIb α blockade in P-selectin deficient mice	50
4.17	Influence of vWF on apoptosis of B16F10 melanoma cells	51
4.18	Experimental pulmonary metastasis under blockade of GPVI	53
5.1	Proposed effect of GPIb α inhibition on tumor-platelet-endothelial and tumor cell-endothelial interactions	64
5.2	Alternative explanation for the effect of GPIb α inhibition on tumor cell arrest	65

List of Tables

3.1	Chemicals	26
3.2	Laboratory equipment	28

Previously published results

Parts of the results presented in this thesis have been published:

Erpenbeck L, Nieswandt B, Schön M, Pozgajova M, Schön MP Inhibition of platelet GPIb alpha and promotion of melanoma metastasis (2009): J Invest Dermatol 130, 576-586

Acknowledgements

I would like to thank Prof. Dr. med MP Schön for his generous support and scientific guidance throughout this project.

Furthermore, I would like to express my sincere appreciation to all those who have contributed, directly or indirectly, to this thesis in form of technical or other support. I particularly want to thank my former laboratory colleagues, Margarete Schön, Katharina Amschler, Katrin Wallbrecht, Stefanie Schlickum, Kai Michaelis, Gregor Wienrich, Helga Sennefelder as well as Attila Braun for the support they gave me and for providing a very pleasant working atmosphere.

I would also like to thank Sebastian Kruss and Vera Keil for proofreading of this manuscript.

Curriculum Vitae

My name is Luise Erpenbeck, I was born in Berlin (Germany) on the 30th of December 1983.

In 2003 I graduated from the Friedrich-Engels-Gymnasium (Berlin, Germany, all-gemeine Hochschulreife) and enrolled for Medical School in Magdeburg (Otto-von-Guericke-Universität, Magdeburg) in autumn 2003 (Wintersemester 2003/04), where I took my pre-clinical examination (1. ärztliche Prüfung) in September 2005.

After completing the first year of clinical education in Magdeburg, I took leave of absence from my university studies for one year in order to work on my medical thesis in the Rudolf-Virchow-Center for Experimental Biomedicine in Göttingen, in the laboratory of Prof. MP Schön and in cooperation with the group of Prof. B Nieswandt.

In September 2007 I resumed my clinical studies in Göttingen (Georg-August-Universität Göttingen) and progressed into the practical year (PJ, medical internship) in August 2009. My medical activities during this year comprised internships in the Department of Dermatology and Venereology at the Academic Medical Center Göttingen (UMG), internal medicine and general surgery in the Kantonsspital Schaffhausen (Schaffhausen, Switzerland) and cardiology, again in the Academic Medical Center Göttingen.

I continued my research work on platelet receptors and melanoma metastasis in Göttingen. In spring 2009 I visited the Fuhlbrigge Lab at the Brigham and Women's Hospital in Harvard, Boston, USA. From September 2006 to September 2008 I took part in the study group "Oncological cell biology" of the German National Academic Foundation (Studienstiftung des deutschen Volkes).

In October 2010 I successfully completed my degree in medicine.

My studies and clerkships were supported by the German National Academic Foundation (Studienstiftung des deutschen Volkes).

Publications:

Erpenbeck L and Schön MP (2010): Deadly allies: the fatal interplay between platelets and metastasizing cancer cells, *Blood* 115 (17), 3427-36

Erpenbeck L, Rubant S, Hardt K, Santoso S, Boehncke WH, Schön MP and Ludwig RJ (2010): Constitutive and functionally relevant expression of JAM-C on platelets, *Thromb Haemost* 103, 857-859

Erpenbeck L, Nieswandt B, Schön M, Pozgajova M and Schön MP (2009): Inhibition of platelet GPIb alpha and promotion of melanoma metastasis, *J Invest Dermatol* 130, 576-586

Amschler K, Schön MP, Pletz N, Wallbrecht K, **Erpenbeck L** and Schön M (2009): NF-kappaB inhibition through proteasome inhibition or IKKbeta blockade increases the susceptibility of melanoma cells to cytostatic treatment through distinct pathways, *J Invest Dermatol* 130 1073-1086